

Fréchet topologies on hypoelliptic sheaves

Thomas Krainer

ABSTRACT. In [2] it is shown that on the space of holomorphic functions convergence with respect to any ‘reasonable’ Fréchet topology implies uniform convergence on compact sets. The question is raised whether a result like this can be proved for other natural function spaces.

In this note we give an alternative proof of Krantz’s result and extend it to hypoelliptic sheaves.

1. Topologies on the space of holomorphic functions

The following theorem is proved in [2] with a method based on the Hahn-Banach theorem and Cauchy estimates for holomorphic functions. We give an alternative proof based on the closed-graph theorem. This argument lies at the heart of our extension to more general function spaces.

Theorem 1.1 ([2]). *Let $\Omega \subset \mathbb{C}^n$ be a pseudoconvex domain. Let τ be a Fréchet space topology on the space $\mathcal{H}(\Omega)$ of holomorphic functions in Ω , and assume that each continuous semi-norm with respect to τ can be majorized by a continuous semi-norm with respect to $C^\infty(\Omega)$.*

If a sequence of functions $(f_j)_j \subset \mathcal{H}(\Omega)$ converges in the topology of τ , then it converges uniformly on compact sets.

PROOF. Let $(C^\infty(\Omega), \tau_{C^\infty})$ be the Fréchet space of C^∞ -functions in Ω endowed with the topology τ_{C^∞} of uniform convergence with all derivatives on all compact sets. The space of holomorphic functions $\mathcal{H}(\Omega)$ is a closed subspace of $(C^\infty(\Omega), \tau_{C^\infty})$ by the Weierstrass approximation theorem. Thus $(\mathcal{H}(\Omega), \tau_{C^\infty})$ is a Fréchet space, and by assumption the identity map

$$(\mathcal{H}(\Omega), \tau_{C^\infty}) \rightarrow (\mathcal{H}(\Omega), \tau)$$

is continuous between Fréchet spaces, hence a topological isomorphism by the closed-graph theorem. Consequently, every sequence $(f_j)_j \subset \mathcal{H}(\Omega)$ that converges in the topology of τ converges uniformly with all partial derivatives on compact subsets. $\square$

2000 *Mathematics Subject Classification.* 46E10.

Key words and phrases. Fréchet topology, holomorphic functions, hypoellipticity.

2. Hypoelliptic sheaves

Definition 2.1. Let M be a C^∞ -manifold, and let A be a differential operator with C^∞ -coefficients acting in sections of the smooth vector bundles E and F that satisfies the following hypoellipticity condition:

- Whenever $\Omega \subset M$ is an open subset and $u \in \mathcal{D}'(\Omega, E)$ such that $Au = 0$ in Ω , then $u \in C^\infty(\Omega, E)$.

Let then $\mathcal{H}$ be the sheaf of sections of E given by

$$\mathcal{H}(\Omega, E) = \{u \in C^\infty(\Omega, E) \mid Au = 0 \text{ in } \Omega\}$$

for every open subset $\Omega \subset M$. We call $\mathcal{H}$ a hypoelliptic sheaf.

In Definition 2.1 and further below, we tacitly assume that an arbitrary density on M is chosen, and likewise a metric on the vector bundles, in order to embed functions into distributions. For the purposes of this note, the concrete choices do not matter. Hypoelliptic sheaves arise as examples of a general functional analytic theory of (nuclear) topological sheaves (of functions) as elaborated in [1] and related references.

Example 2.2. a) The sheaf of germs of holomorphic functions on a Stein manifold M is given on every open subset $\Omega \subset M$ as the kernel of the $\bar{\partial}$ -operator

$$\bar{\partial} : C^\infty(\Omega) \rightarrow C^\infty(\Omega, {}^{0,1}T^*\Omega),$$

and the $\bar{\partial}$ -operator enjoys the hypoellipticity condition of Definition 2.1.

- b) If A is elliptic, then the hypoellipticity condition of Definition 2.1 is clearly fulfilled. In particular, the sheaf of germs of harmonic forms on a Riemannian manifold (M, g) is a hypoelliptic sheaf.

We have the following natural Fréchet sheaves of germs of sections of E on M :

- (1) The sheaf $(C^\infty, \tau_{C^\infty})$ of germs of C^∞ -sections, endowed with the Fréchet topology of uniform convergence together with all derivatives on compact subsets.
- (2) The sheaf (C, τ_C) of germs of continuous sections, endowed with the Fréchet topology of uniform convergence on compact subsets.
- (3) The sheaf $(L_{\text{loc}}^p, \tau_{L_{\text{loc}}^p})$ of L_{loc}^p -sections for $1 \leq p \leq \infty$, endowed with local convergence with respect to L^p -norms.

Moreover, we have the sheaf $(\mathcal{D}', \tau_{\mathcal{D}'})$ of distributional sections of E equipped with the weak- $*$ -topology. The latter, however, is not a Fréchet sheaf. We have

$$(C^\infty, \tau_{C^\infty}) \hookrightarrow (C, \tau_C) \hookrightarrow (L_{\text{loc}}^p, \tau_{L_{\text{loc}}^p}) \hookrightarrow (L_{\text{loc}}^1, \tau_{L_{\text{loc}}^1}) \hookrightarrow (\mathcal{D}', \tau_{\mathcal{D}'})$$

with continuous embeddings.

The following proposition is an extended elaboration of Example 1.3(7) in [1] (p. 326).

Proposition 2.3. *The hypoelliptic sheaf $\mathcal{H}$ is a Fréchet subsheaf of $(C^\infty, \tau_{C^\infty})$, (C, τ_C) , as well as $(L_{\text{loc}}^p, \tau_{L_{\text{loc}}^p})$. The induced topologies τ_{C^∞} , τ_C , and $\tau_{L_{\text{loc}}^p}$ all coincide on $\mathcal{H}$.*

PROOF. Let $\Omega \subset M$ be an open subset. Since

$$A : \mathcal{D}'(\Omega, E) \rightarrow \mathcal{D}'(\Omega, F)$$

is continuous, its kernel $\mathcal{H}(\Omega, E)$ is a closed subspace with respect to the topology $\tau_{\mathcal{D}'}$. Note that, by the hypoellipticity condition, the kernel of A on $\mathcal{D}'(\Omega, E)$ is the same as the kernel of A on $C^\infty(\Omega, E)$. Consequently, $\mathcal{H}(\Omega, E)$ is a closed subspace of $C^\infty(\Omega, E)$ with respect to τ_{C^∞} , of $C(\Omega, E)$ with respect to τ_C , and of $L_{\text{loc}}^p(\Omega, E)$ with respect to $\tau_{L_{\text{loc}}^p}$ since it is the preimage of itself with respect to the continuous inclusion maps into $(\mathcal{D}'(\Omega, E), \tau_{\mathcal{D}'})$. Thus the identity maps

$$(\mathcal{H}(\Omega, E), \tau_{C^\infty}) \hookrightarrow (\mathcal{H}(\Omega, E), \tau_C) \hookrightarrow (\mathcal{H}(\Omega, E), \tau_{L_{\text{loc}}^p})$$

are continuous between Fréchet spaces, and so they are topological isomorphisms by the closed-graph theorem. $\square$

The following represents a generalization of Theorem 1.1 to the setting of hypoelliptic sheaves.

Theorem 2.4. *Let $\mathcal{H}$ be a hypoelliptic sheaf, and let $\Omega \subset M$ be an open subset. Let τ be a Fréchet topology on $\mathcal{H}(\Omega, E)$ such that every continuous semi-norm with respect to τ is majorized by a continuous seminorm with respect to τ_{C^∞} .*

Then $\tau = \tau_{C^\infty}$ on $\mathcal{H}(\Omega, E)$, and so every τ -convergent sequence $(f_j)_j \subset \mathcal{H}(\Omega, E)$ converges uniformly with all derivatives on all compact subsets of Ω .

PROOF. The proof is the same as the one of Theorem 1.1. By assumption, the identity map is continuous between the Fréchet spaces

$$(\mathcal{H}(\Omega, E), \tau_{C^\infty}) \rightarrow (\mathcal{H}(\Omega, E), \tau),$$

and thus a topological isomorphism by the closed-graph theorem. Hence $\tau = \tau_{C^\infty}$ on $\mathcal{H}(\Omega, E)$. $\square$

Remark 2.5. The proof of Theorem 2.4 makes only use of the fact that $(\mathcal{H}(\Omega, E), \tau_{C^\infty})$ is a Fréchet space. This is true for the null-spaces in C^∞ of arbitrary differential operators A with smooth coefficients and does not require hypoellipticity. Hypoelliptic sheaves $\mathcal{H}$, however, share the property with the sheaf of germs of holomorphic functions that all natural function space topologies are Fréchet on $\mathcal{H}$ and coincide with the topology τ_{C^∞} by Proposition 2.3.

References

- [1] K.-D. Bierstedt, B. Gramsch, and R. Meise, *Approximationseigenschaft, Lifting und Kohomologie bei lokalkonvexen Produktgarben*, Manuscripta Math. **19** (1976), 319–364.
- [2] S.G. Krantz, *Topologies on the space of holomorphic functions*, Preprint 2007 (arXiv:0707.1876 on arXiv.org).

PENN STATE ALTOONA, 3000 IVYSIDE PARK, ALTOONA, PA 16601, U.S.A.
E-mail address: `krainer@psu.edu`