

δ -QUASI CAUCHY SEQUENCES

HÜSEYİN ÇAKALLI

ABSTRACT. Recently, it is proved that a real-valued function defined on a subset E of $\mathbf{R}$, the set of real numbers, is uniformly continuous on E if and only if it is defined on E and preserves quasi-Cauchy sequences of points in E where a sequence is called quasi Cauchy if (Δx_n) is a null sequence. In this paper we call a real-valued function defined on a subset E of $\mathbf{R}$ δ -ward continuous if it preserves δ -quasi Cauchy sequences where a sequence $\mathbf{x} = (x_n)$ is defined to be δ -quasi Cauchy if the sequence (Δx_n) is quasi Cauchy. It turns out that δ -ward continuity implies uniform continuity, but there are uniformly continuous functions which are not δ -ward continuous. A new type compactness in terms of δ -quasi Cauchy sequences, namely δ -ward compactness is also introduced, and some theorems related to δ -ward continuity and δ -ward compactness are obtained.

1. INTRODUCTION

Following the idea given in a 1946 American Mathematical Monthly problem by Buck [4], a number of authors Posner [20], Iwinski [18], Srinivasan [24], Antoni [1], Antoni and Salat [2], Spigel and Krupnik [23] have studied A -continuity defined by a regular summability matrix A . Some authors, Öztürk [19], Savaş and Das [21], Borsik, and Salat [3] have studied A -continuity for methods of almost convergence or for related methods.

Now we give some definitions and notation which will be needed throughout the paper. $\mathbf{N}$ will denote the set of all positive integers. We will use boldface letters $\mathbf{x}$, $\mathbf{y}$, $\mathbf{z}$, ... for sequences $\mathbf{x} = (x_n)$, $\mathbf{y} = (y_n)$, $\mathbf{z} = (z_n)$, ... of terms in $\mathbf{R}$. c , δ - and Δ^2 will denote the set of all convergent sequences, the set of all quasi Cauchy sequences and the set of all δ -quasi Cauchy sequences of points in $\mathbf{R}$ where

Date: June 25, 2014.

2000 *Mathematics Subject Classification.* Primary: 40A05; Secondary: 26A15, 54C30.

Key words and phrases. Real functions, continuity, sequences.

a sequence $\mathbf{x} = (x_n)$ is called quasi Cauchy sequence if $\lim_{n \rightarrow \infty} \Delta x_n = 0$ which was given in [10] and [9] as being called forward convergent to 0 sequence (see also [5]).

Fast [16] introduced the definition of statistical convergence. Recall that for a subset M of $\mathbf{N}$ the asymptotic density of M , denoted by $\delta(M)$, is given by

$$\delta(M) = \lim_{n \rightarrow \infty} \frac{1}{n} |\{k \leq n : k \in M\}|,$$

if this limit exists, where $|\{k \leq n : k \in M\}|$ denotes the cardinality of the set $\{k \leq n : k \in M\}$. A sequence (x_n) is statistically convergent to ℓ if

$$\delta(\{n : |x_n - \ell| \geq \epsilon\}) = 0,$$

for every $\epsilon > 0$. In this case ℓ is called the statistical limit of $\mathbf{x}$. $st(\mathbf{R})$ will denote the set of statistically convergent sequences. Schoenberg [22] studied some basic properties of statistical convergence and also studied the statistical convergence as a summability method. Fridy [17] gave characterizations of statistical convergence.

A real sequence (x_k) is called lacunary statistically convergent to an element ℓ of $\mathbf{R}$ if

$$\lim_{r \rightarrow \infty} \frac{1}{h_r} |\{k \in I_r : |x_k - \ell| \geq \epsilon\}| = 0,$$

for every $\epsilon > 0$ where $I_r = (k_{r-1}, k_r]$ and $k_0 = 0$, $h_r : k_r - k_{r-1} \rightarrow \infty$ as $r \rightarrow \infty$ and $\theta = (k_r)$ is an increasing sequence of positive integers. For an introduction to lacunary statistical convergence for real or complex number sequences see [15].

Recently, Connor and Grosse-Erdman [6] have given sequential definitions of continuity for real functions calling G -continuity instead of A -continuity and their results cover the earlier works related to A -continuity where a method of sequential convergence, or briefly a method, is a linear function G defined on a linear subspace of s , denoted by c_G , into $\mathbf{R}$. A sequence $\mathbf{x} = (x_n)$ is said to be G -convergent to ℓ if $\mathbf{x} \in c_G$ and $G(\mathbf{x}) = \ell$. In particular, $\lim$ denotes the limit function $\lim \mathbf{x} = \lim_n x_n$ on the linear space c , $st - \lim$ denotes the statistical limit function $st - \lim \mathbf{x} = st - \lim_n x_n$ on the linear space $st(\mathbf{R})$ and $st_\theta - \lim$ denotes the lacunary statistical limit function $st_\theta - \lim \mathbf{x} = st_\theta - \lim_n x_n$ on the linear space $st_\theta(\mathbf{R})$. A real function f is called G -continuous at a point u provided that whenever a sequence $\mathbf{x} = (x_n)$ of terms in the domain of f is G -convergent to u , then the sequence $f(\mathbf{x}) = (f(x_n))$ is G -convergent to $f(u)$. A method G is called regular if every convergent sequence $\mathbf{x} = (x_n)$ is G -convergent with $G(\mathbf{x}) = \lim \mathbf{x}$. A method is called subsequential if

whenever $\mathbf{x}$ is G -convergent with $G(\mathbf{x}) = \ell$, then there is a subsequence (x_{n_k}) of $\mathbf{x}$ with $\lim_k x_{n_k} = \ell$.

The concept of continuity, and any concept involving continuity play very important role not only in pure mathematics but also in other branches of sciences involving mathematics especially in computer sciences and dynamical systems. It is well known that a real function f is continuous if and only if $(f(x_n))$ is convergent whenever (x_n) is, and a subset E of $\mathbf{R}$ is compact if any sequence (x_n) of points in E has a convergent subsequence whose limit is in E where $\mathbf{R}$ is the set of real numbers. A concept of ward continuity, and a concept of ward compactness have been introduced in [10] in the sense that a real function is called ward continuous if $\lim_{n \rightarrow \infty} \Delta f(x_n) = 0$ whenever $\lim_{n \rightarrow \infty} \Delta x_n = 0$, and a subset E of $\mathbf{R}$ is called ward compact if whenever $\mathbf{x} = (x_n)$ is a sequence of points in E there is a subsequence $\mathbf{z} = (z_k) = (x_{n_k})$ of $\mathbf{x}$ with $\lim_{k \rightarrow \infty} \Delta z_k = 0$ where $\Delta y_n = y_{n+1} - y_n$ (see also [5]. Recently, some further results on Quasi-Cauchy sequences are obtained [11].

The purpose of this paper is to introduce a concept of δ -ward continuity of a real function and a concept of δ -ward compactness of a subset of $\mathbf{R}$ which cannot be given by means of any G and prove some important theorems.

2. δ -WARD CONTINUITY

We say that a sequence $\mathbf{x} = (x_n)$ is δ -quasi Cauchy convergent to a number ℓ if $\lim_{k \rightarrow \infty} \Delta^2 x_k = \ell$ where $\Delta^2 x_k = x_{k+2} - 2x_{k+1} + x_k$. For the special case, $\ell = 0$, $\mathbf{x}$ is called δ -quasi Cauchy. We note that any quasi Cauchy convergent to 0 sequence is also δ -quasi Cauchy, but the converse is not always true as it can be seen by considering the sequence (n) .

Now we give the definition of δ -ward compactness of a subset of $\mathbf{R}$.

Definition 1. A subset E of $\mathbf{R}$ is called δ -ward compact if whenever $\mathbf{x} = (x_n)$ is a sequence of points in E there is a subsequence $\mathbf{z} = (z_k) = (x_{n_k})$ of $\mathbf{x}$ with $\lim_{k \rightarrow \infty} \Delta^2 z_k = 0$.

Firstly, we note that any finite subset of $\mathbf{R}$ is δ -ward compact, union of two δ -ward compact subsets of $\mathbf{R}$ is δ -ward compact and intersection of any δ -ward compact subsets of $\mathbf{R}$ is δ -ward compact. Furthermore any subset of a δ -ward

compact set is δ -ward compact and any bounded subset of $\mathbf{R}$ is δ -ward compact. Any compact subset of $\mathbf{R}$ is also δ -ward compact and the converse is not always true. For example, any open interval is δ -ward compact, which is not compact. On the other hand, the set $\mathbf{N}$ is not δ -ward compact. We note that any slowly oscillating compact subset of $\mathbf{R}$ is δ -ward compact (see also [8], and [12] for the results on slowly oscillating compactness), and any quasi-slowly oscillating compact subset of $\mathbf{R}$ is δ -ward compact where a subset E of $\mathbf{R}$ is called quasi-slowly oscillating compact (see [15]) if whenever $\mathbf{x} = (x_n)$ is a sequence of points in E , there is a quasi-slowly oscillating subsequence $\mathbf{y} = (y_{n_k})$ of $\mathbf{x}$.

We note that this definition of δ -ward compactness can not be obtained by any G -sequential compactness, i.e. by any summability matrix A , even by the summability matrix $A = (a_{nk})$ defined by $a_{nk} = 1$ if $k = n$, $a_{kn} = -2$ if $k = n + 1$ and $a_{kn} = 1$ if $k = n + 2$

$$G(x) = \lim A\mathbf{x} = \lim_{k \rightarrow \infty} \sum_{n=1}^{\infty} a_{kn} x_n = \lim_{k \rightarrow \infty} \Delta^2 x_k \quad (*)$$

(see [7] for the definition of G -sequential compactness). In this example, despite that G -sequential compact subsets of $\mathbf{R}$ should include the singleton set $\{0\}$, δ -ward compact subsets of $\mathbf{R}$ do not have to include the singleton $\{0\}$.

The concept of ward continuity suggests us to give a new type continuity, namely, δ -ward continuity, analogous to ward continuity.

Definition 2. A function f is called δ -ward continuous on E if the sequence $f(\mathbf{x}) = (f(x_n))$ is δ -quasi Cauchy whenever $\mathbf{x} = (x_n)$ is a sequence of terms in E which is δ -quasi Cauchy .

We note that this definition of continuity can not be obtained by any G -continuity, i.e. by any summability matrix A , even by the summability matrix $A = (a_{nk})$ defined by (*) however for this special summability matrix A if A -continuity of f at the point 0 implies δ -ward continuity of f , then $f(0) = 0$; and if δ -ward continuity of f implies A -continuity of f at the point 0, then $f(0) = 0$.

We also note that sum of two δ -ward continuous functions is δ -ward continuous and composite of two δ -ward continuous functions is δ -ward continuous but the product of two δ -ward continuous functions need not be δ -ward continuous as it

can be seen by considering product of the δ -ward continuous function $f(x) = x$ with itself.

Now we give the following theorem.

Theorem 1 If f is δ -ward continuous on a subset E of $\mathbf{R}$, then it is ward continuous on E .

Proof. Let (x_n) be any sequence with $\lim_{k \rightarrow \infty} \Delta x_k = 0$. Then the sequence

$$(x_1, x_1, x_2, x_2, \dots, x_n, x_n, \dots)$$

is δ -quasi Cauchy hence, by the hypothesis, the sequence

$$(f(x_1), f(x_1), f(x_2), f(x_2), \dots, f(x_n), f(x_n), \dots)$$

is δ -quasi Cauchy . It follows from this that

$$(f(x_1), f(x_2), \dots, f(x_n), \dots)$$

is quasi Cauchy convergent to 0. This completes the proof of the theorem. $\square$

The converse of this theorem is not always true for the ward continuous function $f(x) = \sin x$ is an example which is uniformly continuous but is δ -ward continuous. This example also explains that not all uniformly continuous functions are δ -ward continuous.

Corollary 2 If f is δ -ward continuous on a subset E of $\mathbf{R}$, then it is continuous on E in the ordinary sense.

Proof. Although, the proof follows from Theorem 1 above and Theorem 7 in [10] with the statement that any ward continuous function is continuous, we give a different proof in the following for completeness.

Let (x_n) be any convergent sequence with $\lim_{k \rightarrow \infty} x_k = \ell$. Then the sequence $(x_1, \ell, x_2, \ell, \dots, \ell, x_n, \ell, \dots)$ also converges to ℓ , hence $\lim_{k \rightarrow \infty} \Delta x_k = 0$. So the sequence $(x_1, \ell, x_2, \ell, \dots, \ell, x_n, \ell, \dots)$ is δ -quasi Cauchy hence, by the hypothesis, the sequence

$$(f(x_1), f(\ell), f(x_2), f(\ell), \dots, f(\ell), f(x_n), f(\ell), \dots)$$

is δ -quasi Cauchy . It follows from this that

$$(f(x_1), f(x_2), \dots, f(x_n), \dots)$$

converges to $f(\ell)$. This completes the proof of the theorem.

The converse is not always true for the function $f(x) = x^2$ is an example since the sequence (n) is δ -quasi Cauchy while $(f(n)) = (n^2)$ is not δ -quasi Cauchy . $\square$

Now we have the following result.

Corollary 3 If f is δ -ward continuous, then it is statistically continuous.

We state much more general case in the following.

Corollary 4 If f is δ -ward continuous, then it is G -continuous for any regular sub-sequential method G .

Theorem 5 δ -ward continuous image of any δ -ward compact subset of $\mathbf{R}$ is δ -ward compact.

Proof. Let f be a δ -ward continuous function and E be a δ -ward compact subset of $\mathbf{R}$. Take any sequence $\mathbf{y} = (y_n)$ of terms in $f(E)$. Write $y_n = f(x_n)$ where $x_n \in E$ for each $n \in \mathbf{N}$. δ -ward compactness of E implies that there is a subsequence $\mathbf{z} = (z_k) = (x_{n_k})$ of $\mathbf{x}$ with $\lim_{k \rightarrow \infty} \Delta^2 z_k = 0$. Since f is δ -ward continuous, $(t_k) = f(\mathbf{z}) = (f(z_k))$ is δ -quasi Cauchy . Thus (t_k) is a subsequence of the sequence $f(\mathbf{x})$ with $\lim_{k \rightarrow \infty} \Delta t_k = 0$. This completes the proof of the theorem. $\square$

Corollary 6 δ -ward continuous image of any compact subset of $\mathbf{R}$ is compact.

The proof of this theorem follows from the preceding theorem.

It is well-known that any continuous function on a compact subset E of $\mathbf{R}$ is uniformly continuous on E . It is also true for a regular subsequential method G that any δ -ward continuous function on a G -sequentially compact subset E of $\mathbf{R}$ is uniformly continuous on E (see [7]). For δ -ward continuous functions we have the following.

Theorem 7 If f is δ -ward continuous on a subset E of $\mathbf{R}$, then it is uniformly continuous on E .

Proof. The proof of this theorem follows from Theorem 2 in [11]. $\square$

It is known that uniform limit of a sequence of continuous functions is continuous. This is also true in case of δ -ward continuity.

Theorem 8 If (f_n) is a sequence of δ -ward continuous functions defined on a subset E of $\mathbf{R}$ and (f_n) is uniformly convergent to a function f , then f is δ -quasi Cauchy continuous on E .

Proof. Let $\mathbf{x} = (x_n)$ be any δ -quasi Cauchy sequence of points in E and $\varepsilon > 0$. Then there exists a positive integer N such that $|f_n(x) - f(x)| < \frac{\varepsilon}{8}$ for all $x \in E$ whenever $n \geq N$. As f_N is δ -quasi Cauchy continuous, there exists a positive integer N_1 , depending on ε and greater than N such that

$$|f_N(x_{n+2}) - 2f_N(x_{n+1}) + f_N(x_n)| < \frac{\varepsilon}{8}$$

for $n \geq N_1$. Now for $n \geq N_1$ we have

$$\begin{aligned} |f(x_{n+2}) - 2f(x_{n+1}) + f(x_n)| &= \\ &\leq |f(x_{n+2}) - f_N(x_{n+2})| + 2|f_N(x_{n+1}) - f(x_{n+1})| + |f_N(x_n) - f(x_n)| \\ &\quad + |f_N(x_{n+2}) - 2f_N(x_{n+1}) + f_N(x_n)| \\ &\leq \frac{\varepsilon}{8} + \frac{\varepsilon}{4} + \frac{\varepsilon}{8} + \frac{\varepsilon}{8} < \varepsilon. \end{aligned}$$

This completes the proof of the theorem. $\square$

Theorem 9 The set of all δ -ward continuous functions on E is a closed subset of the set of all continuous functions on E , i.e. $\overline{C(E)} = C(E)$ where $C(E)$ is the set of all continuous functions on E , $\overline{C(E)}$ denotes the set of all cluster points of $C(E)$ and E is a bounded subset of $\mathbf{R}$.

Proof. Let us denote the set of all δ -ward continuous functions on E by $SFC(E)$ and f be any element in the closure of $SFC(E)$. Then there exists a sequence of points in $SFC(E)$ such that $\lim_{k \rightarrow \infty} f_k = f$. To show that f is δ -ward continuous take any δ -quasi Cauchy sequence (x_n) . Let $\varepsilon > 0$. Since (f_k) converges to f , there exists an N such that for all $x \in E$ and for all $n \geq N$, $|f(x) - f_n(x)| < \frac{\varepsilon}{8}$. As f_N is δ -ward continuous, there is an N_1 , greater than N , such that for all $n \geq N_1$, $|f_N(x_{n+2}) - 2f_N(x_{n+1}) + f_N(x_n)| < \frac{\varepsilon}{4}$. Hence for all $n \geq N_1$,

$$\begin{aligned} &|f(x_{n+2}) - 2f(x_{n+1}) + f(x_n)| \\ &\leq |f(x_{n+2}) - f_N(x_{n+2})| + 2|f_N(x_{n+1}) - f(x_{n+1})| + |f(x_n) - f_N(x_n)| \\ &\quad + |f_N(x_{n+2}) - 2f_N(x_{n+1}) + f_N(x_n)| \\ &\leq \frac{\varepsilon}{8} + \frac{\varepsilon}{4} + \frac{\varepsilon}{8} + \frac{\varepsilon}{8} < \varepsilon. \end{aligned}$$

This completes the proof of the theorem. $\square$

Corollary 10 The set of all δ -ward continuous functions on a subset E of $\mathbf{R}$ is a complete subspace of the space of all continuous functions on E .

Proof. The proof follows from the preceding theorem. □

Finally we note the following further investigation problems arise.

Problem 1. For further study, we suggest to investigate δ -quasi Cauchy sequences of fuzzy points and δ -ward continuity for the fuzzy functions. However due to the change in settings, the definitions and methods of proofs will not always be analogous to those of the present work (for example see [13]).

Problem 2. For another further study we suggest to introduce a new concept in dynamical systems using δ -ward continuity.

REFERENCES

- [1] J.Antoni, On the A -continuity of real functions II, Math. Slovaca, **36**, No.3, (1986), 283-287. MR **88a**:26001
- [2] J.Antoni and T.Salat, On the A -continuity of real functions, Acta Math. Univ. Comenian. **39**, (1980), 159-164. MR **82h**:26004
- [3] J.Borsik and T.Salat, On F -continuity of real functions, Tatra Mt. Math. Publ. **2**, (1993), 37-42. MR **94m**:26006
- [4] R.C.Buck, Solution of problem 4216 Amer. Math. Monthly **55**, (1948), 36. MR **15**: 26874
- [5] D.Burton, and J. Coleman, Quasi-Cauchy Sequences, Amer. Math. Monthly, **117**, 4,328-333, 2010.
- [6] J.Connor, Grosse-Erdmann K.-G. Sequential definitions of continuity for real functions, Rocky Mountain J. Math., **33**, (1), (2003), 93-121. MR **2004e**:26004
- [7] H.Çakallı, Sequential definitions of compactness, Applied Mathematics Letters, 21 , No:6, (2008), 594-598,
- [8], Slowly oscillating continuity, Abstract and Applied Analysis, Volume 2008, Article ID 485706,
- [9], Forward compactness, Conference on Summability and Applications, Shawnee State University, November 6, 2009 November 8, 2009. <http://webpages.math.luc.edu/mgb/ShawneeConference/Articles/HuseyinCakalliOhio.pdf> .
- [10], Forward continuity, Journal of Computational Analysis and Applications, to appear.
- [11], On quasi-Cauchy sequences, MathFest 2010, Pittsburg, USA.
- [12] H.Çakallı, I.Canak and M.Dik, δ -quasi-Slowly oscillating continuity, Applied Mathematics and Computation, **216**, 2010, 2865-2867.
- [13] Hüseyin Çakallı and Pratulananda Das, Fuzzy compactness via summability, Applied Mathematics Letters, **22**, No: 11, November 2009, 1665-1669.
- [14] I.Canak, U. Totur, M. Dik, One-sided Tauberian conditions for (A,k) summability method, Mathematical and Computer Modelling, to appear.
- [15] I.Canak and M.Dik, New types of continuities, Abstract and Applied Analysis, 2010.
- [16] H.Fast, Sur la convergence statistique, Colloq. Math. **2**, (1951), 241-244. MR **14**:29c
- [17] J.A.Fridy, On statistical convergence, Analysis, **5**, (1985), 301-313 . MR **87b**:40001
- [18] T.B.Iwinski, Some remarks on Toeplitz methods and continuity, Comment.Math. Prace Mat. **17**, (1972), 37-43. MR **48**759
- [19] E.Öztürk, On almost-continuity and almost A -continuity of real functions, Comm.Fac.Sci. Univ.Ankara Ser. A1 Math. Stat. **32**, (1983), 25-30. MR **86h**:26003
- [20] E.C.Posner, Summability preserving functions, Proc.Amer.Math.Soc. **12**, (1961), 73-76. MR **22**12327
- [21] E. Savaş and G. Das, On the A -continuity of real functions, İstanbul Univ. Fen Fak. Mat Derg. **53**, (1994), 61-66. MR **97m**:26004

- [22] I.J.Schoenberg, The integrability of certain functions and related summability methods, Amer. Math. Monthly **66**, (1959), 361-375. MR **21**:3696
- [23] E. Spigel and N. Krupnik, On the A -continuity of real functions, J.Anal. **2**, (1994), 145-155. MR **95h**:26004
- [24] V.K.Srinivasan, An equivalent condition for the continuity of a function, Texas J. Sci. **32**, (1980), 176-177. MR **81f**:26001

HÜSEYİN ÇAKALLI, DEPARTMENT OF MATHEMATICS, MALTEPE UNIVERSITY, MARMARA EĞİTİM
KÖYÜ, TR 34857, MALTEPE, İSTANBUL-TURKEY PHONE:(+90216)6261050 EXT:1206,
FAX:(+90216)6261113

E-mail address: hcakalli@maltepe.edu.tr; hcakalli@gmail.com