

ON SOME CHARACTERIZATIONS OF RULED SURFACE OF A CLOSED SPACELIKE CURVE WITH TIMELIKE BINORMAL IN DUAL LORENTZIAN SPACE

Özcan BEKTAŞ *

Süleyman ŞENYURT *

Abstract

In this paper, we investigate the relations between the pitch, the angle of pitch and drall of parallel ruled surface of a closed spacelike curve with timelike binormal in dual Lorentzian space.

Keywords: Spacelike dual curve; ruled surface; Lorentzian space; dual numbers.

2000 AMS Subject Classification: 53A04, 45F10

1 Introduction

Dual numbers were introduced by W.K. Clifford (1849-79) as a tool for his geometrical investigations. After him E.Study used dual numbers and dual vectors in his research on the geometry of lines and kinematics [4]. The pitches and the angles of the pitches of the closed ruled surfaces corresponds to the one parameter dual unit spherical curves in space of lines IR^3 were calculated respectively by Hacısalihoğlu [7] and Gursoy [5]. Definition of the parallel ruled surface were presented by Blaschke (translated by Erim [3]). The integral invariants of the parallel ruled surfaces in the 3-dimensional Euclidean space E^3 corresponding to the unit dual spherical parallel curves were calculated by Senyurt [11]. The set is called dual numbers set [2].

On this set product and addition operations are respectively

$$(\lambda + \varepsilon\lambda^*) + (\beta + \varepsilon\beta^*) = (\lambda + \beta) + \varepsilon(\lambda^* + \beta^*)$$

and

$$(\lambda + \varepsilon\lambda^*)(\beta + \varepsilon\beta^*) = \lambda\beta + \varepsilon(\lambda\beta^* + \lambda^*\beta).$$

$ID^3 = \left\{ \vec{A} = \vec{a} + \varepsilon\vec{a}^* \mid \vec{a}, \vec{a}^* \in IR^3 \right\}$ the elements of ID^3 are called dual vectors. On this set addition and scalar product operations are respectively

*Ordu University, Faculty of Art and Science, Department of Mathematics, 52750, Perşembe, Ordu, Turkey, ozcanbektas1986@hotmail.com, senyurtsuleyman@hotmail.com.

$$\begin{aligned} \oplus : ID^3 \times ID^3 &\rightarrow ID^3 \\ (\vec{A}, \vec{B}) &\rightarrow \vec{A} \oplus \vec{B} = \vec{a} + \vec{b} + \varepsilon (\vec{a}^* + \vec{b}^*) \end{aligned}$$

$$\begin{aligned} \odot : ID \times ID^3 &\rightarrow ID^3 \\ (\lambda, \vec{A}) &\rightarrow \lambda \odot \vec{A} = (\lambda + \varepsilon \lambda^*) \odot (\vec{a} + \varepsilon \vec{a}^*) = \lambda \vec{a} + \varepsilon (\lambda \vec{a}^* + \lambda^* \vec{a}) \end{aligned}$$

The set $(ID^3, \oplus)$ is a module over the ring $(ID, +, \cdot)$. (*ID-Modul*).

The Lorentzian inner product of dual vectors $\vec{A}, \vec{B} \in ID^3$ is defined by

$$\langle \vec{A}, \vec{B} \rangle = \langle \vec{a}, \vec{b} \rangle + \varepsilon (\langle \vec{a}, \vec{b}^* \rangle + \langle \vec{a}^*, \vec{b} \rangle)$$

with the Lorentzian inner product $\vec{a} = (a_1, a_2, a_3)$ and $\vec{b} = (b_1, b_2, b_3) \in IR^3$

$$\langle \vec{a}, \vec{b} \rangle = -a_1 b_1 + a_2 b_2 + a_3 b_3.$$

Therefore, ID^3 with the Lorentzian inner product $\langle \vec{A}, \vec{B} \rangle$ is called 3-dimensional dual Lorentzian space and denoted by of $ID_1^3 = \{ \vec{A} = \vec{a} + \varepsilon \vec{a}^* \mid \vec{a}, \vec{a}^* \in IR_1^3 \}$ [14].

A dual vector $\vec{A} = \vec{a} + \varepsilon \vec{a}^* \in ID_1^3$ is called A dual space-like vector if $\langle \vec{A}, \vec{A} \rangle > 0$ or $\vec{A} = 0$, A dual time-like vector if $\langle \vec{A}, \vec{A} \rangle < 0$, A dual null(light-like) vector if $\langle \vec{A}, \vec{A} \rangle = 0$ for $\vec{A} \neq 0$. For $\vec{A} \neq 0$, the norm $\|\vec{A}\|$ of $\vec{A} = \vec{a} + \varepsilon \vec{a}^* \in ID^3$ is defined by

$$\|\vec{A}\| = \sqrt{|\langle \vec{A}, \vec{A} \rangle|} = \|\vec{a}\| + \varepsilon \frac{\langle \vec{a}, \vec{a}^* \rangle}{\|\vec{a}\|}, \quad \|\vec{a}\| \neq 0.$$

The dual Lorentzian cross-product of $\vec{A}, \vec{B} \in ID^3$ is defined as

$$\vec{A} \wedge \vec{B} = \vec{a} \wedge \vec{b} + \varepsilon (\vec{a} \wedge \vec{b}^* + \vec{a}^* \wedge \vec{b})$$

with the Lorentzian cross-product $\vec{a}, \vec{b} \in IR^3$

$$\vec{a} \wedge \vec{b} = (a_3 b_2 - a_2 b_3, a_1 b_3 - a_3 b_1, a_1 b_2 - a_2 b_1) [14].$$

1.1. Theorem(E. Study): The oriented lines in are in one to one correspondence with the points of the dual unit sphere where ID-Modul, see [9].

Dual number $\Phi = \phi + \varepsilon \phi^*$ is called dual angle between $\vec{A}$ ve $\vec{B}$ unit dual vectors. In this place

$$\sinh(\phi + \varepsilon \phi^*) = \sinh \phi + \varepsilon \phi^* \cosh \phi \quad \text{and} \quad \cosh(\phi + \varepsilon \phi^*) = \cosh \phi + \varepsilon \phi^* \sinh \phi.$$

2 ON SOME CHARACTERIZATIONS OF RULED SURFACE OF A CLOSED SPACELIKE CURVE IN DUAL LORENTZIAN SPACE (ID_1^3)

$\vec{U} = \vec{U}_1(t)$, $\|\vec{U}(t)\| = 1$ is a differentiable spacelike curve in the one parameter dual unit spherical motion K/K' . The closed ruled surface ($\vec{U}$) corresponds to this curve in IR^3 .

Let the dual orthonormal system of curve $\vec{U} = \vec{U}_1(t)$ as

$$\vec{U}_1 = \vec{U}(t) \quad , \quad \vec{U}_2 = \vec{U}'(t) / \|\vec{U}(t)\| \quad , \quad \vec{U}_3 = \vec{U}_1 \wedge \vec{U}_2$$

2.1. Definition: The closed ruled surface ($\vec{U}$) corresponds to the dual spacelike curve $\vec{U}(t)$ which makes the fixed dual angle $\Phi = \phi + \varepsilon\phi^*$ with $\vec{U}(t)$ and defines by

$$(2.1) \quad \vec{V} = \cosh \Phi \vec{U}_1 + \sinh \Phi \vec{U}_3$$

This surface ($\vec{V}$) corresponds to dual spacelike vector $\vec{V}$ is called the parallel ruled surface of surface ($\vec{U}$) in dual Lorentzian space ID_1^3 .

Let $\vec{U}(t)$ be a closed spacelike curve with curvature $\kappa = k_1 + \varepsilon k_1^*$ and torsion $\tau = k_2 + \varepsilon k_2^*$. Let Frenet frames of $\vec{U}(t)$ be $\{\vec{U}_1, \vec{U}_2, \vec{U}_3\}$. In this trihedron, $\vec{U}_1$ and $\vec{U}_2$ are spacelike vectors, $\vec{U}_3$ is timelike vector. For this vectors, we can write

$$(2.2) \quad \vec{U}_1 \wedge \vec{U}_2 = \vec{U}_3 \quad , \quad \vec{U}_2 \wedge \vec{U}_3 = -\vec{U}_1 \quad , \quad \vec{U}_3 \wedge \vec{U}_1 = -\vec{U}_2$$

where $\wedge$ is the Lorentzian cross product, in space ID_1^3 . In this situation, the Frenet formulas are given by

$$(2.3) \quad \vec{U}_1' = \kappa \vec{U}_2 \quad , \quad \vec{U}_2' = -\kappa \vec{U}_1 + \tau \vec{U}_3 \quad , \quad \vec{U}_3' = \tau \vec{U}_2, [15]$$

The Frenet instantaneous rotation vector for the timelike curve is given by

$$(2.4) \quad \vec{\Psi} = \tau \vec{U}_1 - \kappa \vec{U}_3, [13]$$

Let be $\vec{V}_1 = \vec{V}$. Differentiating of the vector $\vec{V}_1$ with respect the parameter and using the equation (2.3) we get

$$(2.5) \quad \vec{V}_1' = (\kappa \cosh \Phi + \tau \sinh \Phi) \vec{U}_2$$

If the norm of the vector denotes by P , we get

$$(2.6) \quad P = (\kappa \cosh \Phi + \tau \sinh \Phi)$$

Then if is known that, substituting by the values of the equations (2.5) and (2.6) into, we get

$$(2.7) \quad \vec{V}_2 = \vec{U}_2$$

Then, for the vector, we get

$$(2.8) \quad \vec{V}_3 = \sinh \Phi \vec{U}_1 + \cosh \Phi \vec{U}_3$$

If the equation (2.1) , (2.7) and (2.8) are written matrix form, we have

$$(2.9) \quad \begin{bmatrix} \vec{V}_1 \\ \vec{V}_2 \\ \vec{V}_3 \end{bmatrix} = \begin{bmatrix} \cosh \Phi & 0 & \sinh \Phi \\ 0 & 1 & 0 \\ \sinh \Phi & 0 & \cosh \Phi \end{bmatrix} \cdot \begin{bmatrix} \vec{U}_1 \\ \vec{U}_2 \\ \vec{U}_3 \end{bmatrix}$$

or

$$(2.10) \quad \begin{bmatrix} \vec{U}_1 \\ \vec{U}_2 \\ \vec{U}_3 \end{bmatrix} = \begin{bmatrix} \cosh \Phi & 0 & -\sinh \Phi \\ 0 & 1 & 0 \\ -\sinh \Phi & 0 & \cosh \Phi \end{bmatrix} \cdot \begin{bmatrix} \vec{V}_1 \\ \vec{V}_2 \\ \vec{V}_3 \end{bmatrix}$$

If the equation (2.10) is separated into real and dual parts, we get

$$(2.11) \quad \begin{cases} \vec{u}_1 = \cosh \varphi \vec{v}_1 - \sinh \varphi \vec{v}_3 \\ \vec{u}_2 = \vec{v}_2 \\ \vec{u}_3 = -\sinh \varphi \vec{v}_1 + \cosh \varphi \vec{v}_3 \\ \vec{u}_1^* = \cosh \varphi \vec{v}_1^* - \sinh \varphi \vec{v}_3^* + \varphi^* (\sinh \varphi \vec{v}_1 - \cosh \varphi \vec{v}_3) \\ \vec{u}_2^* = \vec{v}_2^* \\ \vec{u}_3^* = -\sinh \varphi \vec{v}_1^* + \cosh \varphi \vec{v}_3^* - \varphi^* (\cosh \varphi \vec{v}_1 - \sinh \varphi \vec{v}_3) \end{cases}$$

Let be curvature $P = p + \varepsilon p^*$ and torsion $Q = q + \varepsilon q^*$ of curve $\vec{V}(t)$. Between the vectors $\vec{V}_1, \vec{V}_2, \vec{V}_3$ and the derivate vectors $\vec{V}_1', \vec{V}_2', \vec{V}_3'$ there is following relation

$$(2.12) \quad \begin{cases} \vec{V}_1' = P \vec{V}_2', \quad \vec{V}_1' = -P \vec{V}_1 + Q \vec{V}_3, \quad \vec{V}_3' = Q \vec{V}_2 \\ P = \sqrt{\langle \vec{V}_1', \vec{V}_1' \rangle}, \quad Q = \frac{\det(\vec{V}_1, \vec{V}_1', \vec{V}_1'')}{\langle \vec{V}_1', \vec{V}_1' \rangle}. \end{cases}$$

If the equation (2.12) is separated into the real and dual parts, we can write

$$(2.13) \quad \begin{cases} \vec{v}_1' = p \vec{v}_2, \quad \vec{v}_2' = -p \vec{v}_1 + q \vec{v}_3, \quad \vec{v}_3' = q \vec{v}_2 \\ \vec{v}_1'^* = p \vec{v}_2^* + p^* \vec{v}_2, \\ \vec{v}_2'^* = -p \vec{v}_1^* - p^* \vec{v}_1 + q^* \vec{v}_3, \\ \vec{v}_3'^* = q \vec{v}_2^* + q^* \vec{v}_2 \end{cases}$$

Now, we can calculate the value of Q relative to κ and τ . Derivating the equation (2.5) with respect to the parameter an of making the recesiory operations we may write

$$(2.14) \quad \begin{aligned} \vec{v}_1'' = & (-\kappa^2 \cosh \Phi - \kappa \tau \sinh \Phi) \vec{U}_1 + \\ & + (\kappa \cosh \Phi + \tau \sinh \Phi)' \vec{U}_2 + (\kappa \tau \cosh \Phi + \tau^2 \sinh \Phi) \vec{U}_3 \end{aligned}$$

Substituting by the equations (2.1) , (2.5) and (2.14) into the equation (2.12) , we get

$$(2.15) \quad Q = \kappa \sinh \Phi + \tau \cosh \Phi$$

The equations (2.6) and (2.15) are separated into the dual and real parts, we have

$$(2.16) \quad \begin{cases} p = k_1 \cosh \varphi + k_2 \sinh \varphi \\ p^* = k_1^* \cosh \varphi + k_2^* \sinh \varphi + \varphi^* (k_1 \sinh \varphi + k_2 \cosh \varphi) \\ q = k_1 \sinh \varphi + k_2 \cosh \varphi \\ q^* = k_1^* \sinh \varphi + k_2^* \cosh \varphi + \varphi^* (k_1 \cosh \varphi + k_2 \sinh \varphi) \end{cases}$$

In the dual unit spherical motion K/K' , the dual orthonormal system $\{\vec{V}_1, \vec{V}_2, \vec{V}_3\}$ each time t is make a dual rotation motion around the instantaneous dual Pfaffion vector. This vector is determined the following equation

$$(2.17) \quad \vec{\Psi} = Q \vec{V}_1 - P \vec{V}_3, [13]$$

Fort he Steiner vector of the motion, we can write

$$(2.18) \quad \vec{D} = \oint \vec{\Psi}$$

or

$$(2.19) \quad \vec{D} = \vec{V}_1 \oint Q dt - \vec{V}_3 \oint P dt$$

Setting by the values of the vectors $\vec{U}_1$ and $\vec{U}_3$ as the equations (2.11) into the equations (2.4), necessary operations have been done, we get

$$\vec{\Psi} = Q \vec{V}_1 - P \vec{V}_3.$$

In this case, we consider the last equation and equation (2.17), we can write

$$(2.20) \quad \vec{\Psi} = \vec{\Psi}$$

Because of the equations $\vec{D} = \oint \vec{\Psi}$ and $\vec{D} = \oint \vec{\Psi}$ we can obtain $\vec{D} = \vec{D}$. Then, for the dual Steiner vector of the motion, we may write

$$(2.21) \quad \vec{D} = \vec{V}_1 \oint Q dt - \vec{V}_3 \oint P dt$$

It is separated into the dual and real part as

$$(2.22) \quad \begin{cases} \vec{d} = \vec{v}_1 \oint q dt - \vec{v}_3 \oint p dt, \\ \vec{d}^* = \vec{v}_1 \oint q^* dt + \vec{v}_1^* \oint q dt + \vec{v}_1^* \oint q dt - \vec{v}_3^* \oint p dt \end{cases}$$

$$(2.23) \quad \vec{D} = \vec{U}_1 \oint \tau dt - \vec{U}_3 \oint \kappa dt$$

The equation (2.23) can be written type of the dual and real part as follow

$$(2.24) \quad \begin{cases} \vec{d} = \vec{u}_1 \oint k_2 dt - \vec{u}_3 \oint k_1 dt, \\ \vec{d}^* = \vec{u}_1^* \oint k_2 dt + \vec{u}_1 \oint k_2^* dt - \vec{u}_3^* \oint k_1 dt - \vec{u}_3 \oint k_1^* dt \end{cases}$$

If the equation (2.3) is separated into the dual and real part, we can obtain

$$(2.25) \quad \begin{cases} \vec{u}_1' = k_1 \vec{u}_2, \quad \vec{u}_2' = -k_1 \vec{u}_1 + k_2 \vec{u}_3, \quad \vec{u}_3' = k_2 \vec{u}_2 \\ \vec{u}_1'^* = k_1^* \vec{u}_2 + k_1 \vec{u}_2^*, \\ \vec{u}_2'^* = -k_1^* + k_2^* \vec{u}_3 - k_1 \vec{u}_1^* + k_2 \vec{u}_3^* \\ \vec{u}_3'^* = k_2^* \vec{u}_2 + k_2 \vec{u}_2^* \end{cases}$$

Now, let is calculate the integral invariants of the closed ruled surfaces respectively. The pitch of the closed surface $\vec{C}$ is obtained as

$$L_{u_1} = \langle \vec{d}, \vec{u}_1^* \rangle + \langle \vec{d}^*, \vec{u}_1 \rangle,$$

$$(2.26) \quad L_{u_1} = \oint k_2^* dt.$$

For the dual angle of the pitch of the closed surface , we may write

$$\Lambda_{U_1} = -\langle \vec{D}, \vec{U}_1 \rangle$$

Because of the equation (2.23) we can obtain

$$(2.27) \quad \Lambda_{U_1} = -\oint \tau dt.$$

If the equation (2.27) is separated into the dual and real part, we can obtain

$$(2.28) \quad \lambda_{u_1} = -\oint k_2 dt \quad , \quad L_{u_1} = \oint k_2^* dt$$

For the drall of the closed surface , we may write

$$P_{U_1} = \frac{\langle d\vec{u}_1, d\vec{u}_1^* \rangle}{\langle d\vec{u}_1, d\vec{u}_1 \rangle}$$

Setting by the values of the statements $\vec{d\bar{u}}_1$ and $\vec{d\bar{u}}_1^*$ as the equations (2.25) into the last equations, we get

$$(2.29) \quad P_{U_1} = \frac{k_1^*}{k_1}$$

The pitch of the closed surface is obtained as

$$L_{u_2} = \langle \vec{d}, \vec{u}_2^* \rangle + \langle \vec{d}^*, \vec{u}_2 \rangle,$$

$$(2.30) \quad L_{u_2} = 0.$$

For the dual angle of the pitch of the closed surface , we may write

$$\Lambda_{U_2} = -\langle \vec{D}, \vec{U}_2 \rangle,$$

$$(2.31) \quad \Lambda_{U_2} = 0.$$

For the drall of the closed surface , we may write

$$P_{U_2} = \frac{\langle \vec{d\bar{u}}_2, \vec{d\bar{u}}_2^* \rangle}{\langle \vec{d\bar{u}}_2, \vec{d\bar{u}}_2 \rangle}$$

Setting by the values of the statements $\vec{d\bar{u}}_2$ and $\vec{d\bar{u}}_2^*$ as the equations (2.25) into the last equations, we get

$$(2.32) \quad P_{U_2} = \frac{k_1 k_1^* - k_2 k_2^*}{k_1^2 - k_2^2}$$

The pitch of the closed surface is obtained as

$$L_{u_3} = \langle \vec{d}, \vec{u}_3^* \rangle + \langle \vec{d}^*, \vec{u}_3 \rangle,$$

$$(2.33) \quad L_{u_3} = \oint k_1^* dt$$

For the dual angle of the pitch of the closed surface , we may write

$$\Lambda_{U_3} = -\langle \vec{D}, \vec{U}_3 \rangle$$

Because of the equation (2.23) we can obtain

$$(2.34) \quad \Lambda_{U_3} = -\oint \kappa dt$$

If the equation (2.34) is separated into the dual and real part, we can obtain

$$(2.35) \quad \lambda_{u_3} = - \oint k_1 dt \quad , \quad L_{u_3} = \oint k_1^* dt$$

For the drall of the closed surface , we may write

$$P_{U_3} = \frac{\langle d\vec{u}_3, d\vec{u}_3^* \rangle}{\langle d\vec{u}_3, d\vec{u}_3 \rangle}$$

Setting by the values of the statements $d\vec{u}_3$ and $d\vec{u}_3^*$ as the equations (2.25) into the last equations, we get

$$(2.36) \quad P_{U_3} = \frac{k_2^*}{k_2}.$$

Let $\Omega(t) = \omega(t) + \varepsilon\omega^*(t)$ be Lorentzian timelike angle of between the instantaneous dual Pfaffion vector $\vec{\Psi}$ and the vector $\vec{U}_3$ [10].

a) If the instantaneous dual Pfaffion vector $\vec{\Psi}$ is spacelike ($|\tau| > |\kappa|$)

$$\kappa = \|\vec{\Psi}\| \sinh \Omega \quad , \quad \tau = \|\vec{\Psi}\| \cosh \Omega$$

On the way $\vec{C} = \vec{c} + \varepsilon \vec{c}^*$, unit vector about the vector $\vec{\Psi}$ direction is

$$(2.37) \quad \vec{C} = \cosh \Omega \vec{U}_1 - \sinh \Omega \vec{U}_3$$

If the equation (2.37) is separated into the dual and real part, we can obtain

$$(2.38) \quad \begin{cases} \vec{c} = \cosh \omega \vec{u}_1 - \sinh \omega \vec{u}_3 \\ \vec{c}^* = \cosh \omega \vec{u}_1^* - \sinh \omega \vec{u}_3^* + \omega^* \sinh \omega \vec{u}_1 - \omega^* \cosh \omega \vec{u}_3 \end{cases}$$

The pitch of the closed surface $\vec{C}$ is obtained as

$$(2.39) \quad L_C = \cosh \omega \oint k_2^* dt - \sinh \omega \oint k_1^* dt - \omega^* (\cosh \omega \oint k_1 dt - \sinh \omega \oint k_2 dt)$$

$$(2.40) \quad \begin{aligned} L_C = & \cosh \omega \oint k_2^* dt - \sinh \omega \oint k_1^* dt - \\ & - \omega^* \left(\cosh \omega \oint k_1 dt - \sinh \omega \oint k_2 dt \right) \end{aligned}$$

If we use the equations (2.28) and (2.35) into the equation (2.39) we get

$$(2.41) \quad L_C = \cosh \omega L_{u_1} - \sinh \omega L_{u_3} - \omega^* (\sinh \omega \lambda_{u_1} - \cosh \omega \lambda_{u_3})$$

For the dual angle of the pitch of the closed ruled surface , we may write

$$\Lambda_C = -\langle \vec{D}, \vec{C} \rangle$$

Because of the equations (2.23) and (2.37) we can obtain

$$(2.42) \quad \Lambda_C = -\cosh \Omega \oint \tau dt + \sinh \Omega \oint \kappa dt$$

If we use the equations (2.27) and (2.34) into the last equations, we get

$$(2.43) \quad \Lambda_C = \cosh \Omega \Lambda_{U_1} - \sinh \Omega \Lambda_{U_3}$$

For the drall of the closed surface , we may write

$$(2.44) \quad P_C = \frac{\langle d\vec{C}, d\vec{C}^* \rangle}{\langle d\vec{C}, d\vec{C} \rangle}$$

$$P_C = \frac{(k_1 \cosh \omega - k_2 \sinh \omega) [(k_1^* - k_2 \omega^*) \cosh \omega + (k_1 \omega^* - k_2^*) \sinh \omega] - \omega' \omega^{*'}}{(k_1 \cosh \omega - k_2 \sinh \omega)^2 - \omega'^2}$$

b) If the instantaneous dual Pfaffion vector $\vec{\Psi}$ is timelike ($|\tau| < |\kappa|$)

$$\kappa = \|\vec{\Psi}\| \cosh \Omega \quad , \quad \tau = \|\vec{\Psi}\| \sinh \Omega$$

On the way $\vec{C} = \vec{C} + \varepsilon \vec{C}^*$, unit vector about the vector $\vec{\Psi}$ direction is

$$(2.45) \quad \vec{C} = \sinh \Omega \vec{U}_1 - \cosh \Omega \vec{U}_3$$

If the equation (2.45) is separated into the dual and real part, we can obtain

$$(2.46) \quad \begin{cases} \vec{C} = \sinh \omega \vec{u}_1 - \cosh \omega \vec{u}_3 \\ \vec{C}^* = \sinh \omega \vec{u}_1^* - \cosh \omega \vec{u}_3^* + \omega^* \cosh \omega \vec{u}_1 - \omega^* \sinh \omega \vec{u}_3 \end{cases}$$

The pitch of the closed surface is obtained as

$$(2.47) \quad L_C = \langle \vec{d}, \vec{C}^* \rangle + \langle \vec{d}^*, \vec{C} \rangle$$

$$L_C = \sinh \omega \oint k_2^* dt - \cosh \omega \oint k_1^* dt -$$

$$- \omega^* \left(\sinh \omega \oint k_1 dt - \cosh \omega \oint k_2 dt \right)$$

or

$$(2.48) \quad L_C = \sinh \omega L_{u_1} - \cosh \omega L_{u_3} - \omega^* (\cosh \omega \lambda_{u_1} - \sinh \omega \lambda_{u_3})$$

For the dual angle of the pitch of the closed ruled surface , we may write

$$\Lambda_C = - \left\langle \vec{D}, \vec{C} \right\rangle$$

Because of the equations (2.23) and (2.37) we can obtain

$$(2.49) \quad \Lambda_C = \cosh \Omega \oint \tau dt - \sinh \Omega \oint \kappa dt$$

If we use the equations (2.27) and (2.34) into the last equations, we get

$$(2.50) \quad \Lambda_C = - \cosh \Omega \Lambda_{U_1} + \sinh \Omega \Lambda_{U_3}$$

For the drall of the closed surface , we may write

$$P_C = \frac{\langle d\vec{c}, d\vec{c}^* \rangle}{\langle d\vec{c}, d\vec{c} \rangle}$$

(2.51)

$$P_C = \frac{\omega' \omega^{*'} + (k_1 \sinh \omega - k_2 \cosh \omega) [(k_1^* - k_2 \omega^*) \sinh \omega + (k_1 \omega^* - k_2^*) \cosh \omega]}{(k_1 \sinh \omega - k_2 \cosh \omega)^2 + \omega'^2}$$

Now, let is calculate the integral invariants of the closed ruled surfaces respectively. The pitch of the closed surface is obtained as

$$L_{V_1} = \left\langle \vec{d}, \vec{v}_1^* \right\rangle + \left\langle \vec{d}^*, \vec{v}_1 \right\rangle,$$

(2.52)

$$L_{V_1} = \oint q^* dt.$$

Substituting by the value into the equation (2.52)

$$(2.53) \quad L_{V_1} = \sinh \varphi \oint k_1^* dt + \cosh \varphi \oint k_2^* dt + \varphi^* (\cosh \varphi \oint k_1 dt + \sinh \varphi \oint k_2 dt)$$

or

$$(2.54) \quad L_{V_1} = \cosh \phi L_{u_1} + \sinh \phi L_{u_3} - \phi^* (\sinh \phi \lambda_{u_1} + \cosh \phi \lambda_{u_3}).$$

For the dual angle of the pitch of the closed ruled surface , we may write

$$\Lambda_{V_1} = - \left\langle \vec{D}, \vec{V}_1 \right\rangle$$

Because of the equation (2.21) we can obtain

$$(2.55) \quad \Lambda_{V_1} = - \oint Q dt.$$

Substituting by the equation (2.15) into the last equation, we get

$$\Lambda_{V_1} = -\sinh \Phi \oint \kappa dt - \cosh \Phi \oint \tau dt$$

or

$$(2.56) \quad \Lambda_{V_1} = \cosh \Phi \wedge_{U_1} + \sinh \Phi \wedge_{U_3}$$

If the equation (2.56) is separated into the dual and real part, we can obtain

$$(2.57) \quad \begin{cases} \lambda_{v_1} = \cosh \phi \lambda_{u_1} + \sinh \phi \lambda_{u_3} \\ L_{v_1} = \cosh \phi L_{u_1} + \sinh \phi L_{u_3} - \phi^* (\sinh \phi \lambda_{u_1} + \cosh \phi \lambda_{u_3}) \end{cases}$$

For the drall of the closed surface , we may write

$$P_{V_1} = \frac{\langle d\vec{v}_1, d\vec{v}_1^* \rangle}{\langle d\vec{v}_1, d\vec{v}_1^* \rangle}$$

Setting by the values of the statements $d\vec{v}_1$ and $d\vec{v}_1^*$ as the equations (2.13) into the last equations, we get

$$(2.58) \quad P_{V_1} = \frac{p^*}{p}$$

Setting by the values of p and p^* as the equations (2.16) into the last equations, we get

$$(2.59) \quad P_{V_1} = \frac{k_1^* \cosh \phi + k_2^* \sinh \phi}{k_1 \cosh \phi + k_2 \sinh \phi} + \phi^* \frac{k_1 \sinh \phi + k_2 \cosh \phi}{k_1 \cosh \phi + k_2 \sinh \phi}$$

2.2. Theorem: Let (V_1) be the parallel surface of the surface (U_1) . The pitch, drall and the dual of the pitch of the ruled surface (V_1) are

$$1-) L_{V_1} = \oint q^* dt \quad 2-) \Lambda_{V_1} = - \oint Q dt \quad 3-) P_{V_1} = \frac{p^*}{p}$$

2.3. Corollary: Let (V_1) be the parallel surface of the surface (U_1) . The pitch and the dual of the pitch of the ruled surface (V_1) related to the invariants of the surface (U_1) are written as follow

$$1-) L_{V_1} = \cosh \phi L_{u_1} + \sinh \phi L_{u_3} - \phi^* (\sinh \phi \lambda_{u_1} + \cosh \phi \lambda_{u_3})$$

$$2-) \Lambda_{V_1} = \cosh \phi \wedge_{U_1} + \sinh \phi \wedge_{U_3}$$

The pitch of the closed surface (V_2) is obtained as

$$L_{V_2} = \langle \vec{d}, \vec{v}_2^* \rangle + \langle \vec{d}^*, \vec{v}_2 \rangle$$

$$(2.60) \quad L_{V_2} = 0$$

For the dual angle of the pitch of the closed ruled surface (V_2) , we may write

$$\Lambda_{V_2} = -\langle \vec{D}, \vec{V}_2 \rangle$$

Because of the equation (2.21) we can obtain

$$(2.61) \quad \Lambda_{V_2} = 0$$

For the drall of the closed surface (V_2) , we may write

$$P_{V_2} = \frac{\langle d\vec{v}_2, d\vec{v}_2^* \rangle}{\langle d\vec{v}_2, d\vec{v}_2 \rangle}$$

Setting by the values of the statements $d\vec{v}_2$ and $d\vec{v}_2^*$ as the equations (2.13) into the last equations, we get

$$(2.62) \quad P_{V_2} = \frac{pp^* - qq^*}{p^2 - q^2}$$

Setting by the values of p, p^*, q and q^* as the equations (2.16) into the last equations, we get

$$(2.63) \quad P_{V_2} = \frac{k_1 k_1^* - k_2 k_2^*}{k_1^2 - k_2^2}$$

2.4. Theorem: Let (V_1) be the parallel surface of the surface (U_1) . The pitch, drall and the dual of the pitch of the ruled surface (V_2) are

$$1-) L_{V_2} = 0 \quad 2-) \Lambda_{V_2} = 0 \quad 3-) P_{V_2} = \frac{pp^* - qq^*}{p^2 - q^2}$$

The pitch of the closed surface (V_3) is obtained as

$$L_{V_3} = \langle \vec{d}, \vec{v}_3^* \rangle + \langle \vec{d}^*, \vec{v}_3 \rangle,$$

$$(2.64) \quad L_{V_3} = \oint p^* dt$$

Substituting by the value into the equation (2.64)

$$(2.65) \quad L_{V_3} = \cosh \varphi \oint k_1^* dt + \sinh \varphi \oint k_2^* dt + \varphi^* (\sinh \varphi \oint k_1 dt + \cosh \varphi \oint k_2 dt)$$

or

$$(2.66) \quad L_{V_3} = \sinh \phi L_{u_1} + \cosh \phi L_{u_3} - \phi^* (\cosh \phi \lambda_{u_1} + \sinh \phi \lambda_{u_3})$$

For the dual angle of the pitch of the closed ruled surface , we may write

$$\Lambda_{V_3} = - \langle \vec{D}, \vec{V}_3 \rangle$$

Because of the equation (2.21) we can obtain

$$(2.67) \quad \Lambda_{V_3} = - \oint P dt.$$

Substituting by the equation (2.6) into the last equation, we get

$$\Lambda_{V_3} = - \cosh \Phi \oint \kappa dt - \sinh \Phi \oint \tau dt$$

or

$$(2.68) \quad \Lambda_{V_3} = \sinh \Phi \Lambda_{U_1} + \cosh \Phi \Lambda_{U_3}$$

If the equation (2.68) is separated into the dual and real part, we can obtain

$$(2.69) \quad \begin{cases} \lambda_{v_3} = \sinh \phi \lambda_{u_1} + \cosh \phi \lambda_{u_3} \\ L_{v_3} = \sinh \phi L_{u_1} + \cosh \phi L_{u_3} - \phi^* (\cosh \phi \lambda_{u_1} + \sinh \phi \lambda_{u_3}) \end{cases}$$

For the drall of the closed surface , we may write

$$P_{V_3} = \frac{\langle d\vec{v}_3, d\vec{v}_3^* \rangle}{\langle d\vec{v}_3, d\vec{v}_3 \rangle}$$

Setting by the values of the statements $d\vec{v}_3$ and $d\vec{v}_3^*$ as the equations (2.13) into the last equations, we get

$$(2.70) \quad P_{V_3} = \frac{q^*}{q}$$

Setting by the values of q and q^* as the equations (2.16) into the last equations, we get

$$(2.71) \quad P_{V_3} = \frac{k_1^* \sinh \phi + k_2^* \cosh \phi}{k_1 \sinh \phi + k_2 \cosh \phi} + \phi^* \left(\frac{k_1 \cosh \phi + k_2 \sinh \phi}{k_1 \sinh \phi + k_2 \cosh \phi} \right)$$

2.5. Theorem: Let (V_1) be the parallel surface of the surface (U_1) . The pitch, drall and the dual of the pitch of the ruled surface (V_3) are

$$1-) L_{V_3} = \oint p^* dt \quad 2-) \Lambda_{V_3} = - \oint P dt \quad 3-) P_{V_3} = \frac{q^*}{q}$$

2.6. Corollary: Let (V_1) be the parallel surface of the surface (U_1) . The pitch and the dual of the pitch of the ruled surface (V_3) related to the invariants of the surface (U_1) are written as follow

$$\begin{aligned} 1-) L_{V_3} &= \sinh \phi L_{u_1} + \cosh \phi L_{u_3} - \phi^* (\cosh \phi \lambda_{u_1} + \sinh \phi \lambda_{u_3}) \\ 2-) \wedge_{V_3} &= \sinh \Phi \wedge_{U_1} + \cosh \Phi \wedge_{U_3} \end{aligned}$$

Let $\Theta(t) = \theta(t) + \varepsilon \theta^*(t)$ be Lorentzian timelike angle of between the instantaneous dual Pfaffion vector $\vec{\Psi}$ and the vector $\vec{V}_3$ [10].

a) If the instantaneous dual Pfaffion vector $\vec{\Psi}$ is spacelike ($|Q| > |P|$)

$$\left\{ P = \left\| \vec{\Psi} \right\| \sinh \Theta \quad , \quad Q = \left\| \vec{\Psi} \right\| \cosh \Theta \right.$$

On the way $\vec{C} = \vec{c} + \varepsilon \vec{c}^*$, unit vector about the vector $\vec{\Psi}$ direction is

$$(2.72) \quad \vec{C} = \cosh \Theta \vec{V}_1 - \sinh \Theta \vec{V}_3$$

Setting by the values of the vectors $\vec{V}_1$ and $\vec{V}_3$ as the equations (2.9) into the equation (2.72), we get

$$\vec{C} = \cosh \Theta (\cosh \Phi \vec{U}_1 + \sinh \Phi \vec{U}_3) - \sinh \Theta (\sinh \Phi \vec{U}_1 + \cosh \Phi \vec{U}_3)$$

$$(2.73) \quad \vec{C} = \cosh (\Theta - \Phi) \vec{U}_1 - \sinh (\Theta - \Phi) \vec{U}_3$$

If the equation (2.72) is separated into the dual and real part, we can obtain

$$(2.74) \quad \begin{cases} \vec{c} = \cosh \theta \vec{v}_1 - \sinh \theta \vec{v}_3 \\ \vec{c}^* = \cosh \theta \vec{v}_1^* - \sinh \theta \vec{v}_3^* + \theta^* \sinh \theta \vec{v}_1 - \theta^* \cosh \theta \vec{v}_3 \end{cases}$$

The pitch of the closed surface is obtained as

$$L_{\vec{C}} = \langle \vec{d}, \vec{c}^* \rangle + \langle \vec{d}^*, \vec{c} \rangle$$

Setting by the values of the statements $\vec{d}$ and $\vec{d}^*$ as the equations (2.22) into the last equations and if we do the necessary operations, we get

$$(2.75) \quad L_{\vec{C}} = -\sinh \theta \oint p^* dt + \cosh \theta \oint q^* dt + \theta^* \left(\sinh \theta \oint q dt - \cosh \theta \oint p dt \right)$$

or

$$(2.76) \quad L_{\vec{C}} = \cosh \theta L_{V_1} - \sinh \theta L_{V_3} + \theta^* (-\sinh \theta \lambda_{V_1} + \cosh \theta \lambda_{V_3})$$

If we use the equations (2.57) and (2.69) into the equation (2.76) and necessary operations have been done, we get

$$(2.77) \quad \begin{aligned} L_{\vec{C}} &= \cosh (\theta - \phi) L_{U_1} - \sinh (\theta - \phi) L_{U_3} + \\ &\quad (\phi^* - \theta^*) (\sinh (\theta - \phi) \lambda_{U_1}) - \cosh (\theta - \phi) \lambda_{U_3} \end{aligned}$$

For the dual angle of the pitch of the closed ruled surface , we may write

$$\Lambda_{\overrightarrow{C}} = - \left\langle \overrightarrow{D}, \overrightarrow{C} \right\rangle$$

Because of the equations (2.21) and (2.72) we can obtain

$$(2.78) \quad \Lambda_{\overrightarrow{C}} = - \cosh \Theta \oint Q dt + \sinh \Theta \oint P dt$$

Setting by the values of the statements $\Lambda_{V_1} = - \oint Q dt$ and $\Lambda_{V_3} = - \oint P dt$ as the equations (2.55) and (2.67) into the last equations, we get

$$(2.79) \quad \Lambda_{\overrightarrow{C}} = \cosh \Theta \Lambda_{V_1} - \sinh \Theta \Lambda_{V_3}$$

If we use the equations (2.56) and (2.68) into the equations (2.79), we get

$$(2.80) \quad \Lambda_{\overrightarrow{C}} = \cosh (\Theta - \Phi) \Lambda_{U_1} - \sinh (\Theta - \Phi) \Lambda_{U_3}$$

For the drall of the closed surface , we may write

$$P_{\overrightarrow{C}} = \frac{\left\langle d \overrightarrow{C}, d \overrightarrow{C}^* \right\rangle}{\left\langle d \overrightarrow{C}, d \overrightarrow{C} \right\rangle}$$

(2.81)

$$P_{\overrightarrow{C}} = \frac{-\theta' \theta^{*'} + (p \cosh \theta - q \sinh \theta) [(p^* - q \theta^*) \cosh \theta + (p \theta^* - q^*) \sinh \theta]}{(p \cosh \theta - q \sinh \theta)^2 - \theta'^2}$$

b) If the instantaneous dual Pfaffion vector $\overrightarrow{\Psi}$ is timelike ($|Q| < |P|$)

$$P = \left\| \overrightarrow{\Psi} \right\| \cosh \Theta, \quad Q = \left\| \overrightarrow{\Psi} \right\| \sinh \Theta$$

On the way $\overrightarrow{C} = \overrightarrow{C} + \varepsilon \overrightarrow{C}^*$, unit vector about the vector $\overrightarrow{\Psi}$ direction is

$$(2.82) \quad \overrightarrow{C} = \sinh \Theta \overrightarrow{V_1} - \cosh \Theta \overrightarrow{V_3}$$

Setting by the values of the vectors $\overrightarrow{V_1}$ and $\overrightarrow{V_3}$ as the equations (2.9) into the equation (2.82), we get

$$\overrightarrow{C} = \sinh \Theta \left(\cosh \Phi \overrightarrow{U_1} + \sinh \Phi \overrightarrow{U_3} \right) - \cosh \Theta \left(\sinh \Phi \overrightarrow{U_1} + \cosh \Phi \overrightarrow{U_3} \right)$$

$$(2.83) \quad \overrightarrow{C} = \sinh (\Theta - \Phi) \overrightarrow{U_1} + \cosh (\Theta - \Phi) \overrightarrow{U_3}$$

If the equation (2.82) is separated into the dual and real part, we can obtain

$$(2.84) \quad \begin{cases} \overrightarrow{C} = \sinh \theta \overrightarrow{v_1} - \cosh \theta \overrightarrow{v_3} \\ \overrightarrow{C}^* = \sinh \theta \overrightarrow{v_1}^* - \cosh \theta \overrightarrow{v_3}^* + \theta^* \cosh \theta \overrightarrow{v_1} - \theta^* \sinh \theta \overrightarrow{v_3} \end{cases}$$

The pitch of the closed surface is obtained as

$$L_{\overline{C}} = \langle \vec{d}, \vec{c}^* \rangle + \langle \vec{d}^*, \vec{c} \rangle$$

Setting by the values of the statements $\vec{d}$ and $\vec{d}^*$ as the equations (2.22) into the last equations and if we do the necessary operations , we get

$$(2.85) \quad L_{\overline{C}} = -\cosh \theta \oint p^* dt + \sinh \theta \oint q^* dt + \theta^* \left(\cosh \theta \oint q dt - \sinh \theta \oint p dt \right)$$

or

$$(2.86) \quad L_{\overline{C}} = \sinh \theta L_{V_1} - \cosh \theta L_{V_3} + \theta^* (-\cosh \theta \lambda_{V_1} + \sinh \theta \lambda_{V_3})$$

If we use the equations (2.57) and (2.69) into the equation (2.86) and necessary operations have been done, we get

$$(2.87) \quad \begin{aligned} L_{\overline{C}} = & \sinh (\theta - \phi) L_{U_1} - \cosh (\theta - \phi) L_{U_3} + \\ & + (\phi^* - \theta^*) (\cosh (\theta - \phi) \lambda_{U_1} - \sinh (\theta - \phi) \lambda_{U_3}) \end{aligned}$$

For the dual angle of the pitch of the closed ruled surface , we may write

$$\Lambda_{\overline{C}} = - \left\langle \vec{D}, \vec{C} \right\rangle$$

Because of the equations (2.21) and (2.82) we can obtain

$$(2.88) \quad \Lambda_{\overline{C}} = -\sinh \Theta \oint Q dt + \cosh \Theta \oint P dt$$

If we use the equations (2.55) and (2.67) into the last equations, we get

$$(2.89) \quad \Lambda_{\overline{C}} = \sinh \Theta \Lambda_{V_1} - \cosh \Theta \Lambda_{V_3}$$

If we use the equations (2.56) and (2.68) into the equations (2.89), we get

$$(2.90) \quad \Lambda_{\overline{C}} = \sinh (\Theta - \Phi) \Lambda_{U_1} - \cosh (\Theta - \Phi) \Lambda_{U_3}$$

For the drall of the closed surface , we may write

$$P_{\overline{C}} = \frac{\langle d \vec{c}, d \vec{c}^* \rangle}{\langle d \vec{c}, d \vec{c} \rangle}$$

$$(2.91) \quad P_{\overline{C}} = \frac{\theta' \theta^{*'} + (p \sinh \theta - q \cosh \theta) [(p \theta^* - q^*) \cosh \theta + (p^* - q \theta^*) \sinh \theta]}{(p \sinh \theta - q \cosh \theta)^2 + \theta'^2}$$

References

- [1] Birman, G.S. and Nomizu, K., *Trigonometry in Lorentzian Geometry*, Am. Math. Mont., 91 (1984), 543-549.
- [2] Clifford, W.K., *Preliminary sketch of biquaternions*, Proceedings of London Math. Soc. 4 (1873) 361-395.
- [3] Erim, K., *Lessons of Differential Geometry*, Istanbul, 1949, 399p (in Turkish).
- [4] Study, E., *Geometrie der Dynamen*, Leipzig, 1903.
- [5] Gursoy, O., *The Dual Angle of Pitch of a Closed Ruled Surface*, Mech. and Mach. Theory, 25(1990),2, 131-140.
- [6] Hacısalihoğlu, H. H., *Moving Geometry and Quaternion Theory*, Publishings of Science and Art Faculty of Gazi University, Math. No: 2, Ankara,(1983) 338p (in Turkish).
- [7] Hacısalihoğlu, H. H., *On the Pitch of a Closed Ruled Surface*, Mech. and Mach. Theory, Pargaman Press. Printed in Great Britian, 7(1972)291-305.
- [8] Hacısalihoğlu, H. H., *Differential Geometry*, Publishings of Science and Art Faculty of Inonu University, Math. No:2, Malatya (1983), 894p.
- [9] Muller, H. R., *Kinematic Lessons*, Publishings of Science Faculty of Ankara University, Mat.27, Ankara (1963), 292p(in Turkish).
- [10] Ratcliffe, J. G., *Foundations of Hyperbolic Manifolds*, Springer-Verlag New York, Inc., New York,(1994) 736p.
- [11] Senyurt, S., *Parallel Regle Surfaces and Properties of Some Characteristic*, Institute of Science of Ondokuz Mayıs University Ph. D. Thesis(1999).
- [12] Turgut, A., *Spacelike and Timelike Regle Surfaces in 3-dim. Minkowski Space*, Institute of Science of Ankara University, Ankara Ph. D. Thesis, (1995).
- [13] Ugurlu, H. H., *On the Geometry of Timelike Surfaces*, Commun. Fac. Sci. Ank. Series A1 V.46(1997), 211-223.
- [14] Ugurlu, H. H., Çalışkan, A., *The study mapping for directed spacelike and timelike lines in Minkowski 3-space.*, Math. Comp. Appl. 1, 2 (1996), 142-148.
- [15] Woestijne, V. D. I., *Minimal Surfaces of the 3-dimensional Minkowski space*, Proc. Congres "Géométrie différentielle et applications" Avignon Word Scientific Publishing., Singapore(1988), 344-369.