

PRIMITIVE REPRESENTATIONS OF INTEGERS BY $x^3 + y^3 + 2z^3$

SAMIR SIKSEK

ABSTRACT. A well-known open problem is to show that the cubic form $x^3 + y^3 + 2z^3$ represents all integers. An obvious variant of this problem is whether every integer can be *primitively* represented by $x^3 + y^3 + 2z^3$. In other words, given an integer n , are there coprime integers x, y, z such that $x^3 + y^3 + 2z^3 = n$? In this note we answer this variant question negatively. Indeed, we use cubic reciprocity to show that for every integral solution to $x^3 + y^3 + 2z^3 = 8^m$, the unknowns x, y, z are divisible by 2^m .

Guy [2, Problem D5] asks if every number is the sum of four cubes with two of them equal. In other words, can every integer be represented by the cubic form $x^3 + y^3 + 2z^3$. Guy notes that this has been shown to be the case for all positive integers < 1000 except for 148, 671, 788 which are still open.

An obvious variant of this problem is whether every integer can be *primitively* represented by $x^3 + y^3 + 2z^3$. In other words, given an integer n , are there coprime integers x, y, z such that $x^3 + y^3 + 2z^3 = n$? In this note we answer this question negatively, as the following theorem shows.

Theorem 1. *Let $m \geq 1$. If x, y, z are integers satisfying*

$$(1) \quad x^3 + y^3 + 2z^3 = 8^m$$

then x, y and z are divisible by 2^m .

The proof, given in Section 2, uses cubic reciprocity. The use of cubic reciprocity is inspired by Cassels' [1] where he shows that any integral solution to $x^3 + y^3 + z^3 = 3$ satisfies $x \equiv y \equiv z \pmod{9}$. We summarize what we need from cubic reciprocity in Section 1.

1. CUBIC RECIPROCITY

Good references for cubic reciprocity are [3, Chapter 9] and [4, Chapter 7]. We work over the Eisenstein ring $\mathbb{E} = \mathbb{Z}[\varrho]$ where $\varrho = (-1 + \sqrt{-3})/2$. Recall that $\mathbb{E}$ is a unique factorisation domain. The prime 2 is inert in $\mathbb{E}$ while 3 ramifies in $\mathbb{E}$ as $3\mathbb{E} = (1 - \varrho)^2\mathbb{E}$.

We use $(\div)$ to denote the *cubic* residue symbol which takes values in the cube roots of unity $\mathcal{P} = \{1, \varrho, \varrho^2\}$. One way of defining the cubic residue symbol is as follows: let α, π be coprime elements of $\mathbb{E}$ with π prime, and suppose $\pi \nmid 3$. Clearly

$$\alpha^{(N\pi-1)/3} \equiv \epsilon \pmod{\pi}$$

Date: August 8, 2011.

2000 *Mathematics Subject Classification.* Primary 11D85, Secondary 11D25, 11G35.

The author is supported by an EPSRC Leadership Fellowship and by a Marie-Curie International Reintegration Grant (MIRG-CT-2006-044530).

for some unique $\epsilon \in \mathcal{P}$, where $N\pi$ is the norm of π . We define $\left(\frac{\alpha}{\pi}\right)$ to be this unique ϵ . If π and π' are associates (that is the same up to units) then clearly

$$\left(\frac{\alpha}{\pi}\right) = \left(\frac{\alpha}{\pi'}\right).$$

If α, β are coprime elements of $\mathbb{E}$, with $(1 - \varrho) \nmid \beta$ then we define

$$\left(\frac{\alpha}{\beta}\right) = \left(\frac{\alpha}{\pi_1}\right) \cdots \left(\frac{\alpha}{\pi_n}\right),$$

where $\beta = \pi_1 \cdots \pi_n$ is a prime factorisation of β ; replacing one prime factorisation by another only changes the order and the values of the primes up to units and so does not affect the definition. If $\alpha \equiv \alpha' \pmod{\beta}$ then

$$\left(\frac{\alpha}{\beta}\right) = \left(\frac{\alpha'}{\beta}\right).$$

Moreover, it is easy to see that,

$$\left(\frac{\alpha_1 \alpha_2}{\beta}\right) = \left(\frac{\alpha_1}{\beta}\right) \left(\frac{\alpha_2}{\beta}\right), \quad \left(\frac{\alpha}{\beta_1 \beta_2}\right) = \left(\frac{\alpha}{\beta_1}\right) \left(\frac{\alpha}{\beta_2}\right).$$

The only non-elementary fact we need is Eisenstein's Cubic Reciprocity Law.

Theorem (Eisenstein's Cubic Reciprocity Law). *Suppose $\alpha \in \mathbb{E}$ and $a \in \mathbb{Z}$ are coprime and are both $\equiv \pm 1 \pmod{3}$. Then*

$$\left(\frac{\alpha}{a}\right) = \left(\frac{a}{\alpha}\right).$$

Moreover, if $a \in \mathbb{Z}$ satisfies $a \equiv 1 \pmod{3}$ then

$$\left(\frac{\varrho}{a}\right) = \varrho^{(1-a)/3}, \quad \left(\frac{1-\varrho}{a}\right) = \varrho^{(a-1)/3}.$$

The second part of the theorem is often called the 'supplement to the law of cubic reciprocity'. For a proof see [4, pages 213–214].

2. PROOF OF THEOREM 1

The proof is by induction on m . If one of x, y is even then all three unknowns x, y, z are even and we reduce to the case where m is replaced by $m - 1$. Thus suppose that x and y are odd and it is sufficient to derive a contradiction.

Note that if a is an integer then $a^3 \equiv 0, 1, -1 \pmod{9}$. Considering (1) modulo 9 we see that either $x^3 \equiv \pm 1 \pmod{9}$ and $y^3 \equiv 0 \pmod{9}$ or the other way round. We swap x and y if necessary so that these two congruences hold. Then

$$(2) \quad x \equiv \pm 1 \pmod{3}, \quad \text{and} \quad y \equiv 0 \pmod{3}.$$

From (1) we see that

$$2z^3 \equiv 8^m \pmod{x + y\varrho}.$$

Since x, y are odd, $x + y\varrho \equiv 1 + \varrho \pmod{2}$. Moreover, $x + y\varrho$ and $2z^3$ are coprime. It follows that

$$\left(\frac{2}{x + y\varrho}\right) = \left(\frac{2z^3}{x + y\varrho}\right) = \left(\frac{8^m}{x + y\varrho}\right) = \left(\frac{2}{x + y\varrho}\right)^{3m} = 1.$$

Now, by (2), $x + y\rho \equiv \pm 1 \pmod{3}$. Applying Eisenstein's Cubic Reciprocity Law we have

$$\left(\frac{x + y\rho}{2}\right) = 1.$$

However,

$$\left(\frac{x + y\rho}{2}\right) = \left(\frac{1 + \rho}{2}\right) = \left(\frac{-\rho^2}{2}\right) = \left(\frac{\rho}{2}\right)^2 = \left(\frac{\rho}{-2}\right)^2 = \rho^2,$$

using the supplement to Eisenstein's Cubic Reciprocity Law. This contradiction completes the proof of Theorem 1.

REFERENCES

- [1] J. W. S. Cassels, A note on the Diophantine equation $x^3 + y^3 + z^3 = 3$, *Mathematics of Computation* **44** (1985), 265–266.
- [2] R. K. Guy, *Unsolved Problems in Number Theory*, third edition, Springer, 2003.
- [3] K. Ireland and M. Rosen, *A Classical Introduction to Modern Number Theory*, third edition, GTM **84**, Springer-Verlag, New York, 1990.
- [4] F. Lemmermeyer, *Reciprocity Laws: from Euler to Eisenstein*, Springer, Berlin, 2000.

MATHEMATICS INSTITUTE, UNIVERSITY OF WARWICK, COVENTRY, CV4 7AL, UNITED KINGDOM
E-mail address: `s.siksek@warwick.ac.uk`