

A Maximal Large Deviation Inequality for Sub-Gaussian Variables

Dotan Di Castro¹, Claudio Gentile², and Shie Mannor¹

¹Technion, Israel Institute of Technology, Haifa, Israel

²Universita' dell'Insubria, Varese, Italy

dot@tx.technion.ac.il, claudio.gentile@uninsubria.it,
shie@ee.technion.ac.il

Abstract. In this short note we prove a maximal concentration lemma for sub-Gaussian random variables implying that for independent sub-Gaussian random variables we have

$$P\left(\max_{1 \leq i \leq N} S_i > \epsilon\right) \leq \exp\left(-\frac{1}{N^2} \sum_{i=1}^N \frac{\epsilon^2}{2\sigma_i^2}\right),$$

where S_i is the sum of the first i zero-mean independent sub-Gaussian random variables, and σ_i^2 is the variance of the i th random variable.

Definition 1. Let X be a random variable, and σ^2 be a positive constant. If X satisfies

$$E[e^{sX}] \leq \exp\left\{\frac{1}{2}s^2\sigma^2\right\} \quad \forall s \in \mathcal{R}$$

then X is a sub-Gaussian random variable with parameter σ^2 .

Lemma 1. Let $X_1, \dots, X_N$ be independent (zero-mean) sub-Gaussian random variables with parameters $\sigma_1^2, \dots, \sigma_N^2$, respectively. Let $S_i = \sum_{j=1}^i X_j$. Then,

$$P\left(\max_{1 \leq i \leq N} S_i > \epsilon\right) \leq \exp\left(-\frac{1}{N^2} \sum_{i=1}^N \frac{\epsilon^2}{2\sigma_i^2}\right).$$

We begin by stating a general maximal bound on a martingale difference sequence using rate functions. Then, we apply it to the rate function for sub-Gaussian variables, and end up proving Lemma 1.

Theorem 1. Let $X_1, \dots, X_N$ be independent random variables where $\mathbb{E}[X_i] = 0$ for $1 \leq i \leq N$. For each $1 \leq i \leq N$ define the logarithmic generating function $\rho_i(s) = \log(\mathbb{E}[e^{sX_i}])$, and assume that they are finite in the neighborhood of 0. Let $\epsilon_1, \dots, \epsilon_N$ be a sequence of positive numbers, and define the i -th rate function f_i as $f_i(\epsilon_i) = \sup_s(\epsilon_i s - \rho_i(s))$. Let $\bar{X}_i = \sum_{j=1}^i X_j$ and $\bar{\epsilon}_i = \sum_{j=1}^i \epsilon_j$. Then, for $1 \leq i \leq N$ we have $f_i(\epsilon_i) > 0$ and

$$P\left(\max_{1 \leq i \leq N} \bar{X}_i \geq \bar{\epsilon}_N\right) \leq \exp\left(-\sum_{j=1}^N f_j(\epsilon_j)\right). \quad (1)$$

Proof. Define $S_0 = 0$ and

$$S_{k+1} = \begin{cases} S_k + X_{k+1} & \text{if } S_k < \sum_{i=1}^N \epsilon_i, \\ S_k & \text{if } S_k \geq \sum_{i=1}^N \epsilon_i, \end{cases} \quad (2)$$

i.e., S_k is identical to $\bar{X}_k$ until S_k exceeds $\sum_{i=1}^N \epsilon_i$, and afterward it maintains the highest value. Note that the following two events are equal:

$$\left\{ \max_{1 \leq i \leq N} \bar{X}_i > \sum_{i=1}^N \epsilon_i \right\} = \left\{ S_N \geq \sum_{i=1}^N \epsilon_i \right\}. \quad (3)$$

For the rest of the proof we concentrate on the r.h.s. of (3). Notice that $\rho_i(0) = \log(\mathbb{E}[e^{0 \cdot X_i}]) = \log(1) = 0$. Moreover, using the Jensen's inequality and the fact that the r.v. averages are 0 we have for all $s \neq 0$

$$\mathbb{E}[\exp(sX_i)] \geq \exp(\mathbb{E}[sX_i]) = e^0 = 1. \quad (4)$$

Taking the log of (4) and using $\rho_i(0) = 0$ yields $\rho_i(s) \geq 0$. Let us fix some s where $\rho_i(s)$ is finite (such s exists by the theorem's assumptions) and define the following stopping time

$$L = \begin{cases} N & \text{if } \bar{X}_n < \sum_{i=1}^n \epsilon_i \quad \text{for } n = 1, \dots, N, \\ \min \{n : \bar{X}_n \geq \sum_{i=1}^n \epsilon_i\} & \text{otherwise.} \end{cases}$$

Define Y_k to be

$$Y_k = \exp\left(sS_k - \sum_{i=1}^{\min(k, L)} \rho_i(s)\right). \quad (5)$$

We claim that Y_k is a martingale. Trivially, if $k \geq L$ then $\min(L, k) = N$, by (2) we have $S_{k+1} = S_k$, and by (5) $Y_{k+1} = Y_k$. Let us define the σ -algebra $\mathcal{F}_k \triangleq \sigma(X_1, \dots, X_k)$. If $k < N$ then

$$\begin{aligned} \mathbb{E}[Y_{k+1} | \mathcal{F}_k] &= \mathbb{E}\left[\exp\left(sS_{k+1} - \sum_{i=1}^{k+1} \rho_i(s)\right) \middle| \mathcal{F}_k\right] \\ &= \mathbb{E}\left[\exp\left(sS_k - \sum_{i=1}^k \rho_i(s)\right) \cdot \exp(sX_{k+1} - \rho_{k+1}(s)) \middle| \mathcal{F}_k\right] \\ &= \exp\left(sS_k - \sum_{i=1}^k \rho_i(s)\right) \cdot \mathbb{E}[\exp(sX_{k+1} - \rho_{k+1}(s)) | \mathcal{F}_k] \\ &= Y_k \mathbb{E}[\exp(sX_{k+1} - \log(\mathbb{E}[e^{sX_{k+1}}]))] \\ &= Y_k \mathbb{E}[\exp(sX_{k+1}) (\mathbb{E}[e^{sX_{k+1}}])^{-1}] \\ &= Y_k, \end{aligned}$$

and $(Y_k)_{k=1}^N$ is a martingale. Next, we claim that $\mathbb{E}[Y_1] = 1$:

$$\begin{aligned}\mathbb{E}[Y_1] &= \mathbb{E} \left[\exp \left(sX_1 - \sum_{i=1}^{\min(1,L)} \rho_i(s) \right) \right] \\ &= \mathbb{E} [\exp(sX_1) \exp(-\log(\mathbb{E}[e^{sX_1}]))] \\ &= 1,\end{aligned}$$

therefore, $\mathbb{E}[Y_N] = \mathbb{E}[Y_{N-1}] = \dots = \mathbb{E}[Y_1] = 1$. Now,

$$\mathbb{E} \left[\exp \left(sX_N - \sum_{i=1}^N \rho_i(s) \right) \right] \leq \mathbb{E} \left[\exp \left(sX_N - \sum_{i=1}^{\min(L,N)} \rho_i(s) \right) \right] = \mathbb{E}[Y_N] = 1,$$

which yields

$$\mathbb{E}[\exp(sX_N)] \leq \mathbb{E} \left[\exp \left(\sum_{i=1}^N \rho_i(s) \right) \right]. \quad (6)$$

Next, using Markov inequality and (6) we can write

$$\begin{aligned}P\left(S_N \geq \sum_{i=1}^N \epsilon_i\right) &= P\left(\exp(sS_N) \geq \exp\left(s \sum_{i=1}^N \epsilon_i\right)\right) \\ &\leq \frac{\mathbb{E}[\exp(sS_N)]}{\exp\left(s \sum_{i=1}^N \epsilon_i\right)} \\ &\leq \exp\left(\sum_{i=1}^N \rho_i(s) - s \sum_{i=1}^N \epsilon_i\right) \\ &= \exp\left(-\sum_{i=1}^N (s\epsilon_i - \rho_i(s))\right)\end{aligned} \quad (7)$$

Because this is true for every s for which $\rho(s)$ is finite, we can take the infimum of the r.h.s. over s , which yields (1). $\square$

The following lemma characterizes the rate function of a sub-Gaussian random variable.

Lemma 2. *Let X be a sub-Gaussian random variable with parameter σ^2 , and let $\epsilon > 0$. Then,*

$$f(\epsilon) \triangleq \sup_s (\epsilon s - \log(\mathbb{E}[e^{sX}])) \geq \frac{\epsilon^2}{2\sigma^2}.$$

Proof. We have

$$\begin{aligned} & \sup_s \left\{ \epsilon s - \log \left(\mathbb{E} \left[e^{sX} \right] \right) \right\} \\ & \geq \sup_s \left\{ \epsilon s - \log \left(e^{\frac{1}{2} \sigma^2 s^2} \right) \right\} \\ & = \sup_s \left\{ \epsilon s - \frac{1}{2} \sigma^2 s^2 \right\} = \frac{\epsilon^2}{2\sigma^2}, \end{aligned}$$

where the inequality stems from variable X to be a sub-Gaussian random variable. $\square$

Next, we prove Lemma 1.

Proof. (Lemma 1) Directly follows from Theorem 1 and Lemma 2. Let $\epsilon_i = \epsilon$. Then by Theorem 1

$$P\left(\max_{1 \leq i \leq N} \bar{X}_i \geq N\epsilon\right) \leq \exp\left(-\sum_{j=1}^N f_i(\epsilon)\right).$$

Slightly abusing notation and using Lemma 2 we have

$$\begin{aligned} P\left(\max_{1 \leq i \leq N} \bar{X}_i \geq \epsilon\right) & \leq \exp\left(-\sum_{j=1}^N f_i(\epsilon/N)\right) \\ & \leq \exp\left(-\sum_{j=1}^N \frac{(\epsilon/N)^2}{2\sigma_i^2}\right). \end{aligned}$$