arXiv:1105.3150v2 [math.DG] 28 May 2011

The space of Constant Mean Curvature surfaces in compact Riemannian Manifolds

José M. Espinar [†1]

December 31, 2013

[†] Departamento de Geometría y Topología, Universidad de Granada, 18071 Granada, Spain; e-mail: jespinar@ugr.es

Abstract

The main point of this paper is that, under suitable conditions on the mean curvature and the Ricci curvature of the ambient space, we can extend Choi-Schoen's Compactness Theorem [1] to compact embedded minimal surfaces to simple immersed compact $H-$surfaces in a Riemannian manifold with positive Ricci curvature (the mean curvature small depending on the Ricci curvature).

Also, we prove that the space of convex embedded (fixed) constant mean curvature hypersurfaces in a simply connected $1/4-$pinched manifold is compact.

Mathematical Subject Classification (2010): 53A10, 53C21, 47H11.

Keywords: Constant Mean Curvature, Compactness Theorem, Degree Theory.

1 Introduction

One of the most celebrated compactness results for minimal surfaces is Choi-Schoen's Compactness Theorem [1] for compact minimal surfaces embedded in a Riemannian manifold with positive Ricci curvature. Embeddedness is crucial since it implies area estimates, and area estimates allow to get curvature estimates. Another tool used in Choi-Schoen's Compactness Theorem is the Maximum Principle, this ensures that the limit of a sequence of embedded compact minimal surfaces is embedded. Such a compactness theorem allowed to develop an integer

[1]The author is partially supported by Spanish MEC-FEDER Grant MTM2010-19821, and Regional J. Andalucia Grants P06-FQM-01642 and FQM325

degree theory for compact minimal surfaces embedded in Riemannian manifolds with positive Ricci curvature [9].

But, what is the situation for constant mean curvature surfaces? We can think in the following two examples. First, consider a constant mean curvature Clifford torus in the standard three-sphere. So, we can deform such a Clifford torus as *Delaunay-type* surfaces with the same constant mean curvature and the background metric on the sphere such that the limit is a stack of spheres with the same mean curvature. Second, we can consider a constant mean curvature sphere in the standard three-sphere. We can deform it as embedded spheres with the same constant mean curvature (varying the background metric) until the limit is a constant mean curvature sphere with a point where two points touch , i.e., the limit is not embedded.

The second example tells us that, even thinking about a compactness result for constant mean curvature surfaces, the family of embedded constant mean curvature surfaces is too small. At least, we shall consider those surfaces that arise as limit of embedded surfaces, that is, we shall consider the space of simple immersed constant mean curvature surfaces, i.e., loosely speaking, those that are embbeded up to a set of measure zero and bound a compact connected component (see Definition 3.1 for details). Nonetheless, we still have the first example, that is, it seems that area and curvature estimates are not always valid for simple immersed constant mean curvature surfaces. Actually, using [8], we only need area estimates when we deal with constant mean curvature surfaces.

The main point of this paper is that, under suitable conditions on the mean curvature and the Ricci curvature of the ambient space, we can obtain an uniform area estimate (depending on the genus, the mean curvature and the Ricci curvature) for simple immersed constant mean curvature surfaces, the mean curvature fixed. Hence, we can extend Choi-Schoen's Compactness Theorem to simple immersed compact constant mean curvature surfaces in a Riemannian manifold with positive Ricci curvature (the mean curvature small depending on the Ricci curvature).

The paper is organized as follows. In Section 2 we introduce the necessary notation and tools we will need along this paper. Section 3 is devoted to establish the compactness result for simple immersed compact constant mean curvature surfaces. First, we shall extend the first eigenvalue estimate to simple immersed surface, i.e.,

> **Theorem 3.1.** *Suppose that Σ is a compact orientable simple immersed hypersurface in a compact $(n+1)-$dimensional orientable Riemannian manifold $\mathcal{N}$. If the Ricci curvature of $\mathcal{N}$ is bounded below by a positive constant k, then*
>
> $$2\lambda_1 \geq k - n \max_{\Sigma} \{|H|\},$$
>
> *where λ_1 is the first nonzero eigenvalue of the Laplacian of Σ and $|H|$ is the absolute value of the mean curvature of Σ.*

And Theorem 3.1 gives us to an uniform area estimate

Theorem 3.2. *Let Σ and $\mathcal{N}$ be the same as in Theorem 3.1. If $n = 2$ and Σ has genus g with area A such that $k - 2\max_\Sigma \{|H|\} > 0$, then*

$$A \leq \frac{16\pi(1+g)}{k - 2\max_\Sigma \{|H|\}}.$$

Therefore, such an area estimate, and the curvature estimate in [8], is the key for obtaining the extension of the Choi-Schoen Compactness Theorem for simple immersed compact constant mean curvature surfaces, specifically:

Theorem 3.5. *Set k and H two positive real numbers so that $k - 2H > 0$. Let $(\mathcal{N}, \gamma)$ be a compact orientable three-manifold with Ricci curvature bounded below by k. Then, the space, $\mathcal{M}_s(H, g)$, of oriented compact simple immersed $H-$surfaces of fixed genus g is compact in the C^j topology for any $j \geq 2$.*

One remark is important at this point. It looks like the above result leds us to use the Degree Theory developed by B. White in [9]. Unfortunately, this is not the case since the space $\mathcal{M}_s(H, g)$ (as defined here) is not an open set of the Banach space of compact immersed constant mean curvature surfaces with fixed topological type.

One might have the following example. Suppose one has a generic one parameter family of metrics $\{\gamma_t\}_{t\in[0,1]}$ so that each γ_t has exactly one constant mean curvature torus $T(t)$, and suppose that $T(t)$ has no Jacobi fields for each $t \in [0, 1]$. Then, the following could happen:

- $T(t)$ is embedded for $t \in [0, 1/2)$.
- $T(1/2)$ is simple immersed but not embedded.
- $T(t)$ is immersed but not simple immersed for $t \in (1/2, 1]$.

Section 4 is devoted to convex embedded constant mean curvature surfaces. We prove a compactness result for such kind of surfaces.

Theorem 4.1. *Let $(\mathcal{N}, \gamma)$ be a $1/4-$pinched orientable simply connected $n-$manifold. Then, the space, $\mathcal{M}_c(H, g)$, of oriented compact embedded convex $H-$hypersurfaces of fixed topological type is compact in the C^j topology for any $j \geq 2$.*

Again, we could try to use the Degree Theory developed in [9], but $\mathcal{M}_c(H, g)$ is not an open set of the Banach space of compact embedded constant mean curvature hypersurfaces.

2 Set up

Let $(\mathcal{N}, \gamma)$ be a Riemannian $(n+1)$-dimensional manifold. Let $\Sigma \subset \mathcal{N}$ be a connected two-sided hypersurface and let N be a unit normal vector field along Σ. We denote the induced metric on Σ by γ as well, and II denotes the second fundamental form, i.e., $II(u,v) := \gamma(\nabla_u N, v)$ where ∇ denotes the Levi-Civita connection of $(\mathcal{N}, \gamma)$. Σ has constant mean curvature H, in short $H-$surface, iff it is a critical point of the functional

$$\mathcal{F} := \text{Area} - 2H\text{Volume}, \tag{2.1}$$

that is,

$$\left.\frac{d}{dt}\right|_{t=0} \mathcal{F}(t) = 0,$$

for all compactly supported variations of Σ

$$\Sigma(t) := \left\{\exp_p(tf(p)N(p)) \ : \ p \in \Sigma\right\}, \ f \in H_0^{1,2}(\Sigma),$$

where $\exp$ is the exponential map on $\mathcal{N}$.

Let $\Sigma \subset \mathcal{N}$ be an $H-$surface and let ψ_t, $t \in (-\epsilon, \epsilon)$, be a smooth family of diffeomorphisms of $\mathcal{N}$ so that ψ_0 is the identity. Assume that

$$\left.\frac{d}{dt}\right|_{t=0} \psi_{t\,|\Sigma} = Y + fN,$$

where Y is a tangential vector field along Σ, N is the normal unit vector field along Σ and f is a smooth function with compact support on Σ. Then

$$\left.\frac{d}{dt}\right|_{t=0} H(t) := L_\Sigma f := \Delta f + (|II|^2 + \text{Ric}(N,N))f, \tag{2.2}$$

where Δ and $|II|^2$ denote the Laplacian with respect to (Σ, γ), the square of the length of the second fundamental form A, and $\text{Ric}(N,N)$ is the Ricci curvature of $\mathcal{N}$ in the direction of N. Here, $H(t)$ denotes the mean curvature of $\Sigma(t)$ for each $t \in (-\epsilon, \epsilon)$.

3 Small $H-$surfaces

The aim of this section is to extend the compactness result of Choi-Schoen [1] to simple immersions with small mean curvature with respect to the Ricci curvature of the ambient manifold. First, we define a simple immersion.

Definition 3.1. *Let $\mathcal{N}$ be a Riemannian manifold of dimension n. We say that an immersed hypersurface $\Sigma \subset \mathcal{N}$ is* **simple** *if there exist a compact connected Riemannian manifold $\Omega \subset \mathcal{N}$ with boundary $\partial\Omega := \Sigma$ and a set $\mathcal{O} \subset \Sigma$ of measure zero so that $\Sigma \setminus \mathcal{O}$ is an embedding.*

To do so, we need to establish area and curvature estimates for simple immersions with constant mean curvature. Once we obtain these estimates, with fixed mean curvature, we obtain the compactness result for simple immersions with small constant mean curvature.

3.1 Area and Curvature Estimate for small $H-$surfaces

In order to get area estimates, we first extend the first eigenvalue estimate of the Laplacian developed in [5] (based on the Choi-Wang estimate for minimal hypersurfaces [2]) to simple immersions.

Theorem 3.1. *Suppose that Σ is a compact orientable simple immersed hypersurface in a compact $(n+1)-$dimensional orientable Riemannian manifold $\mathcal{N}$. If the Ricci curvature of $\mathcal{N}$ is bounded below by a positive constant k, then*

$$2\lambda_1 \geq k - n \max_{\Sigma} \{|H|\},$$

where λ_1 is the first nonzero eigenvalue of the Laplacian of Σ and $|H|$ is the absolute value of the mean curvature of Σ.

Proof. Since Σ is a compact orientable simple immersion, then Σ bounds a compact connected component $\Omega \subset \mathcal{N}$ so that $\partial\Omega = \Sigma$. Consider the inward orientation N along Σ, i.e., the one pointing into Ω, and set

$$\Sigma_t := \left\{\exp_p(tN(p)) \ : \ p \in \Sigma\right\}, \ t \geq 0.$$

Therefore, since Σ is compact, there exists $\delta > 0$ so that Σ_t is an embedding for all $0 < t < \delta$. Hence, from [5, Theorem 2.1], we have

$$2\lambda_1^t > k - n \max_{\Sigma_t} \{|H_t|\}, \ 0 < t < \delta,$$

where λ_1^t is the first nonzero eigenvalue of the Laplacian of Σ_t and $|H_t|$ is the absolutte value of the mean curvature of Σ_t. Thus, taking limit as t goes to zero, we obtain

$$2\lambda_1 \geq k - n \max_{\Sigma} \{|H|\},$$

as desired. □

And, as a direct consequence from [10], we obtain.

Theorem 3.2. *Let Σ and $\mathcal{N}$ be the same as in Theorem 3.1. If $n = 2$ and Σ has genus g with area A such that $k - 2\max_{\Sigma}\{|H|\} > 0$, then*

$$A \leq \frac{16\pi(1+g)}{k - 2\max_{\Sigma}\{|H|\}}.$$

We recover here a curvature estimate for surfaces in three-manifolds that are stationary for a parametric elliptic functional [8, Theorem 0]. We particularize such a general result for constant mean curvature surfaces.

Lemma 3.1 ([8])**.** *Let $\mathcal{N}$ be a three-dimensional Riemannian manifold. Let $q \in \mathcal{N}$ and $r > 0$ such that $B(q,r)$ has compact closure in $\mathcal{N}$, here $B(q,r)$ denotes the geodesic ball in $\mathcal{N}$ centered at q and radius r. Suppose Σ is a compact immersed surface (with possibly empty boundary) in $\mathcal{N}$ such that $q \in \Sigma$ and $\partial\Sigma \cap B(q,r) = \emptyset$.*

There exists $\epsilon_0 > 0$ depending on the geometry of $B(q,r)$ and H so that if

$$\int_{\Sigma \cap B(q,r)} |II|^2 \, dv \leq \epsilon_0 \text{ and } r \leq \epsilon_0,$$

then the following inequality holds:

$$\max_{0 \leq \sigma \leq r} \left\{ \sigma^2 \sup_{B(q,r-\sigma)} \left\{ |II|^2 \right\} \right\} \leq C,$$

where C is a constant depending on the geometry of $B(q,r)$ and H.

3.2 Compactness result for small $H-$surfaces

Now, we establish the aforementioned compactness result:

Theorem 3.3. *Set k and H two positive real numbers so that $k - 2H > 0$. Let $(\mathcal{N}, \gamma)$ be a compact three-manifold with $\mathrm{Ric}_\gamma \geq k$. Let $\Sigma_i \subset \mathcal{N}$ be a sequence of simple immersed $H-$surfaces of fixed topological type, then a subsequence of the $\{\Sigma_i\}$ converges to a smooth compact simple immersed $H-$surface in the sense of $C^{2,\alpha}$ graphs.*

Proof. Area and curvature estimates are given in the previous section. We only need to check that the limit surface Σ, that exists from [8, Theorem 3], is one sheeted and a simple immersion.

Claim 1. Σ is one sheeted, that is, it has multiplicity one.

Proof of Claim 1. Assume some point $p \in \Sigma$ is covered, at least, twice. Then, there are, at least, two sheets of Σ_i approaching to $p \in \Sigma$. Since Σ_i is an orientable simple immersion, we can find two sheets so that the normal vector field at each sheet points in opposite directions. Otherwise we would violate that Σ_i are compact orientable surfaces.

Consider those two sheets $p_{k;i} \in D_{k;i} \subset \Sigma_i$, $k = 1, 2$, and $p_{k;i} \to p$ as $i \to +\infty$. First, observe that, taking a subsequence if necessary, the unit normal $N_{k;i}(p_{k;i})$ of $D_{k;i}$ at $p_{k;i}$ converges to a unit normal vector $N_{k;\infty}$ at p, $k = 1, 2$, so that $N_{1;\infty} = -N_{1;\infty}$. By curvature estimates and for i big enough, each $D_{k;i}$ can be written as a graph (in exponential coordinates) over a uniform disk, $\mathcal{D}(p,\delta) \subset T_p\Sigma$, in the tangent plane of Σ at p and, moreover, $D_{k;i} \to D_{k;\infty}$ as $i \to +\infty$, $D_{k;\infty}$ is also a graph over $\mathcal{D}(p,\delta)$.

Note that $p_{1;i}$ can not apporach to p toward the mean convex side of $D_{1;\infty}$ (recall that this makes sense locally), since otherwise, $D_{1;i}$ and $D_{2;i}$ would intersect for i big enough. Therefore, $D_{1;\infty}$ and $D_{2;\infty}$ intersect at p, the normals point at opposite directions and the mean convex sides (locally at p) are disjoint.

So, we only need to prove that $D_{1;\infty} \cap D_{2;\infty}$ has empty interior. Assume this were not the case then, in a neighborhood of p, the mean curvature of Σ should vanish, a contradiction. □

Clearly, at the self-intersecting points of Σ, no more than two sheet can meet, otherwise we would contradict the fact that Σ_i is a simple immersion for some i big enough. Morever, at those points, the normal vector field a each sheet points at opposite directions. Hence, the measure of self-intersecting points is zero, otherwise we have a contradiction as in Claim 1. □

An inmediate consequence of Theorem 3.3 is the following curvature estimate (see [1, Theorem 2] for details):

Theorem 3.4. *Set k and H two positive real numbers so that $k-2H>0$, and $g \geq 0$ an integer. Let $(\mathcal{N},\gamma)$ be compact three-manifold with $\mathrm{Ric}_g \geq k$. There exists a constant C depending on the ambient geometry of $\mathcal{N}$, H and g such that if Σ is a compact embedded $H-$surface of genus g in $\mathcal{N}$, then*

$$\max_{\Sigma}\{|II|\} \leq C.$$

More generally, we can assert

Theorem 3.5. *Set k and H two positive real numbers so that $k-2H>0$. Let $(\mathcal{N},\gamma)$ be a compact orientable three-manifold with Ricci curvature bounded below by k. Then, the space, $\mathcal{M}_s(H,g)$, of oriented compact simple immersed $H-$surfaces of fixed genus g is compact in the C^j topology for any $j \geq 2$.*

4 Compactness result for convex $H-$surfaces

In this Section, we focus our attention on convex $H-$hypersurfaces immersed in a $1/4-$pinched manifold, that is,

Definition 4.1. *Let $\mathcal{N}$ be a compact $n-$manifold whose sectional curvatures, K_s, are strictly positive. Let κ^+ and κ^- denote the maximum and minimum of the sectional curvatures on $\mathcal{N}$ respectively. Then, we say that $\mathcal{N}$ is $1/4-$pinched if $\kappa^-/\kappa^+ > 1/4$.*

First, we establish an improvement of [4, Lemma 2.2] about embeddedness of a family of closed convex submanifolds in $1/4-$pinched manifolds, which will be the key result for applications in what follows, and it is, in fact, of independent interest. For us, convex means that all the principal curvatures (w.r.t. a fixed orientation) are nonnegative.

Lemma 4.1. *Let D and $\mathcal{N}$ be orientable $n-$manifolds, $\mathcal{N}$ simply connected and D compact with non-empty boundary Σ. Assume $\gamma(t)$ and $h(t)$, $0 \leq t \leq 1$, are continuous families of metrics on D and $\mathcal{N}$ respectively, and each $h(t)$ is $1/4-$pinched.*

Let $f_t : (D, \gamma(t)) \to (\mathcal{N}, h(t))$ be isometric immersions, $0 \leq t \leq 1$, continuous in t. Suppose $f_t(\Sigma) := \Sigma(t)$ has nonnegative principal curvatures for all t (w.r.t. the normal pointing into D).

If f_0 is an embedding, then so is f_t for all t.

Proof. Since D is compact, there exists $\delta > 0$ such that f_t is an embedding for $0 \leq t < \delta$. It suffices to show f_δ is an embedding as well.

Suppose not, let $x, y \in D$ be distinct points such that $f_\delta(x) = f_\delta(y)$. If one of the points $\{x, y\}$ is not on Σ, then one can find open neighborhoods of x and y, U_x and V_y, such that $U_x \cap V_y = \emptyset$ and $f_\delta(U_x) \cap f_\delta(V_y)$ contains an open set of $\mathcal{N}$. But then f_t would not be an embedding for $t < \delta$, t close to δ; a contradiction. Thus, both x and y are on Σ and $f_{\delta|\mathrm{int}D}$ is an embedding.

Consider D with the metric $\gamma(\delta) = f_\delta^*(h(\delta))$. Let β be a minimizing geodesic of $(D, \gamma(\delta))$ joining x to y; β exists because Σ is locally convex and β lies in D (possibly it lies in the boundary of D). Set $l = \mathrm{Length}(\beta)$.

On the one hand, the injectivity radius of $(\mathcal{N}, h(\delta))$, $\mathrm{inj}(\mathcal{N}, h(\delta))$, bounds l from below as (see [3, Theorem C])

$$l/2 \geq \mathrm{inj}(\mathcal{N}, h(\delta)) \geq \frac{\pi}{\sqrt{\kappa^+(\delta)}},$$

here $\kappa^+(\delta)$ denotes the maximum of the sectional curvatures of $(\mathcal{N}, h(\delta))$.

On the other hand, the second-variation-of-length argument of Bonnet-Myers gives an upper bound of l as

$$l \leq \frac{\pi}{\sqrt{\kappa^-(\delta)}},$$

here $\kappa^-(\delta)$ denotes the minimum of the sectional curvatures of $(\mathcal{N}, h(\delta))$.

Thus, joining the above inequalities, we obtain

$$2 \leq \frac{\sqrt{\kappa^+(\delta)}}{\sqrt{\kappa^-(\delta)}},$$

that is

$$\frac{\kappa^-(\delta)}{\kappa^+(\delta)} \leq 1/4,$$

which contradicts the $1/4-$pinched assumption. This proves the Lemma. □

Now, we can state:

Theorem 4.1. *Let $(\mathcal{N},\gamma)$ be a $1/4-$pinched orientable simply connected $n-$manifold. Then, the space, $\mathcal{M}_c(H,g)$, of oriented compact embedded convex $H-$hypersurfaces of fixed topological type is compact in the C^j topology for any $j\geq 2$.*

Proof. Take $\Sigma\in\mathcal{M}_c(H,g)$. Curvature estimates follow from convexity and bounded mean curvature. Actually, $|A|^2\leq (n-1)^2H^2$. Since $(\mathcal{N},\gamma)$ is $1/4-$pinched, by the Gauss equation and convexity, the sectional curvatures of $\Sigma\in\mathcal{M}_c(H,g)$ are uniformly bounded from below by κ^-. Also, by classical Comparisson Theorems, we obtain area estimates.

Let $\{\Sigma_i\}\in\mathcal{M}_c(H,g)$ be a sequence of compact embedded $H-$hypersurfaces of fixed topological type then, following the proof of [8, Theorem 3], a subsequence of the $\{\Sigma_i\}$ converges to a smooth compact convex finite simple immersed $H-$hypersurface in the sense of $C^{2,\alpha}$ graphs. So now, we need to prove that the limit is embedded. This last assertion follows from Lemma 4.1 □

Acknowledgements

The author wishes to express his gratitude to Brian White for his guidance and useful conversations during the preparation of this paper. Also, the author wishes to thank to Joaquín Pérez for his interesting comments and suggestions.

References

[1] H.I. Choi, R. Schoen, *The space of minimal embeddings of a surface in a three-dimensional manifold of positive Ricci curvature*, Invent. Math., **81** (1985), 387–394.

[2] H.I. Choi, A.N. Wang, *A first eigenvalue estimate for minimal hypersurfaces*, J. Diff. Geom., **18** (1983), 559–562.

[3] J. H. Eschenburg, *Local convexity and nonnegative curvature - Gromov's proof of the Sphere Theorem*, Invent. Math., **84** (1986), 507–522.

[4] J.M. Espinar, H. Rosenberg, *When strictly locally convex hypersurfaces are embedded*, To appear in Math. Z.

[5] P. Tung Ho, *A first eigenvalue estimate for embedded hypersurfaces*, Diff. Geom. and its Applications, **26** (2008), 273–276.

[6] H. Rosenberg, G. Smith, *Degree Theory of Immersed Hypersurfaces*. Preprint.

[7] W. Rossman, N. Sultana, *Morse Index of Constant Mean Curvature Tori of Revolution in the $3-$sphere*, Illinois J. Math., **51**, no 4 (2007), 1329–1340.

[8] B. White, *Curvature estimates and compactness theorems in $3-$manifolds for surfaces that are stationary for parametric elliptic functionals*, Invent. Math., **88** (1987), 243–256.

[9] B. White, *The Space of Minimal Submanifolds for Varying Riemannian Metrics*, Indiana Univ. Math. J., **40** no 1 (1991), 161–200.

[10] S.T. Yau, *Seminar on Differential Geometry*, Annals of Math. Studies, **102**, Princeton Univ. Press, Princeton, NJ, 1982.