

Logarithmic and absolute-value like properties of $\pi(x)$ with applications

Boris B. Benyaminov

*6 Moa St, Belmont, North Shore,
Auckland, Postal Code: 0622, New Zealand
boris.b.b@hotmail.com*

Abstract

We derive properties of $\pi(x)$ reminiscent of those of the logarithm and absolute value functions. Two of these properties are similar to the relations defining the linearity of a function. Several applications of these properties of $\pi(x)$ are also examined.

Keywords: prime-counting function, properties, second Hardy-Littlewood conjecture
Mathematics Subject Classification 2010: 11A41, 11M26.

1 Introduction

The prime-counting function $\pi(x) : \mathbb{R} \rightarrow \mathbb{N}$ counts the number of primes less than or equal to x . This function has all the properties of the following functions (listed below explicitly):

1. the logarithm function, $\ln(x)$:

$$\begin{aligned}
 \ln(xy) &= \ln(x) + \ln(y), \quad x, y > 0, \\
 \ln\left(\prod_{i=1}^n x_i\right) &= \sum_{i=1}^n \ln(x_i), \\
 \ln(x^n) &= n \ln(x), \\
 \ln\left(\frac{x}{y}\right) &= \ln(x) - \ln(y), \quad y \neq 0, \\
 \ln(x^{1/m}) &= \frac{\ln(x)}{m};
 \end{aligned} \tag{1.1}$$

2. the absolute-value function, $|x|$:

$$\begin{aligned}
|x + y| &\leq |x| + |y|, \\
\left| \sum_{i=1}^n x_i \right| &\leq \sum_{i=1}^n |x_i|, \\
|x - y| &\geq |x| - |y|, \quad |x| > |y|, \\
|xy| &= |x| |y|, \\
\left| \prod_{i=1}^n x_i \right| &= \prod_{i=1}^n |x_i|, \\
\left| \frac{x}{y} \right| &= \frac{|x|}{|y|}, \quad y \neq 0;
\end{aligned}$$

3. a linear function, $f(x)$:

$$\begin{aligned}
f(x + y) &= f(x) + f(y), \\
f(kx) &= kf(x).
\end{aligned}$$

One logarithmic property of $\pi(x)$ is known [1], similar to (1.1): for $y \geq 2$, $x \geq 6$, $x \geq y$,

$$\pi(xy) > \pi(x) + \pi(y). \quad (1.2)$$

Let us prove all the other properties listed above.

2 Logarithmic properties

Inequality (1.2) can be generalized.

Theorem 2.1. *For $x_i \geq 2$, $i = 1, 2, \dots, n-1$, $x_n \geq 6$, we have*

$$\pi(x_1 x_2 \dots x_n) > \pi(x_1) + \pi(x_2) + \dots + \pi(x_n). \quad (2.3)$$

Proof. By assumption, $x_1 \geq 2$ and $x_2 x_3 \dots x_n \geq 6$, so from (1.2):

$$\pi(x_1 x_2 \dots x_n) = \pi[x_1(x_2 x_3 \dots x_n)] > \pi(x_1) + \pi(x_2 x_3 \dots x_n).$$

Moreover, since $x_2 \geq 2$ and $x_3 x_4 \dots x_n \geq 6$, we also have

$$\pi(x_1 x_2 \dots x_n) > \pi(x_1) + \pi[x_2(x_3 x_4 \dots x_n)] > \pi(x_1) + \pi(x_2) + \pi(x_3 \dots x_n).$$

Continuing this process, we get

$$\pi(x_1 x_2 \dots x_n) > \pi(x_1) + \pi(x_2) + \dots + \pi(x_{n-1} x_n),$$

where $x_{n-1} \geq 2$ and $x_n \geq 6$. Applying inequality (1.2) to the right-hand-side one last time gives inequality (2.3). $\square$

The next property is derived from (2.3).

Theorem 2.2. *If $x \geq 2$ and $n \geq 1$, then*

$$\pi(x^n) \geq n\pi(x). \quad (2.4)$$

Proof. To begin with, assume $x \geq 6$. We would like to use inequality (2.3). To this end, observe that the conditions of Theorem 2.1 are met: if $x_i \geq 6$ for $i = 1, 2, \dots, n$, then clearly $x_i \geq 2$. Therefore, take $x_1 = x_2 = \dots = x_n = x \geq 6$ in (2.3) and get

$$\pi(x \ x \dots x) > \pi(x) + \pi(x) + \dots + \pi(x),$$

which is precisely (2.4), with equality for $n = 1$.

For $2 \leq x < 6$, (2.4) can be readily checked explicitly. It becomes:

- $x = 2 \Rightarrow \pi(2^n) \geq n$. We have equality for $n = 1, 2$ and inequality for $n \geq 3$.
- $x = 3 \Rightarrow \pi(3^n) \geq 2n$. As above, the equality holds for $n = 1, 2$ and the inequality for $n \geq 3$.
- $x = 4 \Rightarrow \pi(4^n) \geq 2n$. Now we have equality for $n = 1$ and inequality for $n \geq 2$.
- $x = 5 \Rightarrow \pi(5^n) \geq 3n$. Here, also, the equality holds for $n = 1$ and the inequality for $n \geq 2$.

Thus, (2.4) is valid for all $x \geq 2$. □

Theorem 2.3. *For $y \geq 2, x \geq 12$ and $x \geq y$, one has*

$$\pi\left(\frac{x}{y}\right) \leq \pi(x) - \pi(y). \quad (2.5)$$

Proof. Let us write (1.2) as $\pi(uv) > \pi(u) + \pi(v)$, where $v \geq 2, u \geq 6, u \geq v$. Take $uv = x$ and $v = y$, so that $u = x/v = x/y$ and $y = v \geq 2, x = uv \geq 6 \cdot 2 = 12, x \geq y$ (i.e. conditions of Theorem 2.3 are met). Substituting these expressions for u and v , we get

$$\pi(x) > \pi(x/y) + \pi(y),$$

whence

$$\pi\left(\frac{x}{y}\right) < \pi(x) - \pi(y). \quad (2.6)$$

In (2.6) above, we in fact have equality whenever $x = y$ as well as for some $x \neq y$, for example, $x = 126$ and $y = 114$. Thus we arrive at (2.5). □

With the help of (2.4) we can prove a few more logarithmic properties of $\pi(x)$. Let us investigate how inequality (2.4) would change if the power on the left-hand-side was less than one. With this in mind, denote $x^n = z$, $z \geq 2$, so that $x = z^{1/n}$. From (2.4) we have $\pi(z) \geq n\pi(z^{1/n})$, whence

$$\pi(z^{1/n}) \leq \frac{1}{n}\pi(z). \quad (2.7)$$

Inequalities (2.4) and (2.7) can be generalized to fractional powers k/m for the two cases $k/m \geq 1$ and $k/m \leq 1$. The resulting expressions are:

$$\begin{aligned} \pi(x^{k/m}) &\geq \frac{k}{m}\pi(x) \quad \text{for } k \geq m, \\ \pi(x^{k/m}) &\leq \frac{k}{m}\pi(x) \quad \text{for } k \leq m. \end{aligned}$$

3 Absolute-value-like properties

For the proof of the first absolute-value property of $\pi(x)$, analogous to the triangle inequality, we need the concept of the density of primes among naturals. The density of primes no greater than x may be defined as $\rho(x) = \pi(x)/x$. Since it is well-known that $\pi(x) = o(x)$, $\rho(x)$ decreases with increasing x . However, this does not happen monotonically. For example, for all twin primes ρ increases. Hence, for real numbers $x, y \in (2, \infty)$ with $y \leq x < x + y$, the relations between $\rho(y)$, $\rho(x)$, $\rho(x + y)$ cannot be stated in general. It is simple to check that the possible relations may be summarized in the following six options:

1.

$$\rho(x + y) \leq \rho(x) \leq \rho(y),$$

2.

$$\rho(x + y) \leq \rho(y) \leq \rho(x),$$

3.

$$\rho(x) \leq \rho(x + y) \leq \rho(y),$$

4.

$$\rho(y) \leq \rho(x + y) \leq \rho(x),$$

5.

$$\rho(x + y) > \rho(x) > \rho(y),$$

6.

$$\rho(x + y) > \rho(y) > \rho(x).$$

In options five and six the case of equality is excluded because it has already been accounted for in previous possibilities.

Theorem 3.1. *[Second Hardy-Littlewood conjecture] For $x, y \geq 2$, $x \geq y$ one has*

$$\pi(x + y) \leq \pi(x) + \pi(y). \quad (3.8)$$

Proof. First of all, we show that the converse of inequality (3.8), that is

$$\pi(x + y) > \pi(x) + \pi(y), \quad (3.9)$$

is impossible. For example, take $y = 1500$, $x = 2000$, leading to $\pi(3500) > \pi(2000) + \pi(1500)$ or $489 > 303 + 239 = 542$ — clearly false.

Next, in order to prove (3.8) we need to consider each of the six options in turn.

1.

$$\frac{\pi(x + y)}{x + y} \leq \frac{\pi(x)}{x} \leq \frac{\pi(y)}{y}. \quad (3.10)$$

From the right-hand inequality in (3.10) we know that $\frac{y\pi(x)}{x} \leq \pi(y)$. Taking this into account, the left-hand part of (3.10) gives

$$\pi(x + y) \leq \frac{x + y}{x} \pi(x) = \pi(x) + \frac{y\pi(x)}{x} \leq \pi(x) + \pi(y),$$

which is of course (3.8).

2.

$$\frac{\pi(x + y)}{x + y} \leq \frac{\pi(y)}{y} \leq \frac{\pi(x)}{x}. \quad (3.11)$$

From the right-hand inequality in (3.11) we know that $\frac{x\pi(y)}{y} \leq \pi(x)$. Taking this into account, the left-hand part of (3.11) gives

$$\pi(x + y) \leq \frac{x + y}{y} \pi(y) = \pi(y) + \frac{x\pi(y)}{y} \leq \pi(x) + \pi(y),$$

which is of course (3.8).

3.

$$\frac{\pi(x)}{x} \leq \frac{\pi(x + y)}{x + y} \leq \frac{\pi(y)}{y}. \quad (3.12)$$

Examine the right-hand inequality in (3.12): $\frac{\pi(x + y)}{x + y} \leq \frac{\pi(y)}{y}$. Let us prove that also

$$\frac{\pi(x) + \pi(y)}{x + y} \leq \frac{\pi(y)}{y}. \quad (3.13)$$

Inequality (3.13) $\Leftrightarrow y\pi(x) + y\pi(y) \leq x\pi(y) + y\pi(y) \Leftrightarrow \frac{\pi(x)}{x} \leq \frac{\pi(y)}{y}$. The last inequality is valid by one of our initial assumptions. Thus, there are two possibilities:

(a)

$$\frac{\pi(x)}{x} \leq \frac{\pi(x+y)}{x+y} \leq \frac{\pi(x) + \pi(y)}{x+y} \leq \frac{\pi(y)}{y},$$

(b)

$$\frac{\pi(x)}{x} \leq \frac{\pi(x) + \pi(y)}{x+y} \leq \frac{\pi(x+y)}{x+y} \leq \frac{\pi(y)}{y}.$$

In case (a) we still must prove that $\frac{\pi(x)}{x} \leq \frac{\pi(x)+\pi(y)}{x+y}$. This last inequality $\Leftrightarrow x\pi(x) + y\pi(x) \leq x\pi(x) + x\pi(y) \Leftrightarrow \frac{\pi(x)}{x} \leq \frac{\pi(y)}{y}$, which is again true. Therefore the inequalities in case (a) can be considered as either assumed or proved. In either case, the central inequality gives $\pi(x+y) \leq \pi(x) + \pi(y)$, exactly (3.8).

In case (b), all inequalities are either assumed or proved earlier. From the central inequality, we get $\pi(x+y) = \pi(x) + \pi(y)$, as $\pi(x+y) > \pi(x) + \pi(y)$ is the incorrect (3.9).

4.

$$\frac{\pi(y)}{y} \leq \frac{\pi(x+y)}{x+y} \leq \frac{\pi(x)}{x}.$$

Examine the right-hand inequality: $\frac{\pi(x+y)}{x+y} \leq \frac{\pi(x)}{x}$. We prove that also

$$\frac{\pi(x) + \pi(y)}{x+y} \leq \frac{\pi(x)}{x}. \quad (3.14)$$

Inequality (3.14) $\Leftrightarrow x\pi(x) + x\pi(y) \leq x\pi(x) + y\pi(x) \Leftrightarrow \frac{\pi(y)}{y} \leq \frac{\pi(x)}{x}$, which is assumed. Again, we have two options:

(a)

$$\frac{\pi(y)}{y} \leq \frac{\pi(x+y)}{x+y} \leq \frac{\pi(x) + \pi(y)}{x+y} \leq \frac{\pi(x)}{x},$$

(b)

$$\frac{\pi(y)}{y} \leq \frac{\pi(x) + \pi(y)}{x+y} \leq \frac{\pi(x+y)}{x+y} \leq \frac{\pi(x)}{x}.$$

In case (a) we still need to confirm that $\frac{\pi(y)}{y} \leq \frac{\pi(x)+\pi(y)}{x+y}$. This last inequality $\Leftrightarrow x\pi(y) + y\pi(y) \leq y\pi(x) + y\pi(y) \Leftrightarrow \frac{\pi(y)}{y} \leq \frac{\pi(x)}{x}$, which was initially assumed. Therefore the inequalities in case (a) can be considered as either assumed or proved. The central inequality gives $\pi(x+y) \leq \pi(x) + \pi(y)$, precisely (3.8).

In case (b), all inequalities are valid. From the central inequality, we get $\pi(x+y) = \pi(x) + \pi(y)$, as $\pi(x+y) > \pi(x) + \pi(y)$ is the incorrect (3.9).

5.

$$\frac{\pi(x+y)}{x+y} > \frac{\pi(x)}{x} > \frac{\pi(y)}{y}.$$

This option is not possible as it leads to (3.9). Indeed, since $\frac{y\pi(x)}{x} > \pi(y)$, we have $\frac{\pi(x+y)}{x+y} > \frac{\pi(x)}{x} \Leftrightarrow \pi(x+y) > \frac{x+y}{x}\pi(x) = \pi(x) + \frac{y\pi(x)}{x} > \pi(x) + \pi(y)$, which is (3.9).

6.

$$\frac{\pi(x+y)}{x+y} > \frac{\pi(y)}{y} > \frac{\pi(x)}{x}.$$

This case is also not possible because it implies (3.9) is true. The proof is analogous to the previous option.

Thus, (3.8) is proved. □

With this result, it is trivial to prove that for any $x_{1,2,3} \geq 2$

$$\pi(x_1 + x_2 + x_3) \leq \pi(x_1) + \pi(x_2) + \pi(x_3).$$

Indeed, since $x_1 + x_2 \geq 4 > 2$, by (3.8) we have

$$\pi(x_1 + x_2 + x_3) = \pi[(x_1 + x_2) + x_3] \leq \pi(x_1 + x_2) + \pi(x_3) \leq \pi(x_1) + \pi(x_2) + \pi(x_3).$$

Analogously, one can prove a generalized version of the above for a sum of k terms. Let $k < \infty$, $x_i \geq 2$, $i = 1, 2, \dots, k$, then

$$\pi\left(\sum_{i=1}^k x_i\right) \leq \sum_{i=1}^k \pi(x_i). \quad (3.15)$$

Furthermore, if we let $k \geq 1$ and take $x_1 = x_2 = \dots = x_k = x \geq 2$ in (3.15), we obtain:

$$\pi(kx) \leq k\pi(x), \quad k \geq 1. \quad (3.16)$$

Inequalities (3.15) and (3.16) demonstrate that $\pi(x)$ has properties similar to the definition of linearity. Finally, with the help of (3.15) and (3.16), it is easy to show that

$$\pi\left(\sum_{i=1}^n k_i x_i\right) \leq \sum_{i=1}^n k_i \pi(x_i),$$

where $x_i \geq 2$, $k_i \geq 1$, $k_i x_i \geq 2$ and $i = 1, 2, \dots, n$.

We now proceed to prove the second absolute-value property of $\pi(x)$, in this case similar to the *reverse* triangle inequality.

Theorem 3.2. *For $y \geq 2$, $x \geq 4$, $x \geq y$, one has*

$$\pi(x - y) \geq \pi(x) - \pi(y). \quad (3.17)$$

Proof. We write inequality (3.8) as $\pi(u+v) \leq \pi(u) + \pi(v)$ where $u, v \geq 2$. Denote $u+v = x$ and $v = y$ so that $u = x - v = x - y$ with $y = v \geq 2$, $x = u + v \geq 4$ and $x \geq y$ (i.e. the conditions of Theorem 3.2 are met). Substituting these expressions for u and v , we get

$$\pi(x) \leq \pi(x-y) + \pi(y),$$

from which (3.17) follows. $\square$

Theorem 3.3. *For $x, y \geq 8$, we have*

$$\pi(xy) > \pi(x)\pi(y). \quad (3.18)$$

Proof. On account of (3.17), we obtain

$$\pi \left[\left(x - \frac{y}{2} \right)^2 \right] = \pi \left(x^2 - xy + \frac{y^2}{4} \right) = \pi \left[\left(x^2 + \frac{y^2}{4} \right) - xy \right] \geq \pi \left(x^2 + \frac{y^2}{4} \right) - \pi(xy),$$

and from here, we see that

$$\pi(xy) \geq \pi \left(x^2 + \frac{y^2}{4} \right) - \pi \left[\left(x - \frac{y}{2} \right)^2 \right].$$

Keeping in mind inequality (3.18), we require

$$\pi(xy) > \pi \left(x^2 + \frac{y^2}{4} \right) - \pi \left[\left(x - \frac{y}{2} \right)^2 \right] \geq \pi(x)\pi(y). \quad (3.19)$$

Then, all values of x and y that satisfy the right-hand side inequality of (3.19) will also clearly satisfy (3.18). However, the right-hand side part of (3.19) contains two variables and is therefore difficult to solve. Nonetheless, we can take $x = y = q$ so that the inequality becomes

$$\pi \left(\frac{5q^2}{4} \right) \geq \pi \left(\frac{q^2}{4} \right) + [\pi(q)]^2. \quad (3.20)$$

Now, (3.20) is satisfied for all $q \geq 8$ and therefore, (3.18) is satisfied for all $x, y \geq 8$. $\square$

Inequality (3.18) may be readily generalized to the case of a product of $n < \infty$ terms. Let $x_i \geq 8$, $i = 1, 2, \dots, n$, then

$$\pi \left(\prod_{i=1}^n x_i \right) > \prod_{i=1}^n \pi(x_i).$$

For the special case when $x_1 = x_2 = \dots = x_n = x \geq 8$, the above becomes

$$\pi(x^n) > [\pi(x)]^n.$$

Finally, let us investigate the final absolute-value property of $\pi(x)$.

Theorem 3.4. For $y \geq 8$, $x \geq 64$, $x \geq y$, we have

$$\pi\left(\frac{x}{y}\right) < \frac{\pi(x)}{\pi(y)}. \quad (3.21)$$

Proof. We write (3.18) as $\pi(uv) > \pi(u)\pi(v)$ with $u, v \geq 8$. Denote $uv = x$ and $v = y$ so that $u = x/v = x/y$. The constraints for x and y are: $y = v \geq 8$, $x = uv \geq 64$ and $x \geq y$ since $uv \geq v \Leftrightarrow u \geq 1$, which is true since $u \geq 8$. Now substitute these expressions for u and v to get $\pi(x) > \pi(x/y)\pi(y)$ — none other than (3.21). $\square$

4 Applications

Consider the implications of inequality (2.5). Various proofs are known for Bertrand's postulate [2], which states that for all $n \geq 1$ there exists a prime number p such that $n < p < 2n$ or, in other words:

$$\pi(2n) - \pi(n) \geq 1. \quad (4.22)$$

However, by far the simplest proof can be achieved through the use of (2.5). To this end, let $y = n \geq 2$ and $x = 2n \geq 12$ or $n \geq 6$, $x \geq y$. Suppose for the moment that $n \geq 6$. Then, from (2.5), we get

$$\pi(2n) - \pi(n) \geq \pi\left(\frac{2n}{n}\right) = \pi(2) = 1,$$

which is exactly inequality (4.22). For $n \leq 5$ the intervals $(n, 2n)$ in question are: $(1, 2)$, $(2, 4)$, $(3, 6)$, $(4, 8)$ and $(5, 10)$. It is obvious that each of these contains at least one prime number.

With the help of property (2.5), it is equally simple to show that there are at least two prime numbers in the interval $(n, 3n)$, $n \geq 1$. To obtain this result, take $y = n \geq 2$ and $x = 3n \geq 12$ [i.e. $n \geq 4$] in (2.5). We get

$$\pi(3n) - \pi(n) \geq \pi\left(\frac{3n}{n}\right) = \pi(3) = 2.$$

For $n < 4$ we have the intervals $(n, 3n)$: $(1, 3)$, $(2, 6)$ and $(3, 9)$, all containing at least two primes.

Analogously, we prove that for $n \geq 1$, the interval $(n, 5n)$ contains no less than three primes. Once again, take $y = n \geq 2$ while $x = 5n \geq 12$ [i.e. $n \geq 3$] and from (2.5), obtain

$$\pi(5n) - \pi(n) \geq \pi\left(\frac{5n}{n}\right) = \pi(5) = 3.$$

For $n < 3$ we have the intervals $(n, 5n)$: $(1, 5)$ and $(2, 10)$, which contain at least three primes.

Exactly in the same manner, we can prove that the interval $(n, 7n)$, $n \geq 1$ contains at

least four primes, and so on. Generalizing, the interval (n, kn) with $k \geq 2$ contains no less than $\pi(k)$ primes.

Let us consider the application of the absolute-value property (3.17). It can be used to bound from above the number of primes in the interval (y, x) . For example, if we wish to estimate the number of primes in the interval $(10^7, 10^8)$, then from (3.17) we get:

$$\pi(10^8) - \pi(10^7) \leq \pi(10^8 - 10^7) = \pi(9 \cdot 10^7) = 5216954.$$

Actually, the given interval contains 5096879 primes. Let us examine some intervals of special interest.

1. Bertrand's interval: $(n, 2n)$, $n = 2, 3, \dots$

By virtue of inequality (3.17), we have

$$\pi(2n) - \pi(n) \leq \pi(2n - n) = \pi(n). \quad (4.23)$$

In particular, from (4.23) it follows that $\pi(2n) \leq 2\pi(n)$, which we could have obtained from (3.16) with $k = 2$. For the interval in question, we get the bounds

$$1 \leq \pi(2n) - \pi(n) \leq \pi(n), \quad n \geq 2.$$

2. Legendre's interval: $(n^2, (n+1)^2)$, $n = 1, 2, 3, \dots$

From (3.17) we find the upper bound

$$\pi((n+1)^2) - \pi(n^2) \leq \pi(n^2 + 2n + 1 - n^2) = \pi(2n + 1).$$

For instance, if $n = 2010$, i.e. the interval is $(2010^2, 2011^2)$, we get

$$\pi(2011^2) - \pi(2010^2) \leq \pi(4021) = 556.$$

This means that in the given interval there are no more than 556 primes.

3. The interval $(x, x + a)$.

Again from (3.17), we have

$$\pi(x + a) - \pi(x) \leq \pi(x + a - x) = \pi(a).$$

Observe that the right-hand side is independent of x , that is, an interval of length a contains no more than $\pi(a)$ primes, irrespective of its position along the real line. If we consider the interval $(y, x + y)$ of length x , we get $\pi(x + y) - \pi(y) \leq \pi(x)$, which is inequality (3.8).

In fact, property (3.17) allows one to estimate the number of primes in more diverse intervals such as $(y, y + \sqrt{y})$, $(n^3, (n+1)^3)$ etc.

Acknowledgements

The author would like to thank Sophie S. Shamaïlov for translating and typing the article.

References

- [1] H. Ishikawa, *Über die Verteilung der Primzahlen*, Sci. Rep. Tokyo Bunrika Daigaku (A) **2**, 27-40 (1934).
- [2] E. Trost, *Primzahlen*, Verlag Birkhäuser, Basel-Stuttgart (1953).