

ASYMPTOTIC ESTIMATES ON THE ERDŐS-STRAUS CONJECTURE

ELIAS RIOS

ABSTRACT. In this paper analyzes *The Erdős-Straus conjecture* asserts that $f(n) > 0$ for every $n \geq 2$, where $f(n)$ indicates the number of solutions to the Diophantine Equation $\frac{4}{n} = \frac{1}{n_1} + \frac{1}{n_2} + \frac{1}{n_3}$. We show that there exists a function $G(p)$ to be a boundary asymptotic of $\sum_{p \leq N} f_I(p)$, which will have an associated error. We analyze the case when n is a prime number, this was separately developed by Terence Tao [8] and Jia [1], [2].

1. INTRODUCTION

Let $n \in \mathbb{N} = \{1, 2, \dots\}$ natural number, let $f(n)$ denote the number of solutions $(n_1; n_2; n_3) \in \mathbb{N}^3$ to the diophantine equation:

$$(1.1) \quad \frac{4}{n} = \frac{1}{n_1} + \frac{1}{n_2} + \frac{1}{n_3}$$

The *Erdős-Straus conjecture* is an open problem in number theory. Recently Terence Tao [8] studied situation in which n is a prime number. Choahua Jia [1],[2], work on the same basis, by replacing n by a prime number p . Let p a prime number, $f(p)$ can be split into $f_I(p)$ and $f_{II}(p)$. In the first instance Terence Tao [8] establish that

$$(1.2) \quad \sum_{p \leq N} f(p) \ll N \log^2 N \log \log N$$

and

$$(1.3) \quad \sum_{p < N} f_I(p) \ll N \exp\left(\frac{c \log N}{\log \log N}\right)$$

other result for Jia [2]

$$(1.4) \quad \sum_{p < x} f_I(p) \ll x \log^5 x \log \log^2 x.$$

We suppose $G(p)$ a bound asymptotic, such that $\sum_{p \leq N} f(p) \sim G(p)$, then

$$(1.5) \quad \lim_{p \rightarrow \infty} \frac{\sum_{p < x} f_I(p)}{G(p)} = 1$$

therefore $\sum_{p < x} f_I(p) - G(p) \rightarrow 0$, and let $\varepsilon(N) \geq 0$ which we can pose the following inequality

Key words and phrases. The Erdős-Straus Conjecture, Diophantine Equations, Prime Numbers

$$\sum_{p < N} f_I(p) - G(p) \leq \varepsilon(N)$$

This asymptotic limit indicates the distance between it, so one can speak of error to the difference of the two. Therefore we can pose the following theorem

Theorem 1.1. *Let $G(p) \geq 0$ exists a function $\varepsilon(N) \geq 0$ such that*

$$(1.6) \quad \sum_{p < N} f_I(p) - G(p) \leq \varepsilon(N)$$

Remark 1.2. An example we can see according to inequality is the following, for N sufficiently large

$$\begin{aligned} \sum_{p < N} f_I(p) - G(p) &\leq \log N \\ \sum_{p < N} f_I(p) - G(p) &\leq N \log N \end{aligned}$$

We could also say that

$$\sum_{p < N} f_I(p) \ll N \log^N N.$$

2. ESTIMATE OF $G(p)$

We know that $G(p)$ is a bound of $f_I(p)$. The question that we would make immediately what would be the nature of $G(p)$. As we know the estimate of this can be complicated algebraically to find a valid estimate. According inequality (1.6) we can state the following inequality

$$(2.1) \quad \sum_{p < N} f_I(p) \leq G(p) + \varepsilon(N)$$

we propose the following equation

$$(2.2) \quad \chi(N, p) = G(p) + \varepsilon(N)$$

therefore

$$(2.3) \quad \sum_{p < N} f_I(p) \leq \chi(N, p)$$

the expression $\chi(N, p)$ indicates the asymptote more error.

By Theorem proposed by Jia [1], we have

$$(2.4) \quad \chi(N, p) = N \log^5 N \log \log^2 N$$

If we replace and get solving for

$$(2.5) \quad G(p) = N \log^5 N \log \log^2 N - \varepsilon(N)$$

in the remark made we proposed that $\varepsilon(N) = \log N$, so we can replace in equation

(2.5)

$$(2.6) \quad G(p) = N \log^5 N \log \log^2 N - \log N$$

therefore, this can be a value of $G(p)$. Also menara could find many ways to determine $G(p)$.

3. SOME EQUALITIES AND INEQUALITIES

According Von Mangoldt function can pose the following inequality with the following equation

$$\sum_{m|N} \Lambda(m) = \log N.$$

therefore

$$(3.1) \quad \sum_{p < N} f_I(p) - G(p) \leq \sum_{m|N} \Lambda(m)$$

For every real number $X \geq 2$, we write

$$(3.2) \quad \pi(X) = \sum_{p \leq X} 1$$

so that $\pi(X)$ denotes the number of primes in the interval $[2, X]$. This function has been studied extensively by number theorists, and attempts to study it in depth have led to major developments in other important branches of mathematics. As can be expected, many conjectures concerning the distribution of primes were made based purely on numerical evidence, including the celebrated Prime number theorem, proved in 1896 by Hadamard and de la Vallée Poussin, that

$$(3.3) \quad \lim_{X \rightarrow \infty} \frac{\pi(X) \log X}{X}$$

we can establish a relationship with the equation (1.5)

$$(3.4) \quad \frac{\sum_{p < x} f_I(p)}{G(p)} \sim \frac{\pi(X) \log X}{X}.$$

Some of the questions that I will open, in future articles try to show is that if you fulfills

$$G(p) \equiv 0 \pmod{N}$$

or if there is a function $\eta(p)$ dependent of p such that the following congruence is

$$G(p) \equiv \eta(p) \pmod{N}$$

and if there fulfill that $N|G(p)$

REFERENCES

1. Chaohua Jia, *A Note on Terence Tao's Paper "On the Number of Solutions to $\frac{4}{p} = \frac{1}{n_1} + \frac{1}{n_2} + \frac{1}{n_3}$ "*, available at <http://arxiv.org/abs/1107.5394>
2. Chaohua Jia, *On the Estimate for a Mean Value Relative to $\frac{4}{p} = \frac{1}{n_1} + \frac{1}{n_2} + \frac{1}{n_3}$* , available at <http://arxiv.org/abs/1107.6039>
3. Li Delang, *On the Equation $4/n=1/x+1/y+1/z$* , Journal of Number Theory, 13, 485-494, (1981).
4. H. Cohn, *Advanced number theory*, Dover Publications, Inc New York, 1965.
5. H. Iwaniec, E. Kowalski, *Analytic number theory* American Mathematical Society Colloquium Publications, 53. American Mathematical Society, Providence, RI, 2004.
6. H. S. Zuckerman, I. Niven, *Introduction to number theory*. Editorial Limusa, 1976.
7. G.H. Hardy, E. M. Wright *An Introduction to the Theory of Number*, Fourth Edition, Oxford at the Clarendon Press, (1960).
8. Terence Tao, *On the number of solutions to $\frac{4}{p} = \frac{1}{n_1} + \frac{1}{n_2} + \frac{1}{n_3}$* , available at <http://arxiv.org/abs/1107.1010>
9. Tom A. Apostol, *Introduction to analytic number theory*, Undergraduate Texts in Mathematics, Springer-Verlag, (1976).
10. W. W. L. Chen, *Elementary and Analytic Number Theory*, University of London, (1981).
11. W. Edwin Clark, *Elementary Number Theory*, Department of Mathematics, University of South Florida, (2003).
12. Yong-Gao Chen, Christian Elsholtz, Li-Li Jiang, *Egyptian fractions with restrictions*, <http://arxiv.org/abs/1108.6118>

UNIVERSIDAD TECNOLÓGICA NACIONAL, FACULTAD REGIONAL AVELLANEDA, BUENOS AIRES,
ARGENTINA

E-mail address: `erios@fra.utn.edu.ar`