arXiv:1507.06389v1 [math.CO] 23 Jul 2015

A formula for the number of the spanning trees of line graphs[1]

Helin Gong, Xian'an Jin[2]
School of Mathematical Sciences
Xiamen University

Abstract

Let $G = (V, E)$ be a loopless graph and $\mathcal{T}(G)$ be the set of all spanning trees of G. Let $L(G)$ be the line graph of the graph G and $t(L(G))$ be the number of spanning trees of $L(G)$. Then, by using techniques from electrical networks, we obtain the following formula:

$$t(L(G)) = \frac{1}{\prod_{v\in V} d^2(v)} \sum_{T\subseteq\mathcal{T}(G)} [\prod_{e=xy\in T} d(x)d(y)][\prod_{e=uv\in E\setminus T} [d(u)+d(v)]].$$

As a result, we provide a very simple and different proof of the formula on the number of spanning trees of some irregular line graphs, and give a positive answer to a conjecture proposed by Yan [J. Combin. Theory Ser. A 120 (2013) no. 7, 1642-1648]. By applying our formula we also derive the number of spanning trees of circulant line graphs.

Keywords: electrical network, spanning tree, line graph, regular and irregular graph, subdivision, circulant graph
AMS MSC 2010: 05C30, 05C76, 05C05, 05C07, 05C22

1. Introduction

Throughout this paper we consider connected graphs that allow parallel edges but have no loops. We always assume that $G = (V, E)$ is a graph with

[1]This work was supported by the National Natural Science Foundation of China (No. 11271307).
[2]E-mail:xajin@xmu.edu.cn

n vertices and m edges, i.e. $|V| = n, |E| = m$. For $v \in V$, we denote by $d(v)$ the degree of v in G. For $e \in E$ with two end-vertices u and v, we shall write $e = uv$. The *line graph* of G, denoted by $L(G)$, is defined as the graph with $V(L(G)) = E$ such that e and f in E is jointed by k edges in $L(G)$ if and only if e and f have k common end-vertices, where $k = 0, 1$ or 2.

Let $t(G)$ and $t(L(G))$ be the number of spanning trees of G and $L(G)$, respectively. One of classic theorems on graph spectra, established originally by Vahovskii[9] and later reported by Kelmans [2] and Cvetković, Doob and Sachs [4], respectively, explains the relation between $t(G)$ and $t(L(G))$ as follows.

$$t(L(G)) = 2^{m-n+1} r^{m-n-1} t(G), \tag{1}$$

where G is a r-regular graph. The *subdivision* of G, denoted by $S(G)$, is the graph obtained from G by inserting a vertex to each edge of G. In [5], Zhang, Chen and Chen studied the spectra of the clique-inserted graphs, i.e. line graphs of subdivisions of graphs. As a corollary, they obtained:

$$t(L(S(G))) = r^{m-n-1}(r+2)^{m-n+1} t(G), \tag{2}$$

where G is a r-regular graph.

To generalize Eq. (1) from regular graphs to irregular graphs, the *revision* of G, denoted by $R(G)$, was introduced by Yan in [6]. It is defined as the graph obtained from G by attaching $\Delta - d(v)$ pendent edges to each vertex v of G, where Δ is the maximum degree of G. Using a factorization theorem in [7] and the famous Matrix-Tree Theorem, Yan [6] proved:

$$t(L(R(G))) = 2^{m-n+1} \Delta^{n\Delta-m-n-1} t(G). \tag{3}$$

Obviously $R(G) = G$ if G is regular and in this case Eq. (3) reduces to Eq. (1). Motivated by Eq. (2) Yan further conjectured:

$$t(L(S(R(G)))) = \Delta^{n\Delta-m-n-1}(\Delta+2)^{m-n+1} t(G). \tag{4}$$

In fact, Eq. (4) reduces to Eq. (2) if G is regular.

Let $\mathcal{T}(G)$ be the set of all spanning trees of G. In this paper, by using techniques from electrical network, we obtain the following formula:

$$t(L(G)) = \frac{1}{\prod_{v\in V} d^2(v)} \sum_{T\subseteq\mathcal{T}(G)} \prod_{e=xy\in T} d(x)d(y) \prod_{e=uv\in E\setminus T} [d(u)+d(v)]. \tag{5}$$

Such techniques have been used to compute various polynomials of graphs and links, see [13, 12, 14, 11]. As a direct consequence, we provide a very simple and different proof of Eq. (3) and give a positive answer to the conjecture proposed by Yan, i.e. Eq. (4).

The enumeration of spanning trees of circulant graphs is well studied, see, for example, [15]. Using our formula we derive a formula on the number of spanning trees of circulant line graphs.

2. Techniques from electrical networks

A weighted graph is a graph G together with a weight function $\omega : E \to [0, \infty)$. Let G be a weighted graph. The *weight* $w(T)$ of a spanning tree T in G is defined as the product of weights of edges in T, i.e.

$$w(T) = \prod_{e \in T} w(e). \tag{6}$$

Let

$$\tau(G) = \sum_{T} w(T), \tag{7}$$

where the summation is over all spanning trees of G. If $\omega(e) = 1$ for all $e \in E$, then $\tau(G) = t(G)$.

An electrical network can be regarded as a weighted graph in which the weights are the conductances of the respective edges. Following [10], two weighted graphs (networks) G and H are called *electrically equivalent* with respect to $D \subseteq V(G) \cap V(H)$, if they cannot be distinguished by applying voltages to D and measuring the resulting currents on D. In [10], Teufl and Wagner explained the effect of electrically equivalent transformations on the sum of weights of spanning trees.

Theorem 2.1 (Theorem 10,[10]). *Let X be a weighted graph that can be partitioned into two edge-disjoint subgraphs G and H (inherit weights in an obvious way) such that $V(X) = V(G) \cup V(H)$, $V(G) \cap V(H) = S$. We assume that $\tau(X) \neq 0$ and $\tau(H) \neq 0$. Let H' be a weighted graph satisfying $E(G) \cap E(H') = \emptyset$ and $V(G) \cap V(H') = S$. Furthermore, suppose H and H' are electrically equivalent with respect to S. Finally, set $X' = G \cup H'$. Then, the following formula holds:*

$$\frac{\tau(X')}{\tau(X)} = \frac{\tau(H')}{\tau(H)}. \tag{8}$$

We shall focus on two kinds of electrically equivalent transformations below that are well-known in the analysis of an electrical network. Applying Theorem 2.1, we can obtain:

- **Serial edges-single edge transformation**: If two serial edges with weights a and b (H) are merged into a single edge with weight $\frac{ab}{a+b}$ (H'). Then

$$\tau(X) = (a+b)\tau(X'). \tag{9}$$

- **Mesh-star transformation**: If X' is obtained from X by replacing a mesh subgraph K_c $(c \geq 2)$ (H) of X with weight w on its every edge by an electrically equivalent star subgraph $K_{1,c}$ (H') with weight cw on its every edge, then

$$\tau(X) = \frac{1}{c^2 w}\tau(X'). \tag{10}$$

  The mesh-star transformation does not always exist, for the condition under which a mesh network has an equivalent star network is well-known, we refer the readers to [8] for details.

3. Main result and its proof

Theorem 3.1. *Let G be a loopless graph, and let $\mathcal{T}(G)$ be the set of spanning trees of G. Then*

$$t(L(G)) = \frac{1}{\prod_{v\in V} d^2(v)} \sum_{T\subseteq\mathcal{T}(G)} [\prod_{e=xy\in T} d(x)d(y)][\prod_{e=uv\in E\setminus T} [d(u)+d(v)]]. \tag{11}$$

Proof.

We use G' to denote the graph obtained from G by subdividing all pendent edges in G. In the case that G has no pendent edge, $G' = G$. It is easy to see that

$$t(L(G)) = t(L(G')). \tag{12}$$

Let X be the weighted graph consisting of G and the weight function ω such that $\omega(e) = \frac{d(u)d(v)}{d(u)+d(v)}$ for $e = uv \in E$. Let X' be the weighted graph consisting of $L(G')$ and the weight function $\omega' : E(L(G')) \to \{1\}$. Let X''

be the weighted graph consisting of $S(G)$ ($V(S(G)) = V(G) \cup S$) and the weight function ω'' such that $\omega''(e) = d(u_e)$ for $e = u_e v_e \in E(S(G))$ with $v_e \in S$. Then X', X'', X are electrical equivalent to each other, i.e. By Eq. (10) and Eq. (9), we obtain:

$$t(L(G')) = \tau(X'') \cdot \prod_{v \in V} \frac{1}{d^2(v)} \tag{13}$$

and

$$\tau(X'') = \tau(X) \cdot \textstyle\prod_{e=uv \in E}[d(u) + d(v)]. \tag{14}$$

Combining Eqs. (12), (13) and (14), we have

$$\begin{aligned} t(L(G)) &= t(L(G')) \\ &= \prod_{v \in V} \frac{1}{d^2(v)} \cdot \tau(X'') \\ &= \prod_{v \in V} \frac{1}{d^2(v)} \cdot \prod_{e=uv \in E} [d(u) + d(v)] \cdot \tau(X) \\ &= \frac{\prod_{e=uv \in E}[d(u) + d(v)]}{\prod_{v \in V} d^2(v)} \cdot \sum_{T \subseteq \mathcal{T}(G)} \prod_{e=uv \in T} \frac{d(u)d(v)}{d(u) + d(v)} \qquad (15) \\ &= \frac{1}{\prod_{v \in V} d^2(v)} \sum_{T \subseteq \mathcal{T}(G)} \prod_{e=xy \in T} d(x)d(y) \prod_{e=uv \in E \setminus T} [d(u) + d(v)]. \end{aligned}$$

This completes the proof of Theorem 11. □

4. Applications

For convenience in this section we shall use Eq. (15).

Note that $R(G)$ has n vertices of degree Δ and $n\Delta - 2m$ vertices of degree 1, m edges whose two end-vertices are of both degree Δ and $n\Delta - 2m$ pendent edges whose two end-vertices are of degree Δ and 1. So we have

$$\begin{aligned} t(L(R(G))) &= \frac{[2\Delta]^m[\Delta + 1]^{n\Delta - 2m}}{[\Delta^2]^n} \sum_{T \subseteq \mathcal{T}(R(G))} [\frac{\Delta^2}{2\Delta}]^{n-1} [\frac{\Delta}{\Delta + 1}]^{n\Delta - 2m} \\ &= 2^{m-n+1} \Delta^{n\Delta - m - n - 1} t(R(G)) \\ &= 2^{m-n+1} \Delta^{n\Delta - m - n - 1} t(G). \end{aligned}$$

Thus Eq. (3) is established.

Now we prove Eq. (4). Note that $S(R(G))$ has n vertices of degree Δ, $n\Delta - m$ vertices of degree 2 and $n\Delta - 2m$ vertices of degree 1. Moreover, if $e \in E(S(R(G)))$ is not a pendent edge then the degrees of its two end-vertices are 2 and Δ and otherwise,the degrees of two end-vertices of e are 2 and 1. The numbers of pendent edges and non-pendent edges of $S(R(G))$ are $n\Delta - 2m$ and $n\Delta$, respectively. So

$$\begin{aligned} t(L(S(R(G)))) &= \frac{[\Delta+2]^{n\Delta}[2+1]^{n\Delta-2m}}{[\Delta^2]^n[2^2]^{n\Delta-m}} \sum_{T\subseteq\mathcal{T}(S(R(G)))} [\frac{2\Delta}{\Delta+2}]^{n+n\Delta-m-1} [\frac{2}{2+1}]^{n\Delta-2m} \\ &= 2^{-m+n-1}\Delta^{n\Delta-m-n-1}(\Delta+2)^{m-n+1}t(S(R(G))) \\ &= 2^{-m+n-1}\Delta^{n\Delta-m-n-1}(\Delta+2)^{m-n+1}t(S(G)) \\ &= \Delta^{n\Delta-m-n-1}(\Delta+2)^{m-n+1}t(G), \end{aligned}$$

The last equation follows from $t(S(G)) = 2^{m-n+1}t(G)$ (see Exercise 2.4.5 b in [1]).

Finally we compute the number of spanning trees of circulant line graphs. A complete characterization of all circulant line graphs was given by Brown and Hoshino in [3]. They proved that if $L(G)$ is circulant then G must be K_4, C_n, or $K_{a,b}$ with $\gcd(a,b) = 1$.

Theorem 4.1. *Let $G = L(K_{a,b})$ be a circulant line graph. Then*

$$t(G) = a^{a-2}b^{b-2}[a+b]^{ab-a-b+1}.$$

Proof. It is well known that $t(K_{a,b}) = a^{b-1}b^{a-1}$. By Eq. (15), we have

$$\begin{aligned} t(L(K_{a,b})) &= \frac{[a+b]^{ab}}{[a^2]^b[b^2]^a} \sum_{T\subseteq\mathcal{T}(K_{a,b})} [\frac{ab}{a+b}]^{a+b-1} \\ &= a^{a-b-1}b^{b-a-1}[a+b]^{ab-a-b+1}t(K_{a,b}) \\ &= a^{a-b-1}b^{b-a-1}[a+b]^{ab-a-b+1}a^{b-1}b^{a-1} \\ &= a^{a-2}b^{b-2}[a+b]^{ab-a-b+1}. \end{aligned}$$

□

References

[1] J.A. Bondy; U.S.R. Murty, Graph Theory with Applications, THE MACMILLAN PRESS LTD, 1976.

[2] A.K. Kelmans, On properties of the characteristic polynomial of a graph, in: Kibernetiku Na Sluzbu Kom., vol. 4, Gosenergoizdat, Moscow 1967, pp. 27-41 (in Russian).

[3] J. Brown; R. Hoshino, Line graphs and circulants, Ars. Comb. **105** (2012) 463-472.

[4] D. Cvetković; M. Doob; H. Sachs, Spectra of graphs. Theory and application., Academic Press, New York-London, 1980. pp. 39, 218.

[5] F.J. Zhang; Y.-C. Chen; Z.B. Chen, Clique-inserted-graphs and spectral dynamics of clique-inserting, J. Math. Anal. Appl. **349** (2009) 211-225.

[6] W.G. Yan, On the number of spanning trees of some irregular line graphs, J. Combin. Theor. Ser. A **120** (2013) 1642-1648.

[7] F.J. Zhang; W.G. Yan, Enumeration of spanning trees of graphs with an involution, J. Combin. Theor. Ser. A **116** (2009) 650-662.

[8] L. Versfeld, Remarks on star-mesh transformation of electrical networks, Electr. Lett. **6** (1970) 597-599.

[9] E.B. Vahovskii, On the characteristic numbers of incidence matrices for non-singular graphs, Sibirsk. Mat. Zh. **6** (1965) 44-49 (in Russian).

[10] E. Teufl; S. Wagner, Determinant identities for Laplace matrices, Linear Algebra Appl. **432** (2010) 441-457.

[11] M.L. Li; Q.Y. Deng; X. Jin, A general method for computing the Homfly polynomial of DNA double crossover 3-regular links, PLoS ONE **10** (2015): e0125184.

[12] X. Jin; F.J. Zhang; F.M. Dong; E.G. Tay, Zeros of the Jones polynomial are dense in the complex plane, Electr. J. Comb. **17** (2010): #R94.

[13] X. Jin; F.J. Zhang, The replacements of signed graphs and Kauffman brackets of link families, Adv. in Appl. Math. **39** (2007): 155-172.

[14] X. Jin; F.J. Zhang, The Homfly and dichromatic polynomials, Proc. Amer. Math. Soc **140** (2012): 1459-1472.

[15] Y.P. Zhang; X.R. Yong; M. Golin, Chebyshev polynomials and spanning tree formulas for circulant and related graphs, Discrete Math. **298** (2005): 334-364.