arXiv:1605.09314v1 [math.CV] 30 May 2016

WEIGHTED POLYA INEQUALITY IN $\mathbb{C}^n$

OZAN GÜNYÜZ

Abstract. In this article, we obtain a weighted version of Polya inequality in $\mathbb{C}^n$ whose unweighted form was established by V. Zakharyuta in USSR Sbornik, 25 (1975), 350-364.

1. Preliminaries and Introduction

We will denote by $A(\mathbb{C}^n)^*$ the dual space to the space $A(\mathbb{C}^n)$ of all entire functions on $\mathbb{C}^n$, equipped with the locally convex topology of locally uniform convergence in $\mathbb{C}^n$. Following Hörmander([11], Section 4.5), we call the elements of $A(\mathbb{C}^n)^*$ *analytic functionals*.

Let $\mathbb{Z}_+^n$ be the collection of all n-dimensional vectors with non-negative integer coordinates. For $k=(k_1,\ldots,k_\nu,\ldots,k_n)\in\mathbb{Z}_+^n$ and $z=(z_1,\ldots,z_n)\in\mathbb{C}^n$, let $z^k=z_1^{k_1}\ldots z_n^{k_n}$ and $|k|:=k_1+\ldots+k_n$ be the degree of the monomial z^k. We consider the enumeration $\{k(i)\}_{i\in\mathbb{N}}$ of the set $\mathbb{Z}_+^n$ such that $|k(i)|\le|k(i+1)|$ and on each set $\{|k(i)|=s\}$ the enumeration coincides with the lexicographic order relative to $k_1,\ldots,k_n$. We will write $s(i):=|k(i)|$. The number of multiindices of degree at most s is $m_s:=C_{s+n}^s$ and the number of those of degree exactly s is $N_s=m_s-m_{s-1}=C_{n+s-1}^s$, $s\ge1$; $N_0=1$. Let $l_s:=\sum_{q=0}^s qN_q$ for $s=0,1,2,\ldots$.

Consider Vandermondians:

$$V(\zeta_1,\ldots,\zeta_i):=\det\left(e_\alpha(\zeta_\beta)\right)_{\alpha,\beta=1}^i,\ i\in\mathbb{N},$$

where $e_\alpha(z):=z^{k(\alpha)}$, $\alpha\in\mathbb{N}$ and $(\zeta_\beta)\in K^i$.

For a compact set $K\subset\mathbb{C}^n$, define *maximal Vandermondians*:

$$V_i:=\sup\left\{|V(\zeta_1,\ldots,\zeta_i)|:(\zeta_j)\in K^i\right\},\ i\in\mathbb{N}.$$

Set $d_s(K):=(V_{m_s})^{1/l_s}$. The transfinite diameter of K is the number:

$$d(K):=\limsup_{s\to\infty}d_s(K). \tag{1.1}$$

This was introduced by Fekete [7] for $n=1$, and by Leja [14] for $n\ge2$. The proof of usual limit in (1.1) was done in [7] for $n=1$ and in [25] for $n\ge2$.

The *pluripotential Green function* of a compact set $K\subset\mathbb{C}^n$ is defined as follows

$$g_K(z)=\limsup_{\zeta\to z}\sup\{u(\zeta):u|_K\le0,\ u\in\mathcal{L}(\mathbb{C}^n)\},$$

where $\mathcal{L}(\mathbb{C}^n)$ represents the Lelong class consisting of all functions $u\in Psh(\mathbb{C}^n)$, where $Psh(\mathbb{C}^n)$ is the set of all plurisubharmonic functions on $\mathbb{C}^n$, such that $u(\zeta)-\ln|\zeta|$ is bounded from above near infinity. Since $d(K)=d(\hat{K})$, with no loss of generality, we are going to consider polynomially convex compact sets. We say a polynomially convex compact set K is *pluriregular* if $g_K(z)=0$ on K. For more detail about the pluripotential Green function, see [12], [20] and [26]. By an

admissible weight function ω on a compact set $K \subseteq \mathbb{C}^n$, we mean a continuous, non-negative function on K with $\{z \in K : \omega(z) > 0\}$ nonpluripolar. Next we will give the definition of transfinite diameter in weighted version. It was defined by Bloom and Levenberg in [2]. There are two approaches in [2] for the definition of *weighted transfinite diameter*. One of them uses the weighted Vandermonde determinant and the other one is defined by using Chebyshev constant. We are going to use the definition in terms of Vandermonde determinants. Let $K \subseteq \mathbb{C}^n$ be a compact set and ω an admissible weight function K. Given $z_1, z_2, \ldots, z_i \in K$, consider

$$V^{\omega}(z_1, \ldots, z_i) := V(z_1, \ldots, z_i).\omega(z_1)^{|k(i)|} \ldots \omega(z_i)^{|k(i)|}.$$

This is called as *weighted Vandermonde determinant*. Put

$$V_i^{\omega} := \max_{z_1,\ldots,z_i \in K} |V^{\omega}(z_1, \ldots, z_i)|.$$

The weighted transfinite diameter is defined as follows:

$$d^{\omega}(K) := \limsup_{i \to \infty} V_{m_{s(i)}}^{\omega}{}^{\frac{1}{l_{s(i)}}}. \tag{1.2}$$

In the same paper, they proved that there is a limit in (1.2), and they denoted (1.2) by $\delta^{\omega}(K)$. We record the next result from [16]:

Proposition 1.1. Let $K \subseteq \mathbb{C}^n$ be a polynomially convex compact set and ω be an admissible weight on K. There exists a sequence of locally regular compacta $\{K_j\}$ decreasing to K and a sequence of weight functions $\{\omega_j\}$ with ω_j continuous, admissible on K_j, $\omega_{j+1} \leq \omega_j|_{K_{j+1}}$ and $\omega_j \downarrow \omega$. Then

$$\lim_{j \to \infty} d^{\omega_j}(K_j) = d^{\omega}(K).$$

Let K be a compact set in $\mathbb{C}^n$. $A(K)$ represents the locally convex space of all germs of analytic functions on K, equipped with the countable inductive limit topology, i.e.,

$$A(K) = \liminf_{j \to \infty} A(D_j)$$

considered in regard to the inclusion of sets. D_j are open sets such that $D_{j+1} \Subset D_j$ for each $j \in \mathbb{N}$ and $K = \bigcap_{j=1}^{\infty} D_j$. Thus, in this setting, a sequence $\{u_j\}$ of germs converges to a germ u in this topology in case there exists an open neighbourhood $V \supset K$ and functions g_j, $g \in A(V)$ being the representatives of the germs u_j, u respectively, such that g_j converges uniformly to g on any compact subset of V.

2. Polya's Theorem

The following result is due to Polya [19].

Theorem 2.1. *Let K be a polynomially convex compact set in $\mathbb{C}$ and $f \in A\left(\overline{\mathbb{C}} \setminus K\right)$ have the following expansion in a neighbourhood of ∞:*

$$f(z) = \sum_{k=0}^{\infty} \frac{a_k}{z^{k+1}}. \tag{2.1}$$

Let $A_s(f) := \det(a_{k+m})_{k,m=0}^{s-1}$, $s \in \mathbb{N}$, be a sequence of Hankel determinants composed from the coefficients of the expansion (2.1). Then,

$$D(f) := \limsup_{s \to \infty} |A_s(f)|^{1/s^2} \leq d(K). \tag{2.2}$$

A direct multivariate analog of the inequality (2.2) makes no sense, since there are functions analytic on the complement of K but constants only. Schiffer and Siciak ([21]) proved some analog for the product of plane compact sets $K = K_1 \times K_2 \times \ldots \times K_n \subset \mathbb{C}^n$ and functions $f \in A((\overline{\mathbb{C}} - K_1) \times \ldots \times (\overline{\mathbb{C}} - K_n))$. Sheinov ([22], [23]) considered another analog of Polya's inequality for a *linearly convex* compact set K, considering the Taylor expansion at the origin for functions analytic in the domain $D = K^*$ linearly convex adjoint (conjugate) to K (projective complement of K by Martineau [18]).

The case of an arbitrary compact set $K \subset \mathbb{C}^n$ was studied in [25]. It was suggested there, instead of analytic functions on some artificial "complement" of K, to deal with those analytic functionals in $\mathbb{C}^n$ that are extendible continuously onto the space $A(\widehat{K})$. We denote by $A_0(\{\infty^n\})$ the space of all analytic germs f' at $\infty^n = (\infty, \infty, \ldots, \infty) \in \overline{\mathbb{C}^n}$ with Taylor expansion of the form

$$f'(z) = \sum_{k \in \mathbb{Z}^n} \frac{a_k}{z^{k+I}}, \quad I = (1, 1, ..., 1), \tag{2.3}$$

converging in some neighborhood of ∞^n.

Lemma 2.2. *There is an isomorphism,*

$$T : A(\mathbb{C}^n)^* \to A_0(\{\infty^n\}), \tag{2.4}$$

such that, for each f^* and $f' = Tf^*$, we have

$$f^*(\varphi) = \langle \varphi, f' \rangle := \left(\frac{1}{2\pi i}\right)^n \int_{\mathbb{T}_R^n} \varphi(\zeta) f'(\zeta) \, d\zeta, \ \varphi \in A(\mathbb{C}^n),$$

where

$$\mathbb{T}_R^n := \{z = (z_\nu) \in \mathbb{C}^n : |z_\nu| = R, \ \nu = 1, \ldots, n\}, \ R = R(f^*). \tag{2.5}$$

Proof. See, e.g., [6], Chapter 3. □

Let us define, for every analytic functional f^*, a related sequence of multivariate Hankel-like determinants constructed from the coefficients of the expansion (2.3):

$$H_i = H_i(f^*) := \det\left(a_{k(\alpha)+k(\beta)}\right)_{\alpha,\beta=1}^{i}, \ i \in \mathbb{N} \tag{2.6}$$

with $a_{k(\alpha)} := f^*(e_\alpha) = \langle e_\alpha, f' \rangle$, $\alpha \in \mathbb{N}$, $f' = Tf^*$. Now we are ready to formulate the general form of multivariate Polya's inequality.

Theorem 2.3. *Suppose that K is a compact set in $\mathbb{C}^n$, f^* is an analytic functional which has a continuous extension onto $A(K)$ and $f' = Tf^*$ is the corresponding analytic germ at ∞^n. Then for the determinants (2.6), the inequality holds:*

$$D(f') := \limsup_{i \to \infty} |H_i(f^*)|^{\frac{1}{2l_{s(i)}}} \leq d(K). \tag{2.7}$$

It has been proved in [25] a weaker result using the outer transfinite diameter $\widehat{d}(K)$ instead of $d(K)$, however it was proved later that $\widehat{d}(K) = d(K)$ by Znamenskii and Levenberg independently.(see [29], [30] and [15]).

The proof of Theorem 2.3 is essential for the present paper. The basic tool which we will use from that proof is the following equality:

$$i! \, |H_i(f^*)| = |f^*_{\zeta^{(i)}}(\ldots f^*_{\zeta^{(j)}}(\ldots (f^*_{\zeta^{(1)}}([V(\zeta^{(1)}, \zeta^{(2)}, \ldots, \zeta^{(i)})]^2)\ldots)\ldots)|, \tag{2.8}$$

$i \in \mathbb{N}$, here the notation $f^*_{\zeta^{(j)}}$ means that the functional f^* is applied sequentially to a function of the variable $\zeta^{(j)}$ by keeping the other variables fixed.

Let now K be a compact set in $\mathbb{C}^n$ with a decreasing sequence $\{K_j\}$ of locally pluriregular compact sets such that $K_j \downarrow K$. Let $\{\omega_j\}$ be a sequence of admissible weight functions on K_j with $\omega_j \downarrow \omega$, where ω is an admissible weight on K. Considering the coefficients $a_{k(\alpha)}$ defined above, we define the following ones:

$$b_{k(\alpha)} := \omega_j(z)^{2|k(i)|}.a_{k(\alpha)}, \tag{2.9}$$

We form the weighted Hankel-like determinants,

$$H_i^{\omega_j}(f^*) := \det\left(b_{k(\alpha)+k(\beta)}\right)_{\alpha,\beta=1}^{i} \tag{2.10}$$

Now we are ready to give the main theorem:

Theorem 2.4. *Let K, $\{K_j\}$ and $\{\omega_j\}$ be as in Proposition 1.1. Let f^* be an analytic functional which is continuously extendible onto $A(K)$, $f' = T(f^*)$ the analytic germ at ∞^n corresponding to f^*. Then we have, for the determinants (2.10), the following asymptotic inequality*

$$D^{\omega}(f') := \limsup_{i\to\infty} |H_j^{\omega}|^{\frac{1}{i_{s(i)}}} \leq d^{\omega}(K) \tag{2.11}$$

Proof. The proof will be very short because every step follows from the proof of Theorem 2.3. By the setting of what we give in the preliminaries and assumptions, writing the determinant explicitly and examining the proof of Theorem 2.3 especially the relation (2.8), for a given $\epsilon > 0$ we obtain the following estimate

$$\limsup_{i\to\infty} |H_i^{\omega_j}|^{\frac{1}{2i_{s(i)}}} \leq d^{\omega_j}(K_j) + \epsilon. \tag{2.12}$$

Because $\epsilon > 0$ is arbitrary, taking the limit as $j \to \infty$ and using the Proposition 1.1 we have the desired inequality. □

References

[1] T. Bloom, N. Levenberg, *Transfinite Diameter Notions in $\mathbb{C}^N$ and Integrals of Vandermonde Determinants*, Ark.Mat., **48** (2010), 17–40.

[2] T. Bloom, N. Levenberg, *Pluripotential theory and large deviation*, Ind. U. Math. J. 62 no2. (2013) 523-550.

[3] E.Bedford, B.A. Taylor, *A New Capacity for Plurisubharmonic Functions*, Acta Mathematica, **149**, (1982), 1-40.

[4] E.Bedford, B.A. Taylor, *Plurisubharmonic Functions with Logarithmic Singularities*, **38**, 4 (1988), 133-171.

[5] R.Berman, S.Boucksom, *Capacities and Weighted Volumes of Line Bundles*, arXiv:0803.1950

[6] S. Dineen, *Complex Analysis in Locally Convex Spaces*, North-Holland Publishing Company, 1981.

[7] M. Fekete, *Über die Verteilung der Wurzen bei gewissener algebraischen Gleichungen mit ganzzahligen Koefficienten, Math. Z.* **17** (1923), 228-249.

[8] G. M. Goluzin, *On some properties of polynomials,* Mat. Sbornik, **18 (60) (**1946), 227-236.

[9] G.M.Goluzin, *Geometric Theory of Functions of a Complex Variable* , 2nd ed., Nauka, Moscow 1966 (In Russian); English transl., Transl.Math.Monographs, vol.26, Amer. Math. Soc., 1969.

[10] A. Gröthendieck, *Sur certain espaces de fonctions holomorphes*, J. Reine Angew. Math., **192** (1953), 35-64; 77-95.

[11] L.Hörmander, *An Introduction to Complex Analysis in Several Variables*, North Holland and American Elsevier, 1990.

[12] M.Klimek, *Pluripotential Theory*, Oxford Univ. Pres, Oxford, 1991.
[13] G. Köthe, *Dualität in der Funktionentheorie*, J. Reine Angew. Math.**191** (1953), 30-39.
[14] F.Leja, *Problèmes à résoudre posés à la conférence*, Colloqi.Math.7 (1959), 151–153
[15] N. Levenberg, *Capacities in Several Complex Variables*, Thesis, University of Michigan, 1984
[16] N. Levenberg, *Weighted Pluripotential Theory Results of Berman-Boucksom*, arXiv:1010.4035, 2010
[17] N. Levenberg and B. A. Taylor, *Comparison of capacities in* $\mathbb{C}^n$, in: Complex analysis (Toulouse 1983), Springer Lect. Notes Math. 1094 (1984), 162172.
[18] A. Martineau, *Sur la Notion D'ensemble Fortement Lineelement Convexe*, Anais Acad. Brasill. Cienc., 4, No.4, 427-435 (1968).
[19] G. Polya, *Beitrag zur Verallgemeinerung des Verzerrungssatzes auf mehrfach zusammenhängende,Gebiete III*, Sitzber.Preus.Akad. Wiss., Phys.-Math.. Kl., 55–62 (1929).
[20] A. Sadullaev, *Plurisubharmonic Measures and Capacities on Complex Manifolds*, Russian Math.Surveys 36 (1981), no.4, 61–119.
[21] M. Schiffer, J. Siciak, *Transfinite Diameter and Analytic Continuation of Functions of Two Complex Variables*, Studies Math. Analysis and Related topics, Stanford (1962), pp.341–358.
[22] V. P. Sheinov, *Transfinite diameter and Some Theorems of Polya in the case of Several Complex Variables*, Siberian Mathematical Journal, Vol.12, No.6, pp.999-1004, November-December, 1971 (English Translation).
[23] V. P. Sheinov, *Invariant Form of Polya's Inequalities*, Vol.14, No.1, pp.194–203, January-February, 1973 (English Translation).
[24] J. Sebastião e Silva, *Analytic Functions in Functional Analysis*, Portug. Math. **9** (1950), pp.1–130.
[25] V. Zakharyuta, *Transfinite Diameter, Chebyshev Constants and Capacities in* $\mathbb{C}^n$, Math Ussr Sbornik, Vol.25 (1975), 350–364.
[26] V. Zakharyuta, *Extremal Plurisubharmonic Functions, Orthogonal Polynomials and the Bernstein-Walsh Theorem for Analytic Functions of Several Complex Variables*, Ann.Polon.Math. 33 (1976), 137–148(In Russian).
[27] V. Zakharyuta, *Spaces of Analytic Functions and Complex Potential Theory*, Linear Topological Spaces and Complex analysis.1 (1994), 74–146.
[28] V. Zakharyuta, *Transfinite Diameter, Chebyshev Constants and Capacities in* $\mathbb{C}^n$, Ann.Polon.Math. 106 (2012), 293-313.
[29] V. A. Znamenskii, *The Stability of the Transfinite Diameter for a Compact Set in* $\mathbb{C}^n$, Izv. Severo-Kavkaz. Nauchn.Tsentra Vyssh Shkoly Ser. Estestv. Nauk. 1978, no.4, 14–16(Russian)
[30] V. A. Znamenskii, *Chebyshev Constants for a Compact Set in* $\mathbb{C}^n$, Turkish J.Math. 18 (1994), no.3, 229-237

Sabanci University, 34956 Tuzla/Istanbul, Turkey
E-mail address: ozangunyuz@sabanciuniv.edu