National Conference on Gravity and Cosmology, Shahid Beheshti University (2016)

Investigation of energy conditions in the vicinity of future cosmological singularities

Shojai Baghini, Fatimah[1]; Sanati, Mahsa[2]
[1,2]*Department of Physics, University of Tehran, Tehran, Iran*

Abstract

In this paper, considering the energy conditions we shall show that some of the energy conditions are violated in the vicinity of future singularities with the scale factor suggested by Dabrowski - Marosek.

I. Introduction

In cosmology, until the nineties of twentieth century, mainly standard singularities were considered. These singularities started with Big Bang, and according to the curvature of space: a flat or closed universe could expand indefinitely, and an open universe could recollapse to Big Crunch. In 1990's by observing standard candles such as type *Ia* supernovae, the expansion of the universe was discovered. This result was approved by next observations such as the anisotropies in the cosmic microwave background (CMB) radiation and baryon acoustic oscillations (BAO), and cosmologists conclude that Dark Energy is the dominant mass- energy constituent of the universe. As this component must lead to the expansion of the universe, doesn't necessarily satisfy all of the energy conditions; furthermore, many different singularities will appear in cosmological models. Hence, the search and classification of different cosmological singularities began.

II. Cosmological singularities

Type I (Big Rip) [1]: these singularities happen in phantom dark energy models with $w < -1$. In such a universe the very negative equation of state parameter results in a super negative pressure (resulting in super acceleration following the second Friedmann equation) of the dominating dark energy component which together with its ever increasing energy density dissociate everything in the universe.

$$a(t_s) \to \infty\,, \rho(t_s) \to \infty, |p(t_s)| \to \infty$$

Type II (Sudden) [2]: Sudden singularities were first discovered by Barrow as pressure singularities which are accompanied by the divergence of the second derivative of the scale factor.

$$a(t_s) \to a_s\,, \rho(t_s) \to \rho_s, |p(t_s)| \to \infty$$

Type III [3]: these singularities, which are called Big Freeze, happen in phantom models of dark energy. At a Big Freeze singularity, the phantom energy density diverges, filling whole of the universe which means that nothing can move any further.

$$a(t_s) \to a_s\,, \rho(t_s) \to \infty, |p(t_s)| \to \infty$$

Type IV [4]: Such singularities, which were first discovered in the tachyonic cosmological models, were named Big Brake singularities. The first time derivative of the scale factor is zero at this singularity; therefore, at a Big Brake singularity the universe stops its evolution to zero speed.

$$a(t_s) \to a_s\,, \rho(t_s) \to 0, |p(t_s)| \to 0$$

w-singularity [5]: This type of singularity is similar to type IV singularity but with the difference that the higher derivatives of the Hubble parameter do not diverge. w-singularity

would be interpreted as a Big Bang type beginning, fixed and finite size for the scale factor at time of the singularity (t_s), halt of expansion at t_s and end of acceleration at t_s respectively. It can be shown that the equation of state diverges despite the fact that the pressure and the energy density both vanish:

$$w = {}^{p}/_{\rho} = -{}^{2a\ddot{a}}/_{3\dot{a}^2} - {}^{1}/_{3} \quad (1)$$

By making use of the l'Hopital's rule of calculus:

$$\lim_{t\to t_s} -\frac{1}{3}\left(\frac{a}{\dot{a}}+1\right) = -\frac{1}{3}\left(\frac{a_s}{0}+1\right) \to \infty \quad (2)$$

That blows up in the singularity.

III. Energy conditions

Null energy condition (NEC): $\rho + p \geq 0$. According to the Friedmann equations (3, 4), not violating this condition corresponds to:

$$\rho = \frac{3}{8\pi G}\left(\frac{\dot{a}^2}{a^2}+\frac{kc^2}{a^2}\right) \quad (3)$$

$$p = -\frac{c^2}{8\pi G}\left(2\frac{\ddot{a}}{a}+\frac{\dot{a}^2}{a^2}+\frac{kc^2}{a^2}\right) \quad (4)$$

$$\rho + p = 2\frac{\dot{a}^2}{a^2} - 2\frac{\ddot{a}}{a} + \frac{k}{a^2} \geq 0$$

$$\Rightarrow k \geq \ddot{a}a - \dot{a}^2 \quad (5)$$

Weak energy condition (WEC): This condition embodies the NEC and further adds to it the fact that the energy density is positive: $\rho \geq 0$. By taking the velocity of light equal to 1 in the energy density equation:

$$\rho = \frac{3}{8\pi G}\left(\frac{\dot{a}^2}{a^2} - + \frac{k}{a^2}\right) \geq 0 \;\Rightarrow k \geq -\dot{a}^2 \quad (6)$$

Strong energy condition (SEC): $\rho + 3p \geq 0$. Gravity is an attractive force. The violation of this condition comes through negative energy density or a large negative pressure which could occur in inflationary processes such as that in the early universe. Not violating the SEC results in a decelerating universe. Moreover, the SEC provides a lower bound on the energy density.

$$\ddot{a} \leq 0 \quad (7)$$

Dominant energy condition (DEC): $\rho \pm p \geq 0$. Matter cannot travel faster that light. The DEC provides an upper bound on the energy density. This energy condition is interpreted as expressing the same conditions as the WEC plus the fact that pressure should not exceed energy density.

$$\rho - p = \ddot{a}a - \dot{a}^2 \geq 0 \Rightarrow k \geq -\left(\frac{a\ddot{a}}{2} + \dot{a}^2\right) \quad (8)$$

All of the energy conditions are violated for Big Rip, in contrast, for Sudden and Type III just dominant energy condition will be violated in a flat universe model: $k = 0$.

IV. Investigation of energy conditions in the vicinity of singularities

In the Dabrowski– Marosek model, the scale factor is [6]:

$$a(t) = a_s\left(\frac{t}{t_s}\right)^m exp\left(1-\frac{t}{t_s}\right)^n \quad (9)$$

This scale factor is sufficient to encompass all the models we are aware of in the literature, and exponents *m and n* will be used to classify the type of cosmological milestone we are dealing with. In table 1, constraints on the exponents in order to reaching different milestones are shown [7]. Now by using 5-8 conditions and equation (9), energy conditions in the vicinity of singularities will be investigated.




Fig.1. NEC in the vicinity of sudden singularity $(1 < n < 2, a_s = 1, t_s = 10, m = 2/3)$




Fig.2. WEC in the vicinity of sudden singularity $(1 < n < 2, a_s = 1, t_s = 10, m = 2/3)$




Fig.3. SEC in the vicinity of sudden singularity $(1 < n < 2, a_s = 1, t_s = 10, m = 2/3)$

In accordance with Figs. 1-3, in the vicinity of Sudden and Type III singularities:

$lim_{t\to t_s}\ \ddot{a}a - \dot{a}^2 = 0^-$, $lim_{t\to t_s} -\dot{a}^2 = 0^-$,

$$\lim_{t\to t_s} \ddot{a} = 0^-.$$

In contrast, according to Figs.4 and 5:

$$\lim_{t\to t_s} -\left(\frac{a\ddot{a}}{2} + \dot{a}^2\right) = 0^+$$

Therefore, under 5-8 conditions, just in the last equation (DEC), the curvature scalar can't be equal to zero, because according to the equation (8): $k \geq 0^+$. So in the vicinity of these two singularities, in the flat and closed universe, $k = 0,1$, just the dominant energy condition will be violated.




Fig.4. DEC in the vicinity of sudden singularity $(1 < n < 2, a_s = 1, t_s = 10, m = 2/3)$




Fig.5. DEC in the vicinity of Type III singularity $(0 < n < 1, a_s = 1, t_s = 10, m = 2/3)$

In Fig. 6 (NEC), $\ddot{a}a - \dot{a}^2$ as a function of time and the parameter n in the vicinity of Big Rip, and in Fig. 7 as a function of time and discrete values of $n > 2$ in the vicinity of w-singularity is shown. The first of the conditions, (5) must be satisfied in order not to violate the null energy condition. On the other hand, for all values of parameter n:

$$\lim_{t\to t_s} \ddot{a}a - \dot{a}^2 = 0^+$$

Therefore, the curvature scalar can't be equal to zero, and for $k = 0$, which demonstrate the flat universe in FLRW metric, null energy condition and following all of the energy conditions in the vicinity of Big Rip and w-singularity will be violated.




Fig.6. NEC in the vicinity of Big Rip $(n = 1, a_s = 1, t_s = 10, -\infty < m < 0)$




Fig.7. $\ddot{a}a - \dot{a}^2$ as a function of time and discrete values of $n > 2$

	Sudden Singularity	Big Rip	Type III	w-singularity
m	arbitrary	$(-\infty, 0)$	arbitrary	0
n	(1,2)	arbitrary	(0,1)	$(2, \infty)$, $integer$
NEC	Not violated for $k = 0,1$	Violated	Not violated for $k = 0,1$	Violated
WEC	Not violated for $k = 0,1$	Violated	Not violated for $k = 0,1$	Violated
SEC	Not violated for $k = 0,1$	Violated	Not violated for $k = 0,1$	Violated
DEC	Violated	Violated	Violated	Violated

Table1. Energy conditions in the vicinity of Sudden, Big Rip, Type III and w-singularity

[1] R. R. Caldwell, M. Kaminkowski and N. N. Weinberg, Phys. Rev. Lett. 91, 071301 (2003)
[2] J. D. Barrow, Classical Quantum Gravity 21, L79
[3] J. Copeland, M. Sami and Shijini Tsujikawa, hep-th0603057 (2006)
[4] V. Gorini, A.Y. Kamenshchik, U. Moschella, and V. Pasquier, Phys. Rev. D 69, 123512 (2004)
[5] M. P. Dabrowski and T. Denkiewicz, Phys. Rev. D 79, 063521 (2009)
[6] M.P. Dabrowski, K. Marosek, JCAP 02, 012 (2013)
[7] F. Shojai, A. Shojai, M. Sanati, Eur. Phys. J. C,75,568, (2015)