

An Extension of Waring's Problem in $\mathbb{Z}_{p^k}$

Li An-Ping

Beijing 100085, P.R. China
apli0001@sina.com

Abstract

For a set X of integers, and two positive integers $q, a, (q, a) = 1$, denote by

$$X_{q,a} = \{\lambda \mid \lambda q + a \in X\}, \quad X_q = \bigcup_a X_{q,a}, \quad X(q) = \{x \mid x \in X, (x, q) = 1\}.$$

An integer set X in $\mathbb{Z}_{p^k}$ is called strongly uniform, if for arbitrary positive integer

$r \leq k, q = p^r$, it has $|X_q| \ll \lceil |X(q)| / \varphi(q) \rceil$. Denote by Ω the reduced residue

system of $\mathbb{Z}_{p^k}$. For a strongly uniform subset $\mathcal{A} \subseteq \Omega$, $n \in \Omega$, as in Waring's

problem, we will mainly investigate solvability of equation

$$x_1^k + x_2^k + \cdots + x_s^k = n, \quad x_i \in \mathcal{A}, \quad 1 \leq i \leq s.$$

Denote by $\psi_s(n, \mathcal{A})$ the number of solutions of the equation. In this paper, we

will give an upper-bound of $\psi_2(n, \mathcal{A})$.

Keywords: Waring's problem, uniformly distribution, exponential sum, auxiliary equation.

1. Introduction

We know that Waring's problem is to investigate the solvability of the following Diophantine equation

$$x_1^k + x_2^k + \cdots + x_s^k = n, \quad n \in \mathbb{N}, x_i \in (\mathbb{N} \cup 0), 1 \leq i \leq s. \quad (1.1)$$

In 1770, Waring asserted that for each natural number $k \geq 3$, there is a natural number $g(k)$, when $s \geq g(k)$, the equation (1.1) is solvable. The existences of $g(k)$ had been proved by Hilbert, and the precise values have been almost known. The modern version is to find the least number $G(k)$ such that the equation (1.1) is solvable for each sufficient large integer n when $s \geq G(k)$.

Now the main method to estimate $G(k)$ is the so called circle method, which was proposed by Hardy, Ramanujan and Littlewood in 1920's.

In the method, it is encountered to deal with equation (1.1) in the domain $\mathbb{Z}_{p^k}$, that is, the equation

$$x_1^k + x_2^k + \cdots + x_s^k = n, \quad n \in \mathbb{Z}_{p^k}, x_i \in \mathbb{Z}_{p^k}, 1 \leq i \leq s. \quad (1.2)$$

where p is a prime number. For equation (1.2), Hardy and Littlewood's gave a series of researches [1]~[3], which are also one part of basic content of Hardy-Littlewood method. For the detail of Waring's problem, referred to see the Vaughan and Wooley's survey paper [4].

Suppose that X is a set of integers, for two positive integers q, a , $(q, a) = 1$, denote by

$$X_{q,a} = \{\lambda \mid \lambda q + a \in X\}, \quad X_q = \bigcup_{a, (a,q)=1} X_{q,a}, \quad X(q) = \{x \mid x \in X, (x, q) = 1\}, \quad (1.3)$$

An integer set X in $\mathbb{Z}_{p^k}$ is called uniform or strongly uniform distributed respectively if for each

positive integer $r, r \leq k$, $q = p^r$, satisfying condition i) or ii)

$$\text{i) } |X_{q,a}| \ll \lceil |X(q)| / \varphi(q) \rceil, \quad \text{for each integer } a, (q, a) = 1.$$

$$\text{ii) } |X_q| \ll \lceil |X(q)| / \varphi(q) \rceil.$$

Specially, a set of integers in an interval, or a union of finite intervals is strongly uniform, and a segment of arithmetic progression $\{ax + b \mid x \geq 0, (a, p) = 1\}$ is too, where $0 < a \leq c_0$ a constant.

It is assumed that $k > 1$ in this paper, and simply write $q = p^k$. Denote by Ω the reduced residue system of $\mathbb{Z}_q$.

Let $\mathcal{F}_r$ be the set of all strongly uniform r -subsets of Ω , for a set $\mathcal{A} \in \mathcal{F}_m$, and $n \in \Omega$, we investigate the solvability of equation

$$x_1^k + x_2^k + \cdots + x_s^k = n, \quad x_i \in \mathcal{A}, \quad 1 \leq i \leq s. \quad (1.4)$$

Denote by $\psi_s(n, \mathcal{A})$ the number of solutions of equation (1.4). define

$$\Psi_s(m, q) = \max_{\mathcal{A} \in \mathcal{F}_m} \max_{n \in \Omega} \{\psi_s(n, \mathcal{A})\}. \quad (1.5)$$

It is clear that the expected results are that

$$\Psi_s(m, q) \ll \lceil m^s / \varphi(q) \rceil.$$

In this paper, we will prove it for $s = 2$. That is,

Theorem 1. If $(C_k^2, p) = 1$, then

$$\Psi_2(m, q) \ll \lceil m^2 / \varphi(q) \rceil. \quad (1.6)$$

The result (1.6) can be promoted to more general polynomials, suppose that $f(x) =$

$$\sum_{0 \leq i \leq n} a_i x^{n-i}, \quad n > 1, \quad a_i \in \mathbb{Z}_q, \quad 0 \leq i \leq n, \quad (a_0, p) = 1, \text{ denote by}$$

$$\Omega'(f) = \{x \mid x \in \mathbb{Z}_q, f'(x) \equiv 0 \pmod{p}\}, \quad \Omega(f) = \Omega \setminus \Omega'(f).$$

Similarly, let $\mathcal{F}_r(f)$ be the set of all strongly uniform r -subsets of $\Omega(f)$, for a set $\mathcal{A} \in \mathcal{F}_m(f)$,

and $n \in \Omega$, we investigate the solvability of equation

$$f(x_1) + f(x_2) + \cdots + f(x_s) = n, \quad x_i \in \mathcal{A}, \quad 1 \leq i \leq s. \quad (1.7)$$

Denote by $\psi_s(n, \mathcal{A}; f)$ the number of solutions of equation (1.7), define

$$\Psi_s(m, q; f) = \max_{\mathcal{A} \in \mathcal{F}_m(f)} \max_{n \in \Omega} \{\psi_s(n, \mathcal{A}; f)\}. \quad (1.8)$$

There is

Theorem 2.

$$\Psi_2(m, q; f) \ll \lceil m^2 / \varphi(q) \rceil. \quad (1.9)$$

In the following proof, it will be applied four classic results of exponential sums in elementary number theory. As usual, simply write $e(x) = e^{2\pi i x}$.

Lemma 1.1.

$$\sum_{x \in \Omega} e(ax / p^k) = \begin{cases} \varphi(p^k), & \text{if } a \equiv 0 \pmod{p^k}, \\ -p^{k-1}, & \text{if } p^{k-1} \parallel a, \\ 0, & \text{else.} \end{cases} \quad (1.10)$$

Lemma 1.2. Suppose that m is a odd number, and $(m, a) = 1$, then

$$\sum_{x=0}^{m-1} e(ax^2 / m) = \begin{cases} \left(\frac{a}{m}\right) \sqrt{m}, & \text{if } m \equiv 1 \pmod{4}, \\ i \left(\frac{a}{m}\right) \sqrt{m}, & \text{if } m \equiv 3 \pmod{4}. \end{cases} \quad (1.11)$$

Where $\left(\frac{a}{m}\right)$ is the Jacobi symbol, $\left(\frac{a}{m}\right) = 1$ or -1 , if a is a quadratic residue, or not.

Lemma 1.3. Suppose that $\chi(n)$ is an improper character, $(a, p) = 1$, then

$$\sum_{1 \leq n \leq p^l} e(an / p^l) \chi(n) = 0. \quad (1.12)$$

Lemma 1.4. Suppose that $p > 2$, $a^2 - 4b \not\equiv 0 \pmod{p}$, then

$$\sum_{0 \leq x < p} \left(\frac{x^2 + ax + b}{p} \right) = -1. \quad (1.13)$$

2. An Auxiliary equation.

For $N \in \Omega$, $a, b \in \mathbb{Z}_q \setminus \{0\}$, $z \in \mathbb{Z}_q$, consider following equation

$$\begin{cases} x^k + y^k \equiv N \pmod{q}, & x, y \in \Omega, \\ ax + by \equiv z \pmod{q}. \end{cases} \quad (2.1)$$

Denote by $\mathfrak{S}(z, a, b)$ the set of solutions (x, y) of (2.1), and with respect to $\mathfrak{S}(z, a, b)$, we define

$$\rho_i = \rho_i(x, y; a, b) = x^{k-i} b^i + (-a)^i y^{k-i}, \quad \tau_i = \tau_i(x, y; a, b) \Leftrightarrow p^{\tau_i} \parallel \rho_i, \quad i = 1, 2, \dots.$$

Lemma 2.1. Suppose that $(x, y) \in \mathfrak{S}(z, a, b)$, $p^{r-1} \parallel \rho_1$, if $2r-1 < k$, then for arbitrary

integer t , there is $(u, v) \in \mathfrak{S}(z + tp^{2r-1}, a, b)$, with that $u = x + cp^r$, $v = y + dp^r$.

Proof. Clearly, it suffices to show that for arbitrary integer t , there are integers c, d , such that

$$u = x + cp^r, v = y + dp^r, (u, v) \in \mathfrak{S}(z + tp^{2r-1}, a, b). \quad (2.2)$$

Then

$$au + bv \equiv ax + by + (ac + bd)p^r \equiv w + tp^{2r-1} \pmod{q}.$$

$\Rightarrow$

$$(ac + bd)p^r \equiv tp^{2r-1} \pmod{q}$$

And

$$ac + bd \equiv tp^{r-1} \pmod{q'}, \quad q' = q / p^r.$$

Besides,

$$\begin{aligned} u^k + v^k &= (x + cp^r)^k + (y + dp^r)^k \\ &= (x^k + y^k) + C_k^1 p^r (x^{k-1}c + y^{k-1}d) + C_k^2 p^{2r} (x^{k-2}c^2 + y^{k-2}d^2) + \dots \end{aligned} \quad (2.3)$$

$\Rightarrow$

$$C_k^1 p^r (x^{k-1}c + y^{k-1}d) + C_k^2 p^{2r} (x^{k-2}c^2 + y^{k-2}d^2) + \dots = 0$$

And

$$(x^{k-1}c + y^{k-1}d) \equiv 0 \pmod{p^r}.$$

So,

$$x^{k-1}bc + y^{k-1}bd \equiv x^{k-1}bc + y^{k-1}(tp^{r-1} - ac) \equiv (x^{k-1}b - ay^{k-1})c + tp^{r-1}y^{k-1} \equiv 0 \pmod{p^r}$$

Suppose $(x^{k-1}b - ay^{k-1}) = l \cdot p^{r-1}$, $l \not\equiv 0 \pmod{p}$. It follows

$$lc + ty^{k-1} \equiv 0 \pmod{p}. \quad (2.4)$$

Suppose $c = \sum_{i \geq 0} c_i p^{ir}$, $d = \sum_{i \geq 0} d_i p^{ir}$, from (2.4), c_0 and d_0 are followed. And then consider

the rest terms of (2.3) similarly, c_i, d_i will be followed in turn $i \geq 0$. $\square$

For $a, b \in \mathbb{Z}_q$, here notation $(a, b) = 1$ means $a \not\equiv b \pmod{p}$. Let $\epsilon = 1$, or 2 as k is odd or even.

Lemma 2.2. Suppose that $(a, b) = 1$ and $(x, y) \in \mathfrak{S}(z, a, b)$, $p^r \parallel \rho_1$, $p \nmid \rho_2$, $r \geq \lceil (k+1)/2 \rceil$,

let $\lambda = \lfloor (k-1)/2 \rfloor$, then there are $(u, v) \in \mathfrak{S}(z + tp^{2\lambda}, a, b)$, with that $u = x + cp^\lambda$,

$v = y + dp^\lambda$, and as c is over $\mathbb{Z}_{p^\epsilon}$, t will take over all quadratic (non-)residue classes, depend

on ρ_2 is or not a quadratic residue.

Proof. Assume that $(u, v) \in \mathfrak{S}(z + tp^{2\lambda}, a, b)$, then

$$(ac + bd)p^\lambda - tp^{2\lambda} \equiv 0 \pmod{q},$$

i.e.

$$ac + bd \equiv tp^\lambda \pmod{q'}, \quad q' = p^{k-\lambda}. \quad (2.5)$$

And

$$u^k + v^k = x^k + y^k + C_k^1(x^{k-1}c + y^{k-1}d)p^\lambda + C_k^2(x^{k-2}c^2 + y^{k-2}d^2)p^{2\lambda} + \dots$$

$\Rightarrow$

$$C_k^1(x^{k-1}c + y^{k-1}d)p^\lambda + C_k^2(x^{k-2}c^2 + y^{k-2}d^2)p^{2\lambda} + \dots \equiv 0 \pmod{q}$$

$$\begin{aligned} b(x^{k-1}c + y^{k-1}d)p^\lambda &= (x^{k-1}bc + y^{k-1}bd)p^\lambda \\ &\equiv (x^{k-1}bc + y^{k-1}(tp^\lambda - ac))p^\lambda \equiv ((x^{k-1}b - ay^{k-1})c + tp^\lambda y^{k-1})p^\lambda \pmod{q} \\ &\equiv ty^{k-1}p^{2\lambda} \pmod{q}. \end{aligned}$$

Hence,

$$C_k^2(x^{k-2}c^2 + y^{k-2}d^2)p^{2\lambda} + C_k^1ty^{k-1}p^{2\lambda}/b \equiv 0 \pmod{q}. \quad (2.6)$$

From (2.5), we have

$$(ac)^2 - (bd)^2 \equiv \mu p^\lambda \pmod{q'} \quad (2.7)$$

With (2.6), (2.7), it has

$$b^2 \cdot (x^{k-2}c^2 + y^{k-2}d^2)p^{2\lambda} = (x^{k-2}b^2c^2 + y^{k-2}((ac)^2 - \mu p^\lambda) + \omega)p^{2\lambda} \equiv 0 \pmod{q},$$

where $\omega \equiv ty^{k-1}2b/(k-1)$. i.e.

$$\begin{aligned} (x^{k-2}b^2 + y^{k-2}a^2)c^2 + \omega &\equiv 0 \pmod{p^\epsilon} \\ t \equiv \varrho c^2 \pmod{p}, \quad \varrho &= -(k-1)\rho_2/(2by^{k-1}). \end{aligned}$$

For $\varrho \not\equiv 0 \pmod{p}$, so t will take over all quadratic (non-)residue classes as c is from 1 to p^ϵ ,

rely on ϱ is a quadratic residue or not. □

In fact, there is a little more general result. Define $\Delta = k + \tau_2 - 2\tau_1$, and

$$\lambda = \begin{cases} k - \tau_1 - 1, & \text{if } \Delta \geq 0, \\ \lceil (k - 1 - \tau_2) / 2 \rceil, & \text{if } \Delta < 0. \end{cases}$$

$$\delta = \begin{cases} 2\tau_1 + 1 - k, & \text{if } \Delta > 0, \\ 2\tau_1 - k, & \text{if } \Delta = 0, \\ \tau_2, & \text{if } \Delta < 0. \end{cases}$$

Lemma 2.3. Suppose that $(a, b) = 1$, and $(x, y) \in \mathfrak{S}(z, a, b)$, and $\tau_1 \geq \lceil (k+1)/2 \rceil$.

If $\Delta \geq 2$, then for arbitrary integer t , there is $(u, v) \in \mathfrak{S}(z + tp^{k-1}, a, b)$, with that $u = x + cp^\lambda$, $v = y + dp^\lambda$.

If $\Delta \leq 1$, then there is $(u, v) \in \mathfrak{S}(z + tp^{2\lambda}, a, b)$, with that $u = x + cp^\lambda$, $v = y + dp^\lambda$, and as c is over Z_p or Z_{p^2} , t will take over all quadratic (non-)residue classes, rely on ρ_2 is or not a quadratic residue.

Proof. Let $\delta = 2r + 1 - k$, then $k - 1 = 2\lambda + \delta$. Assume $(u, v) \in \mathfrak{S}(z + tp^{2\lambda+\delta}, a, b)$, then

$$(ac + bd)p^\lambda - tp^{2\lambda+\delta} \equiv 0 \pmod{q},$$

i.e.

$$ac + bd \equiv tp^{\lambda+\delta} \pmod{q'}, \quad q' = p^{k-\lambda}. \quad (2.8)$$

And

$$u^k + v^k = x^k + y^k + C_k^1(x^{k-1}c + y^{k-1}d)p^\lambda + C_k^2(x^{k-2}c^2 + y^{k-2}d^2)p^{2\lambda} + \dots$$

$\Rightarrow$

$$C_k^1(x^{k-1}c + y^{k-1}d)p^\lambda + C_k^2(x^{k-2}c^2 + y^{k-2}d^2)p^{2\lambda} + \dots \equiv 0 \pmod{q}$$

$$b(x^{k-1}c + y^{k-1}d)p^\lambda = (x^{k-1}bc + y^{k-1}bd)p^\lambda \equiv (x^{k-1}bc + y^{k-1}(tp^{\lambda+h} - ac))p^\lambda$$

$$\equiv ((x^{k-1}b - ay^{k-1})c + tp^{\lambda+\delta}y^{k-1})p^\lambda$$

$$\equiv \beta cp^{\lambda+r} + ty^{k-1}p^{2\lambda+\delta} \pmod{q}.$$

Hence,

$$C_k^2(x^{k-2}c^2 + y^{k-2}d^2)p^{2\lambda} + C_k^1(\beta cp^{\lambda+r} + ty^{k-1}p^{2\lambda+\delta})/b \equiv 0 \pmod{q} \quad (2.9)$$

From (2.8), we have

$$(ac)^2 - (bd)^2 \equiv \mu tp^{\lambda+\delta} \pmod{q'} \quad (2.10)$$

Write $\omega \equiv (\beta cp^{r-\lambda} + ty^{k-1}p^\delta)2b/(k-1)$, with (2.9), (2.10), it has

$$(b^2 \cdot (x^{k-2}c^2 + y^{k-2}d^2) + \omega) p^{2\lambda} = (x^{k-2}b^2c^2 + y^{k-2}((ac)^2 - \mu tp^{\lambda+r}) + \omega) p^{2\lambda} \pmod{q},$$

i.e.

$$\rho_2 c^2 + \omega \equiv 0 \pmod{p^{k-2\lambda}} \quad (2.11)$$

If $\Delta \geq 2$, then $\tau_2 + 2\lambda \geq k$, $r - \lambda = \delta = k - 1 - 2\lambda$, so, (2.11) is turn into that

$$\beta c + ty^{k-1} \equiv 0 \pmod{p}.$$

For the rest cases, let $\nu = \rho_2(k-1)/(2bp^{\tau_2})$, then (2.11) will be as respectively that

$$\nu c^2 + \beta c + ty^{k-1} \equiv 0 \pmod{p}, \text{ if } \Delta = 1, \text{ or}$$

$$\nu c^2 + \beta pc + ty^{k-1} \equiv 0 \pmod{p^2}, \text{ if } \Delta = 0, \text{ or}$$

$$\nu c^2 + \beta p^{\epsilon_0} c + ty^{k-1} \equiv 0 \pmod{p^{\epsilon}}, \epsilon_0, \epsilon = 1, \text{ or } 2, \text{ if } \Delta < 0.$$

It should be mentioned that there are possible cases that $\delta \geq \lambda$, then we can yet as Lemma 1 by the rest congruencies to calculate the rest part of c and d . $\square$

Let $\varepsilon = 1$, or 0 rely upon k is odd or even. For $(a, b) = 1$, define

$$\mathfrak{S}'(z, a, b) = \{ (x, y) \mid (x, y) \in \mathfrak{S}(z, a, b), \tau_1(x, y, a, b) \geq (k + \varepsilon)/2 \}.$$

Lemma 2.4. Suppose that $(a, b) = 1$, and $(x, y) \in \mathfrak{S}'(z, a, b)$, then there are just $p^{(k-\varepsilon)/2}$ pairs

$$(x_i, y_i) \in \mathfrak{S}'(z, a, b), 1 \leq i \leq p^{(k-\varepsilon)/2}, \text{ with that } x_i \equiv x \pmod{p^{(k+\varepsilon)/2}}, y_i \equiv y \pmod{p^{(k+\varepsilon)/2}}.$$

Proof. For $\tau_1 \geq (k + \varepsilon)/2$, this indicates that polynomial $f(x) = x^k + (z - ax)^k / b^k - N$ has

a duplicative root $x_0 \pmod{p^{\tau_1}}$, then it is easy to know that $x_0 + \lambda p^{(k+\varepsilon)/2}$, $0 \leq \lambda < p^{(k-\varepsilon)/2}$ will

be the roots of polynomial $f(x)$. $\square$

It is not difficult to know that, for arbitrary $\lambda \in \mathbb{Z}_q$, let $b_\lambda = b + \lambda p^{(k+\varepsilon)/2}$, and for arbitrary

$(x, y) \in \mathfrak{S}'(z, a, b)$, it has

$$\mathfrak{S}'(z, a, b) = \mathfrak{S}'(z_\lambda, a, b_\lambda). \quad (2.12)$$

Where $z_\lambda = z + y\lambda p^{(k+\varepsilon)/2}$.

Suppose that u, v are two constants, let $w_t = au + b_tv$, $\zeta_t = z_t - w_t$, assume that $\zeta_0 \equiv 0 \pmod{p^{(k-1)}}$. Then

$$\zeta_t \equiv (z_t - z_0) - (w_t - w_0) + (w_0 - z_0) \equiv (y - v) \cdot \lambda p^{(k+\varepsilon)/2} \pmod{p^{k-1}}.$$

So, if $y \not\equiv v \pmod{p^{(k-\varepsilon)/2}}$, then ζ_t will run a period of length $p^{(k-\varepsilon)/2}$ as t take from 0 to $p^{(k-\varepsilon)/2}$.

Moreover, let ϵ as defined ahead Lemma 2.2, we have

Lemma 2.5. Suppose that $(a, b) = 1, (x, y) \in \mathfrak{S}(z, a, b)$, $\rho_1(x, y; a, b) = 0$. And $\lambda \in \mathbb{Z}_q$, let

$b' = b + \lambda p^{(k-\epsilon)/2}$, then there is $(x', y') \in \mathfrak{S}(z', a, b')$, with that $x' \equiv x \pmod{p^{(k-\epsilon)/2}}$,

$y' \equiv y \pmod{p^{(k-\epsilon)/2}}$, $z' \equiv z + a\lambda y p^{(k-\epsilon)/2} \pmod{p^{(k-\epsilon)}}$.

Proof. As $(a, b) = a(1, ba^{-1})$, so it suffices to prove the Lemma for the case $a = 1$.

Let $\lambda = bt$, $\omega = tp^{(k-\epsilon)/2}$, $\theta = b^{k/(k-1)}$, $\theta' = (b')^{k/(k-1)}$, then it is easy to know that

$$x = (1/(1+\theta))^{1/k} N^{1/k}, y = (\theta/(1+\theta))^{1/k} N^{1/k}, \text{ and}$$

$$x' = (1/(1+\theta'))^{1/k} N^{1/k}, y' = (\theta'/(1+\theta'))^{1/k} N^{1/k}.$$

So,

$$\begin{aligned} x' &= \left(\frac{1}{1+\theta(1+\omega)^{k/(k-1)}} \right)^{1/k} N^{1/k} \\ &= \frac{1}{(1+\theta)^{1/k}} \left(1 - \frac{\theta}{(1+\theta)(k-1)} \omega + \frac{k\theta^2 - \theta}{2!(1+\theta)^2(k-1)^2} \omega^2 \right) N^{1/k} \\ &= x \left(1 - \frac{\theta}{(1+\theta)(k-1)} \omega + \frac{k\theta^2 - \theta}{2!(1+\theta)^2(k-1)^2} \omega^2 \right) \\ y' &= \left(\frac{\theta(1+\omega)^{k/(k-1)}}{1+\theta(1+\omega)^{k/(k-1)}} \right)^{1/k} N^{1/k} \\ &= \left(\frac{\theta}{(1+\theta)} \right)^{1/k} \left(1 + \frac{1}{(k-1)(1+\theta)} \omega - \frac{(2k-1)\theta + (k-2)}{2!(1+\theta)^2(k-1)^2} \omega^2 \right) N^{1/k} \\ &= y \left(1 + \frac{1}{(k-1)(1+\theta)} \omega - \frac{(2k-1)\theta + (k-2)}{2!(1+\theta)^2(k-1)^2} \omega^2 \right) \end{aligned}$$

So,

$$\begin{aligned}
& (x' - x) + b'(y' - y) \\
&= x \left(-\frac{\theta}{(1+\theta)(k-1)} \omega + \frac{k\theta^2 - \theta}{2!(1+\theta)^2(k-1)^2} \omega^2 \right) + by \left(\frac{1}{(k-1)(1+\theta)} \omega + \frac{k-\theta}{2!(k-1)^2(1+\theta)^2} \omega^2 \right) \\
&= \frac{by - \theta x}{(k-1)(1+\theta)} \omega + \frac{(k\theta^2 - \theta)x + (k-\theta)by}{2!(1+\theta)^2(k-1)^2} \omega^2 \\
&= \frac{by}{2!(1+\theta)(k-1)} \omega^2
\end{aligned}$$

And $z' - z = (x' - x) + b'(y' - y) + \lambda y p^{(k-\epsilon)/2}$. □

Hence,

$$\begin{aligned}
\zeta_t &= a(x_t - u) + b_t(y_t - v) = a(x - u) + a(x_t - x) + b_t(y - v) + b_t(y_t - y) \\
&= a(x - u) + b(y - v) + b\omega(y - v) + a(x_t - x) + b_t(y_t - y) \\
&= \zeta_0 + btp^{(k-\epsilon)/2}(y - v) + a(x_t - x) + b_t(y_t - y)
\end{aligned}$$

Assume that $\zeta_0 = \alpha p^{k-1}$, $y - v = \mu p^{(k-\epsilon)/2}$, by Lemma above, it follows

$$\begin{aligned}
\zeta_t &= \alpha p^{(k-1)} + bt\mu p^{(k-\epsilon)} + \frac{by}{2!(1+\theta)(k-1)} t^2 p^{(k-\epsilon)} \\
&= \left(\alpha p^{\epsilon-1} - \frac{b(1+\theta)(k-1)}{2y} \mu^2 \right) p^{(k-\epsilon)} + \left(\frac{(1+\theta)(k-1)}{y} \mu + t \right)^2 \frac{by}{2(1+\theta)(k-1)} p^{(k-\epsilon)} \\
&= \left(\alpha p^{\epsilon-1} - \frac{b}{2} \xi \mu^2 \right) p^{(k-\epsilon)} + \frac{b}{2\xi} (\xi \mu + t)^2 p^{(k-\epsilon)}, \quad \xi = \frac{(1+\theta)(k-1)}{y}.
\end{aligned}$$

Hence,

$$\begin{aligned}
\sum_{0 \leq t < p^\epsilon} e(a\zeta_t / q) &= e\left(a(\alpha p^{\epsilon-1} - b\xi\mu^2 / 2) / p^\epsilon\right) \sum_{0 \leq t < p^\epsilon} e\left(a \frac{b}{2\xi} (\xi\mu + t)^2 / p^\epsilon\right) \\
&= e\left(a\sigma / p^\epsilon\right) p^{\epsilon/2} \chi(ab / 2\xi), \quad \sigma = \alpha p^{\epsilon-1} - b\xi\mu^2 / 2.
\end{aligned}$$

Where $\sigma = \alpha p^{\epsilon-1} - b\xi\mu^2 / 2$, and $\chi(t) = \left(\frac{t}{p}\right)$, or $i\left(\frac{t}{p}\right)$ depending on $p \equiv 1, \text{ or } 3 \pmod{4}$.

Let ϱ be as in the proof of Lemma 2.2, it is easy to follows that

$$\chi(\varrho)\chi(b / 2\xi) = \chi(\varrho b / 2\xi) = 1. \text{ So,}$$

$$\sum_a \chi(a\varrho) \sum_{0 \leq t < p^\epsilon} e(a\zeta_t / q) = \sum_a e(\sigma a / p^\epsilon) p^{\epsilon/2} = \gamma p^{\epsilon/2}. \quad (2.13)$$

By Lemma 1.1, we know that $\gamma = \varphi(p^k)$, or $-p^{k-1}$ depend on $\sigma = 0$, or $\sigma \neq 0$.

We call the procedure described in the proof of Lemma 2.5 that from (x, y) to $(\tilde{x}, \tilde{y})$ by

changing b to $\tilde{b}(=b(1+\omega))$ as a transformation, (x, y) and $(\tilde{x}, \tilde{y})$ are called to be homotopic. From the proof above, it is easy to know that

Lemma 2.6. Suppose that $u, v \in \mathcal{A}$, $(x, y) \in \mathfrak{S}'(z, 1, b)$, and $y \equiv v \pmod{p^r}$, $x \equiv u \pmod{p^r}$, $r \geq (k - \varepsilon)/2$, $(x - u) + b(y - v) \equiv 0 \pmod{p^{k-1}}$, then (x, y) is homotopic to a $(\tilde{x}, \tilde{y})$ with that $\tilde{y} \equiv v \pmod{p^{k-1}}$, $\tilde{x} \equiv u \pmod{p^{k-1}}$, $(\tilde{x} - u) + \tilde{b}(\tilde{y} - v) \equiv \alpha p^{k-1} \pmod{q}$.

With the Lemma 2.6, we may classify the $\{b, (x, y), (u, v)\}$ satisfying the conditions as in the lemma into p homotopic classes $\mathcal{H}_\alpha$, $\alpha = 0, 1, \dots, p-1$.

For the class $\alpha = 0$, then $\zeta_t = bt^2 p^{(k-\varepsilon)}/2\xi$, and for each t , $0 \leq t < p^{(k+\varepsilon)/2}$, it has

$$\begin{aligned} \sum_{a \in \Omega} e(a\zeta_t/q) \chi(a\varrho) &= \sum_{a \in \Omega} e\left(a \frac{b}{2\xi} t^2 / p^\varepsilon\right) \chi(a\varrho) = \sum_{a \in \Omega} e(a/p^\varepsilon) \chi(abt^2 \varrho / 2\xi) \\ &= \sum_{a \in \Omega} e(a/p^\varepsilon) \chi(a) \chi(bt^2 \varrho / 2\xi) \\ &= \begin{cases} 0, & \text{if } t \equiv 0 \pmod{p}, \\ \sum_{a \in \Omega} e(a/p^\varepsilon) \chi(a) = p^{k-\varepsilon/2}, & \text{if } t \not\equiv 0 \pmod{p}. \end{cases} \end{aligned} \quad (2.14)$$

For the class $\alpha \neq 0$, by Lemma 1.4, we can know

$$\begin{aligned} \sum_{0 \leq t < p^\varepsilon} \sum_{a \in \Omega} e(a\zeta_t/q) \chi(a\varrho) &= \sum_{a \in \Omega} e(a \cdot (\alpha' + t^2) / p^\varepsilon) \chi(ab\varrho / 2\xi) \\ &= \sum_{0 \leq t < p^\varepsilon} \chi(\alpha' + t^2) \sum_{a \in \Omega} e(a/p^\varepsilon) \chi(a) \\ &= - \sum_{a \in \Omega} e(a/p^\varepsilon) \chi(a) \\ &= -p^{k-\varepsilon/2}, \quad (\alpha' = \alpha 2\xi / b). \end{aligned} \quad (2.15)$$

Besides, for $b \in \Omega$, denote by $[b]_r$ the least positive integer with that $b \equiv [b]_r \pmod{p^r}$.

Define $\Theta(r)$ as the number of triples $([b]_r, u, v)$ such that there is $(x, y) \in \mathfrak{S}'(z, 1, b)$, $x \equiv u \pmod{p^r}$, $y \equiv v \pmod{p^r}$. And for a integer δ , $0 \leq \delta < p^d \leq |\mathcal{A}|$, denote by $\Theta(r)|_\delta$ the number of ones in the $\Theta(r)$ triples with that $[b]_d = \delta$.

Lemma 2.7.

$$\begin{aligned}\Theta(r) &\ll \left\lceil |\mathcal{A}|^2 / \varphi(p^r) \right\rceil, \\ \Theta(r)|_\delta &\ll \left\lceil |\mathcal{A}|^2 / \varphi(p^{r+d}) \right\rceil.\end{aligned}\tag{2.16}$$

Proof. Let $r_0 = \lceil \log_p |\mathcal{A}| - 1 \rceil$, if $r \leq r_0$, by the uniformity of set $\mathcal{A}$, then

$$\Theta(r) \ll \varphi(p^r) \cdot \left(|\mathcal{A}| / \varphi(p^r) \right)^2 = |\mathcal{A}|^2 / \varphi(p^r), \text{ and}$$

$$\Theta(r)|_\delta \ll \left\lceil |\mathcal{A}|^2 / \varphi(p^{r+d}) \right\rceil.$$

So, assume that $r > r_0$. Suppose that $|\mathcal{A}| \doteq s \cdot p^{r_0}$, $\bar{q} = p^{r_0}$, and by the definition, without loss the generality, let

$$\mathcal{A}_{\bar{q}} = \{\lambda_i\}_1^s, \lambda_i = \sum_{t \geq 0} c_{i,t} p^t, \quad 0 \leq c_{i,t} < p, \quad 1 \leq i \leq s.$$

For a integer $l, 1 \leq l \leq r$, let $\Omega_l = \{x \mid 0 < x < p^l, (x, p) = 1\}$, considering the equation

$$\begin{cases} x^k + y^k \equiv N \pmod{p^l}, & x, y \in \Omega_l \\ bx^{k-1} - y^{k-1} \equiv 0 \pmod{p^l}. \end{cases}\tag{2.17}$$

Denote by $\mathfrak{M}_l$ the set of solutions of equation (2.17). For a pair $(x, y) \in \mathfrak{M}_l$, then it is easy to know that there is the following expression

$$[x]_l = \frac{([N]_l)^{1/k}}{(1 + \mathcal{G}_l)^{1/k}}, \quad [y]_l = \frac{\mathcal{G}_l^{1/k} ([N]_l)^{1/k}}{(1 + \mathcal{G}_l)^{1/k}}.\tag{2.18}$$

where $\mathcal{G}_l = ([b]_l)^{k/(k-1)}$.

Moreover, for a integer $h, 0 \leq h < p^{r-l}$, define

$$\mathfrak{M}_l(h, r) = \left\{ (x, y) \mid (x, y) \in \mathfrak{M}_l, x^k + y^k \equiv [N]_l + h \cdot p^l \pmod{p^r} \right\}.$$

It is clear that

$$\mathfrak{M}_l = \bigcup_{0 \leq h < p^{r-l}} \mathfrak{M}_l(h, r).\tag{2.19}$$

Clearly,

$$[x]_l = [x]_{r_0} + \sum_{r_0 \leq i < l} x_i p^i, \quad [y]_l = [y]_{r_0} + \sum_{r_0 \leq i < l} y_i p^i, \quad \text{for } l \geq r_0.\tag{2.20}$$

Furthermore, for a pair $(i, j), 1 \leq i, j \leq s$, in the expression (2.20), let $x_{r_0+t} = c_{i,t}, y_{r_0+t} = c_{j,t}$,

$0 \leq t < r - r_0$. i.e.

$$[x]_l = [x]_{r_0} + \sum_{0 \leq t < l - r_0} c_{i,t} p^{r_0+t}, \quad [y]_l = [y]_{r_0} + \sum_{0 \leq t < l - r_0} c_{j,t} p^{r_0+t}. \quad (2.21)$$

With (2.18), it has

$$[x]_l = \frac{[N]_{r_0}^{1/k}}{(1 + \mathcal{G}_{r_0})^{1/k}} + \sum_{0 \leq t < l - r_0} c_{i,t} p^{r_0+t}, \quad [y]_l = \frac{\mathcal{G}_{r_0}^{1/k} [N]_{r_0}^{1/k}}{(1 + \mathcal{G}_{r_0})^{1/k}} + \sum_{0 \leq t < l - r_0} c_{j,t} p^{r_0+t}. \quad (2.22)$$

Substituting (2.22) into the first equation of (2.17), and taking $l = r_0 + 1, r_0 + 2, \dots, r$ respectively, then in each set $\mathfrak{M}_{r_0}(h, r), b_i, i = 0, 1, \dots, (r - r_0 - 1)$, will be determined in turn.

Moreover, let $d = r - r_0$, by the induction on the second expression of (2.16), roughly speaking, the sizes of sets in $\left\{ \mathfrak{M}_{r_0}(h, r) \Big|_{\delta} \mid 0 \leq \delta < p^d \right\}$ are about equal. Therefore, there are at most p^{2r_0-r} solutions for the variables b_i 's.

For there are s^2 selections of pair (λ_i, λ_j) , $1 \leq i, j \leq s$. Hence

$$\Theta(r) \ll s^2 \cdot p^{2r_0-r} \ll \left\lceil |\mathcal{A}|^2 / p^r \right\rceil.$$

□

It may be said that Lemma (2.17) is a crux of the issue, which is once used inadvertently in the previous version yet, however, it is not true actually in general for the uniform sets which merely meet the condition i), that can be perceived from the proof, where is likely the main flaw such that the previous version failed.

3. The Proof of Theorem 1.

Suppose that $\mathcal{A} \in \mathcal{T}_m$, let $g(x)$ be the characteristic function of $\mathcal{A}$ in Ω , i.e.

$$g(x) = \begin{cases} 1, & \text{for } x \in \mathcal{A} \\ 0, & \text{for } x \in \Omega \setminus \mathcal{A}. \end{cases} \quad (3.1)$$

By the method of exponential sum, it may be expressed as

$$\begin{aligned} g(x) &= \frac{1}{q} \sum_{n=0}^{q-1} \sum_{t \in \mathcal{A}} e((x-t)n/q) = \frac{1}{q} \sum_{n=0}^{q-1} e(xn/q) \sum_{t \in \mathcal{A}} e(-tn/q) \\ &= \frac{m}{q} + \frac{1}{q} \sum_{n=1}^{q-1} \sum_{t \in \mathcal{A}} e((x-t)n/q) = \frac{m}{q} + \frac{1}{q} \sum_{n=1}^{q-1} e(xn/q) \sum_{t \in \mathcal{A}} e(-tn/q) \end{aligned} \quad (3.2)$$

Let $G(\zeta) = \sum_{x \in \mathcal{A}} e(x^k \zeta / q)$, by (3.2), it has

$$\begin{aligned}
G(\zeta) &= \sum_{x \in \Omega} e(x^k \zeta / q) \cdot g(x) = \sum_{x \in \Omega} e(x^k \zeta / q) \cdot \left(\frac{m}{q} + \frac{1}{q} \sum_{n=1}^{q-1} e(xn / q) \sum_{t \in \mathcal{A}} e(-tn / q) \right) \\
&= \frac{m}{q} \sum_{x \in \Omega} e(x^k \zeta / q) + \frac{1}{q} \sum_{x \in \Omega} \sum_{n=1}^{q-1} e((x^k \zeta + xn) / q) \sum_{t \in \mathcal{A}} e(-tn / q).
\end{aligned} \tag{3.3}$$

So, it has

$$\begin{aligned}
\psi_s(N, \mathcal{A}) &= \frac{1}{q} \sum_{\zeta=0}^{q-1} \sum_{x \in \Omega} \sum_{y \in \Omega} e((x^k + y^k - N)\zeta / q) = \frac{1}{q} \sum_{\zeta=0}^{q-1} G(\zeta)^2 e(-N\zeta / q) \\
&= \frac{1}{q} \sum_{\zeta=0}^{q-1} \left(\frac{m}{q} \sum_{x \in \Omega} e(x^k \zeta / q) + \frac{1}{q} \sum_{x \in \Omega} \sum_{a=1}^{q-1} e((x^k \zeta + xa) / q) \sum_{t \in \mathcal{A}} e(-ta / q) \right)^2 e(-N\zeta / q) \\
&= \frac{m^2}{q^3} \sum_{\zeta=0}^{q-1} \sum_{x, y \in \Omega} e((x^k + y^k - N)\zeta / q) \\
&\quad + \frac{m}{q^3} \sum_{\zeta=0}^{q-1} \sum_{x, y \in \Omega} e((x^k + y^k - N)\zeta / q) \cdot \left(\sum_{a=1}^{q-1} \sum_{t \in \mathcal{A}} e((x-t)a / q) + \sum_{a=1}^{q-1} \sum_{t \in \mathcal{A}} e((y-t)a / q) \right) \\
&\quad + \frac{1}{q^3} \sum_{\zeta=0}^{q-1} \sum_{x, y \in \Omega} \sum_{1 \leq a, b < q} \sum_{u, v \in \mathcal{A}} e((x^k + y^k - N)\zeta / q) \cdot e((a(x-u) + b(y-v)) / q) \\
&= \mathcal{L}_1 + \mathcal{L}_2 + \mathcal{L}_3
\end{aligned}$$

It is easy to know that $\mathcal{L}_1 \ll m^2 / q$,

and,

$$\begin{aligned}
\mathcal{L}_2 &= \frac{m}{q^3} \sum_{\zeta=0}^{q-1} \sum_{x, y \in \Omega} e((x^k + y^k - N)\zeta / q) \cdot \left(\sum_{a=1}^{q-1} \sum_{t \in \mathcal{A}} e((x-t)a / q) + \sum_{a=1}^{q-1} \sum_{t \in \mathcal{A}} e((y-t)a / q) \right) \\
&\leq \frac{2m}{q^3} (q-m) \sum_{\zeta=0}^{q-1} \sum_{x \in \mathcal{A}, y \in \Omega} e((x^k + y^k - N)\zeta / q) \\
&\ll \frac{m^2}{q}.
\end{aligned}$$

Main calculations are in the estimation for $\mathcal{L}_3$, the results and arguments in § 2 will be used here.

$$\begin{aligned}
\mathcal{L}_3 &= \frac{1}{q^3} \sum_{\zeta=0}^{q-1} \sum_{x, y \in \Omega} \sum_{1 \leq a, b < q} \sum_{u, v \in \mathcal{A}} e((x^k + y^k - N)\zeta / q) \cdot e((ax + by) / q) e(-(au + bv) / q) \\
&= \frac{1}{q^2} \sum_{\substack{x, y \in \Omega \\ x^k + y^k = N}} \sum_{1 \leq a, b < q} \sum_{u, v \in \mathcal{A}} e((ax + by) / q) e(-(au + bv) / q) \\
&= \frac{1}{q^2} \sum_{1 \leq a, b < q} \sum_z \sum_{(x, y) \in \mathfrak{S}(z, a, b)} \sum_{u, v \in \mathcal{A}} e(z / q) e(-w / q), \quad w = w(a, b, u, v) = au + bv. \\
&= \frac{1}{q^2} \sum_{0 \leq i < k} \sum_{1 \leq a, b < q} \sum_z \sum_{(x, y) \in \mathfrak{S}(z, a, b)} \sum_{u, v \in \mathcal{A}} e((z - w) / q) \\
&\quad p^i \parallel (a, b)
\end{aligned}$$

$$= \sum_{0 \leq i < k} \mathcal{L}_{3,i}$$

Where $\mathcal{L}_{3,i} = \frac{1}{q^2} \sum_{\substack{1 \leq a, b < q \\ p^i \parallel (a,b)}} \sum_z \sum_{(x,y) \in \mathfrak{S}(z,a,b)} \sum_{u,v \in \mathcal{A}} e((z-w)/q)$. Specially,

$$\begin{aligned} \mathcal{L}_{3,0} &= \frac{1}{q^2} \sum_{\substack{a,b,z \\ (a,b)=1}} \sum_{\substack{\mathfrak{S}(z,a,b) \\ \tau_1 \geq (k-\varepsilon)/2}} \sum_{u,v \in \mathcal{A}} e((z-w)/q) + \frac{1}{q^2} \sum_{\substack{a,b,z \\ (a,b)=1}} \sum_{\substack{\mathfrak{S}(z,a,b) \\ \tau_1 < (k-\varepsilon)/2}} \sum_{u,v \in \mathcal{A}} e((z-w)/q) \\ &= \Gamma_1 + \Gamma_2. \end{aligned}$$

We at first estimate Γ_1 .

$$\begin{aligned} \Gamma_1 &= \frac{1}{q^2} \sum_{\substack{1 \leq a < q \\ (a,p)=1}} \sum_{1 \leq b < q} \sum_z \sum_{\substack{\mathfrak{S}(z,a,b) \\ \tau_1 \geq (k-\varepsilon)/2}} \sum_{u,v \in \mathcal{A}} e((z-w)/q) \\ &= \frac{1}{q^2} \sum_{\substack{1 \leq a < q \\ (a,p)=1}} \sum_{1 \leq b < q} \sum_{\substack{z \\ \tau_1 \geq (k-\varepsilon)/2}} \sum_{u,v \in \mathcal{A}} e(a(z(b)-w(b))/q) p^r, \quad (z(b) = x + by, w(b) = u + by) \\ &= \frac{p^{r+\varepsilon/2}}{q^2} \sum_{\substack{1 \leq a < q \\ (a,p)=1}} \sum_{1 \leq b < q} \sum_{\substack{z \\ \tau_1 \geq (k+\varepsilon)/2}} \sum_{\substack{u,v \in \mathcal{A} \\ w \not\equiv z \pmod{p^{(k-1)}}}} e(a(z-w)/q) \chi(a\varrho), \quad (z := z(b), w := w(b)) \\ &\quad + \frac{p^{r+\varepsilon/2}}{q^2} \sum_{\substack{1 \leq a < q \\ (a,p)=1}} \sum_{1 \leq b < q} \sum_{\substack{z \\ \tau_1 \geq (k+\varepsilon)/2}} \sum_{\substack{u,v \in \mathcal{A} \\ w \equiv z \pmod{p^{(k-1)}}}} e(a(z-w)/q) \chi(a\varrho) \\ &\leq \frac{p^{r+\varepsilon/2}}{q^2} \sum_{\substack{1 \leq a < q \\ (a,p)=1}} \sum_{1 \leq b < q} \sum_{\substack{z \\ \tau_1 \geq (k+\varepsilon)/2}} \sum_{\substack{u,v \in \mathcal{A} \\ w \equiv z \pmod{p^{(k-1)}}}} e(a(z-w)/q) \chi(a\varrho) \\ &= \frac{p^{r+\varepsilon/2}}{q^2} \sum_{\substack{1 \leq a < q \\ (a,p)=1}} \sum_{1 \leq b < q} \sum_{\substack{z \\ \tau_1 \geq (k+\varepsilon)/2}} \sum_{\substack{u,v \in \mathcal{A} \\ w \equiv z \pmod{p^{(k-1)}} \\ y \equiv v \pmod{p^r}}} e(a(z-w)/q) \chi(a\varrho) \\ &\quad + \frac{p^{r+\varepsilon/2}}{q^2} \sum_{\substack{1 \leq a < q \\ (a,p)=1}} \sum_{1 \leq b < q} \sum_{\substack{z \\ \tau_1 \geq (k+\varepsilon)/2}} \sum_{\substack{u,v \in \mathcal{A} \\ w \equiv z \pmod{p^{(k-1)}} \\ y \not\equiv v \pmod{p^r}}} e(a(z-w)/q) \chi(a\varrho) \\ &= \frac{p^{r+\varepsilon/2}}{q^2} \sum_{\substack{1 \leq a < q \\ (a,p)=1}} \sum_{1 \leq b < q} \sum_{\substack{z \\ \tau_1 \geq (k+\varepsilon)/2}} \sum_{\substack{u,v \in \mathcal{A} \\ w \equiv z \pmod{p^{(k-1)}} \\ y \equiv v \pmod{p^r}}} e(a(z-w)/q) \chi(a\varrho) \\ &= \frac{p^{r+\varepsilon/2}}{q^2} \sum_{\substack{1 \leq a < q \\ (a,p)=1}} \sum_{1 \leq b < q} \sum_{\substack{z \\ \tau_1 \geq (k+\varepsilon)/2}} \sum_{0 \leq \alpha < p} \sum_{\mathcal{H}_\alpha} e(a(z-w)/q) \chi(a\varrho) \\ &= \frac{1}{q^2} (\Lambda_{\alpha=0} + \Lambda_{\alpha \neq 0}) \end{aligned}$$

By (2.15), we know that $\Lambda_{\sigma \neq 0} < 0$, and by (2.14), (2.16), it has

$$\Lambda_{\alpha=0} \ll p^r \cdot \frac{|\mathcal{A}|^2}{\varphi(p^k)} \varphi(p^{r+1})q \ll \left\lceil |\mathcal{A}|^2 \cdot q \right\rceil.$$

Thus,

$$\Gamma_1 \ll \left\lceil |\mathcal{A}|^2 / q \right\rceil.$$

In order to calculate Γ_2 , let $\mathfrak{D}_t = \mathfrak{D}_t(1, b) = \{(x, y) \mid (x, y) \in \mathfrak{S}(z, 1, b), \tau_1(x, y) = t\}$. It is easy

to know $|\mathfrak{D}_t| \ll p^t$. On the other hand, by Lemma 2.1, we know that for each $(x, y) \in \mathfrak{D}_t$ there

is a period of length p^{k-1-2t} and step p^{2t+1} in $\mathfrak{D}_t$, denote by $\mathcal{O}_{(x_0, y_0)}$ the orbit start

from (x_0, y_0) , hence, for each a , $(a, p) = 1$.

$$\begin{aligned} \sum_{(x, y) \in \mathfrak{D}_t} e(a(x + by) / q) &= \sum_{(x_0, y_0)} \sum_{(x, y) \in \mathcal{O}_{(x_0, y_0)}} e(a(x + by) / q) \\ &= \sum_{(x_0, y_0)} e(a(x_0 + by_0) / q) \sum_{\lambda=0}^{p^{k-2t-1}} e(a\lambda p^{2t+1} / q) \\ &= 0. \end{aligned}$$

And so,

$$\begin{aligned} \Gamma_2 &= \frac{1}{q^2} \sum_{\substack{1 \leq a < q \\ (a, p)=1}} \sum_{1 \leq b < q} \sum_z \sum_{\substack{\mathfrak{S}(z, a, b) \\ \tau_1 < (k-\varepsilon)/2}} \sum_{u, v \in \mathcal{A}} e((z - w) / q) \\ &= \frac{1}{q^2} \sum_{\substack{1 \leq a < q \\ (a, p)=1}} \sum_{1 \leq b < q} \sum_z \sum_{\tau_1 < (k-\varepsilon)/2} \sum_{\mathfrak{S}(z, a, b)} \sum_{u, v \in \mathcal{A}} e(a(z(b) - w(b)) / q) \quad (z(b) = x + by, w(b) = u + by) \\ &= \frac{1}{q^2} \sum_{\substack{1 \leq a < q \\ (a, p)=1}} \sum_{1 \leq b < q} e(-aw(b) / q) \sum_{t < (k-\varepsilon)/2} \sum_z \sum_{\substack{\mathfrak{S}(z, a, b) \\ \tau_1 = t}} e(a \cdot z(b) / q) \\ &= \frac{1}{q^2} \sum_{\substack{1 \leq a < q \\ (a, p)=1}} \sum_{1 \leq b < q} e(-aw(b) / q) \sum_{t < (k-\varepsilon)/2} \sum_{(x, y) \in \mathfrak{D}_t} e(a \cdot z(b) / q) \\ &= 0. \end{aligned}$$

By now, we have proven $\mathcal{L}_{3,0} \ll |\mathcal{A}|^2 / q$.

The proof of $\mathcal{L}_{3,i}$, $i > 0$, is similar, or applying induction, for example, for $\mathcal{L}_{3,1}$, let $\bar{q} = q / p$,

it has

$$\begin{aligned} \mathcal{L}_{3,1} &= \frac{1}{q^2} \sum_{\substack{1 \leq a < q' \\ (b, p)=1}} \sum_{1 \leq b < q'} \sum_z \sum_{(x, y) \in \mathfrak{S}(z, a, b)} \sum_{u, v \in \mathcal{A}} e((z - w) / \bar{q}) \\ &= \frac{1}{p^2} \cdot \left(\frac{1}{\bar{q}^2} \sum_{1 \leq a < \bar{q}} \sum_{1 \leq b < \bar{q}} \sum_{1 \leq z < \bar{q}} \sum_{(x, y) \in \mathfrak{S}(z, a, b)} \sum_{u, v \in \mathcal{A}} e((z - w) / \bar{q}) \right) \end{aligned}$$

$$\ll \frac{1}{p^2} \cdot \left(\frac{|\mathcal{A}|^2}{\bar{q}} \times p \right) = \frac{|\mathcal{A}|^2}{q},$$

for there are at most p pairs $(x, y) \in \mathfrak{S}(a, b, z)$ with same an image mod $\bar{q}$. Hence we have

$$\mathcal{L}_3 \ll \frac{|\mathcal{A}|^2}{q},$$

and proof is finished. □

The test data shows that the sets of prime numbers in an interval are submitted to the conclusion (1.6) very well yet, and they are uniform distributed, but it is not known whether they are strongly uniform, we guess it may be.

4. General cases

In fact, the proof for Theorem 2 is similar to the one of Theorem 1 provided with some modifications correspondingly for definitions, and the binomial expansion is replaced by Taylor expansion. For the sake of caution, we also restate them if necessary.

For $N \in \Omega(f)$, $a, b \in \mathbb{Z}_q \setminus \{0\}$, $z \in \mathbb{Z}_q$, consider following equation

$$\begin{cases} f(x) + f(y) \equiv N \pmod{q}, & x, y \in \Omega(f), \\ ax + by \equiv z \pmod{q}. \end{cases} \quad (4.1)$$

Denote by $\mathfrak{S}(z, a, b; f)$ the set of solutions (x, y) of (4.1), and with respect to $\mathfrak{S}(z, a, b; f)$, we define

$$\begin{aligned} \rho_i &= \rho_i(x, y; a, b; f) = f^{(i)}(x)b^i + (-a)^i f^{(i)}(y), \\ \tau_i &= \tau_i(x, y; a, b; f) \Leftrightarrow p^{\tau_i} \parallel \rho_i, \quad i = 1, 2, \dots \end{aligned}$$

Lemma 4.1. Suppose that $(x, y) \in \mathfrak{S}(z, a, b; f)$, $p^{r-1} \parallel \rho_1$, if $2r-1 < k$, then for arbitrary integer t , there is $(u, v) \in \mathfrak{S}(z + tp^{2r-1}, a, b; f)$, with that $u = x + cp^r$, $v = y + dp^r$.

Proof. Clearly, it suffices to show that for arbitrary integer t , there are integers c, d , such that

$$u = x + cp^r, v = y + dp^r, (u, v) \in \mathfrak{S}(z + tp^{2r-1}, a, b). \quad (4.2)$$

Then

$$\begin{aligned} au + bv &\equiv ax + by + (ac + bd)p^r \equiv w + tp^{2r-1} \pmod{q}. \\ &\Rightarrow \end{aligned}$$

$$(ac + bd)p^r \equiv tp^{2r-1} \pmod{q}$$

And

$$ac + bd \equiv tp^{r-1} \pmod{q'}, \quad q' = q / p^r.$$

Besides,

$$\begin{aligned} f(u) + f(v) &= f(x + cp^r) + f(y + dp^r) = \sum_i f^{(i)}(x)(cp^r)^i / i! + \sum_i f^{(i)}(y)(dp^r)^i / i! \\ &= (f(x) + f(y)) + (c \cdot f'(x) + d \cdot f'(y))p^r + (c^2 \cdot f''(x) + d^2 \cdot f''(y))p^{2r} / 2! + \dots \end{aligned} \quad (4.3)$$

$\Rightarrow$

$$(c \cdot f'(x) + d \cdot f'(y))p^r + (c^2 \cdot f''(x) + d^2 \cdot f''(y))p^{2r} / 2! + \dots = 0$$

And

$$c \cdot f'(x) + d \cdot f'(y) \equiv 0 \pmod{p^r}.$$

So,

$$c \cdot f'(x) + d \cdot f'(y) \equiv 0 \pmod{p^r}.$$

$$\begin{aligned} f'(x)bc + f'(y)bd &\equiv f'(x)bc + f'(y)(tp^{r-1} - ac) \\ &\equiv (f'(x)b - af'(y))c + tp^{r-1}y^{k-1} \equiv 0 \pmod{p^r} \end{aligned}$$

Suppose $(f'(x)b - af'(y)) = l \cdot p^{r-1}$, $l \not\equiv 0 \pmod{p}$. It follows

$$lc + ty^{k-1} \equiv 0 \pmod{p}. \quad (4.4)$$

Suppose $c = \sum_{i \geq 0} c_i p^{ir}$, $d = \sum_{i \geq 0} d_i p^{ir}$, from (4.4), c_0 and d_0 are followed. And then consider

the rest terms of (4.3) similarly, c_i, d_i will be followed in turn $i \geq 0$. $\square$

Lemma 4.2. Suppose $(a, b) = 1$ and $(x, y) \in \mathfrak{S}(z, a, b; f)$, $r \geq \lceil (k+1)/2 \rceil$, $p^r \parallel \rho_1$, $p \nmid \rho_2$,

let $\lambda = \lceil (k-1)/2 \rceil$, then there are $(u, v) \in \mathfrak{S}(z + tp^{2\lambda}, a, b; f)$, with that $u = x + cp^\lambda$, $v = y + dp^\lambda$, and as c is over Z_{p^e} , t will take over all quadratic (non-)residue classes, depend

on ρ_2 is or not a quadratic residue.

Proof. Assume that $(u, v) \in \mathfrak{S}(z + tp^{2\lambda}, a, b; f)$, then

$$(ac + bd)p^\lambda - tp^{2\lambda} \equiv 0 \pmod{q},$$

i.e.

$$ac + bd \equiv tp^\lambda \pmod{q'}, \quad q' = p^{k-\lambda}. \quad (4.5)$$

And

$$\begin{aligned} f(u) + f(v) &= f(x + cp^\lambda) + f(y + dp^\lambda) \\ &= \sum_i f^{(i)}(x)(cp^\lambda)^i / i! + \sum_i f^{(i)}(y)(dp^\lambda)^i / i! \\ &= (f(x) + f(y)) + (c \cdot f'(x) + d \cdot f'(y))p^\lambda + (c^2 \cdot f''(x) + d^2 \cdot f''(y))p^{2\lambda} / 2! + \dots \\ &\Rightarrow \end{aligned}$$

$$(c \cdot f'(x) + d \cdot f'(y))p^r + (c^2 \cdot f''(x) + d^2 \cdot f''(y))p^{2r} / 2! \equiv 0 \pmod{q}$$

$$\begin{aligned} b(f'(x)c + f'(y)d)p^\lambda &= (f'(x)bc + f'(y)bd)p^\lambda \\ &\equiv (f'(x)bc + f'(y)(tp^\lambda - ac))p^\lambda \equiv ((f'(x)b - af'(y))c + tp^\lambda y^{k-1})p^\lambda \pmod{q} \\ &\equiv ty^{k-1}p^{2\lambda} \pmod{q}. \end{aligned}$$

Hence,

$$(c^2 \cdot f''(x) + d^2 \cdot f''(y))p^{2\lambda} / 2! + ty^{k-1}p^{2\lambda} / b \equiv 0 \pmod{q}. \quad (4.6)$$

From (4.5), we have

$$(ac)^2 - (bd)^2 \equiv \mu p^\lambda \pmod{q'} \quad (4.7)$$

With (4.6), (4.7), it has

$$b^2 \cdot (f''(x)c^2 + f''(y)d^2)p^{2\lambda} = (f''(x)b^2c^2 + f''(y)((ac)^2 - \mu p^\lambda) + \omega)p^{2\lambda} \equiv 0 \pmod{q},$$

where $\omega \equiv ty^{k-1}2b$. i.e.

$$\begin{aligned} (f''(x)b^2 + f''(y)a^2)c^2 + \omega &\equiv 0 \pmod{p^\epsilon} \\ t &\equiv \varrho c^2 \pmod{p}, \quad \varrho = -\rho_2 / (2by^{k-1}). \end{aligned}$$

For $\varrho \not\equiv 0 \pmod{p}$, so t will take over all quadratic (non-)residue classes as c is from 1 to p^ϵ ,

rely on ϱ is a quadratic residue or not. □

Define

$$\mathfrak{S}'(z, a, b; f) = \{ (x, y) \mid (x, y) \in \mathfrak{S}(z, a, b; f), \tau_1 \geq (k + \varepsilon) / 2 \}.$$

Lemma 4.3. Suppose that $(a, b) = 1$, and $(x, y) \in \mathfrak{S}'(z, a, b; f)$, then there are just

$$\begin{aligned} p^{(k-\varepsilon)/2} \text{ pairs } (x_i, y_i) &\in \mathfrak{S}'(z, a, b), 1 \leq i \leq p^{(k-\varepsilon)/2}, \text{ with that } x_i \equiv x \pmod{p^{(k+\varepsilon)/2}}, y_i \equiv y \\ &\pmod{p^{(k+\varepsilon)/2}}. \end{aligned}$$

Proof. For $\tau_1 \geq (k + \varepsilon)/2$, this indicates that polynomial $\Gamma(x) = f(x) + f((z - ax)/b) - N$ has a duplicative root $x_0 \pmod{p^{\tau_1}}$, then it is easy to know that $x_0 + \lambda p^{(k+\varepsilon)/2}$, $0 \leq \lambda < p^{(k-\varepsilon)/2}$ will be the roots of polynomial $\Gamma(x)$.

□

Suppose that $(x, y) \in \mathfrak{S}'(z, 1, b; f)$ and $(x', y') \in \mathfrak{S}'(z', 1, b'; f)$, $x' = x + \Delta x$, $y' = y + \Delta y$,

$b' = b + \Delta b$. If $p^r \mid \Delta x$, then it is easy to know that $p^r \mid \Delta y$, $p^r \mid \Delta b$.

Suppose that $\Delta x = \lambda p^r$, and $r \geq (k - \varepsilon)/2$, then by Taylor expansion, it has

$$\begin{aligned} f(x + \Delta x) &= f(x) + \Delta x f'(x) + (\Delta x)^2 f''(x)/2 \\ f(y + \Delta y) &= f(y) + \Delta y f'(y) + (\Delta y)^2 f''(y)/2 \\ \Rightarrow \\ 2\Delta x f'(x) + (\Delta x)^2 f''(x) + 2\Delta y f'(y) + (\Delta y)^2 f''(y) &= 0 \end{aligned}$$

It follows

$$\Delta y = -\frac{\Delta x}{b} - (\Delta x)^2 \frac{f''(y) + b^2 f''(x)}{2b^2 f'(y)} \quad (4.8)$$

By $(b + \Delta b) f'(x + \Delta x) - f'(y + \Delta y) = 0$, it follows

$$(\Delta y f''(y) + (\Delta y)^2 f'''(y)/2) = b \Delta x f''(x) + b(\Delta x)^2 f'''(x)/2 + \Delta b f'(x) + (\Delta b)(\Delta x) f''(x)$$

Specially,

$$\Delta y f''(y) - b \Delta x f''(x) \equiv \Delta b f'(x) \pmod{p^{k-\varepsilon}} \quad (4.9)$$

With (4.8) and (4.9), it has

$$\Delta x \equiv -\Delta b \frac{f'(y)}{\rho_2} \pmod{p^{k-\varepsilon}}, \quad \Delta y \equiv \Delta b \frac{f'(x)}{\rho_2} \pmod{p^{k-\varepsilon}}. \quad (4.10)$$

And

$$\begin{aligned} z' &= (x + \Delta x) + (b + \Delta b)(y + \Delta y) \\ &= z + \Delta b y + \Delta x + (b + \Delta b)\Delta y \\ &= z + \Delta b y + \Delta x - (b + \Delta b) \left(\frac{\Delta x}{b} + \frac{\rho_2 (\Delta x)^2}{2b^2 f'(y)} \right) \\ &= z + \Delta b y - \frac{\Delta x \Delta b}{b} - \frac{\rho_2}{2b f'(y)} (\Delta x)^2 \end{aligned}$$

$$\begin{aligned}
&= z + \Delta by - \left((\Delta b)^2 \left(\frac{f'(y)}{\rho_2} \right)^2 \frac{\rho_2}{2bf'(y)} - \frac{(\Delta b)^2 f'(y)}{b\rho_2} \right) \\
&= z + \Delta by + (\Delta b)^2 \frac{f'(y)}{2b\rho_2}.
\end{aligned} \tag{4.11}$$

Hence, we have

Lemma 4.4. Suppose that $(a, b) = 1, (x, y) \in \mathfrak{S}(z, a, b; f)$, $\rho_1(x, y; a, b) = 0$. And $\lambda \in \mathbb{Z}_q$, let $b' = b + \lambda p^{(k-\epsilon)/2}$, then there is $(x', y') \in \mathfrak{S}(z', a, b'; f)$, with that $x' \equiv x \pmod{p^{(k-\epsilon)/2}}$, $y' \equiv y \pmod{p^{(k-\epsilon)/2}}$, $z' \equiv z + a\lambda y p^{(k-\epsilon)/2} \pmod{p^{(k-\epsilon)}}$.

Lemma 4.5. Suppose $u, v \in \mathcal{A}$, $(x, y) \in \mathfrak{S}'(z, 1, b; f)$, and $y \equiv v \pmod{p^r}$, $x \equiv u \pmod{p^r}$, $r \geq (k - \epsilon)/2$, $(x - u) + b(y - v) \equiv 0 \pmod{p^{k-1}}$, then (x, y) is homotopic to a $(\tilde{x}, \tilde{y})$ with that $\tilde{y} \equiv v \pmod{p^{k-1}}$, $\tilde{x} \equiv u \pmod{p^{k-1}}$, $(\tilde{x} - u) + \tilde{b}(\tilde{y} - v) \equiv \alpha p^{k-1} \pmod{q}$.

In order to obtain an analogue of Lemma 2.7, we need following lemma

Lemma 4.6. Let $\mathfrak{R}(x)$ be the ring of polynomials over $\mathbb{Z}_q$, suppose that $f(x), g(x) \in \mathfrak{R}(x)$, with the first coefficients coprime to p , $\deg(f) = k$, $\deg(g) = k - 1$, then there is a non-zero bi-variables polynomial $F(u, v)$ over $\mathbb{Z}_q$ such that $F(f, g) = 0$.

Proof. For the simplicity, we assume that $f(x), g(x)$ are monic. Let

$$\mathfrak{I}(f, g) = \left\{ \sum_{i,j} a_{i,j} f^i(x) g^j(x) \mid a_{i,j} \in \mathbb{Z}_q \right\}, \text{ and } \mathfrak{I}_r = \{ \zeta(x) \mid \zeta(x) \in \mathfrak{I}(f, g), \deg(\zeta) = r \}.$$

suppose that r_0 is the least integer r such that $\mathfrak{I}_r \neq \emptyset$. if $r_0 = 0$, then the proof is over. So, assume $r_0 > 0$, and so $\text{rank}(\mathfrak{I}_{r_0}) = 1$. Suppose that r_1 is the least integer r with $r_1 > r_0$ such that $\mathfrak{I}_r \neq \emptyset$. Similarly, $\text{rank}(\mathfrak{I}_{r_1}) \leq 2$, and so on. On the other hand, $f^{\lambda \deg(g)}$ and $g^{\lambda \deg(f)}$ are two polynomials with same degrees, let $\Gamma_\lambda(x) =$

$f(x)^{\lambda \deg(g)} - g(x)^{\lambda \deg(f)}$, then $\deg(\Gamma_\lambda(x)) < \lambda k(k-1)$. Suppose that

$$\mu(k-1) \leq \deg(\Gamma_\lambda(x)) < \mu k, \text{ say, } \deg(\Gamma_\lambda(x)) = \mu(k-1) + a, \quad 0 \leq a \leq \mu.$$

Then there is a constant $c \in \mathbb{Z}_q$, $\deg(\Gamma_\lambda(x) - c \cdot g^{\mu-a} f^a) < \deg(\Gamma_\lambda(x))$, and so on, it follows that there is $\Lambda_\lambda(x) = \sum_{i,j} a_{i,j} f^i(x) g^j(x)$, $\Delta_\lambda(x) = \Gamma_\lambda(x) - \Lambda_\lambda(x)$, with that $\deg(\Delta_\lambda(x)) < k(k-1)$. By the argument above, we know that $\{\Delta_\lambda(x)\}_1^{k(k-1)}$ are linear dependent, i.e. there are constants $c_\lambda \in \mathbb{Z}_q, 1 \leq \lambda \leq k(k-1)$, such that $\sum_{1 \leq \lambda \leq k(k-1)} c_\lambda \Delta_\lambda(x) = 0$.

Lemma 4.7. Let $\Theta(r)$ as defined in Lemma 2.7, then

$$\begin{aligned} \Theta(r) &\ll \left\lceil |\mathcal{A}|^2 / \varphi(p^r) \right\rceil, \\ \Theta(r)|_\delta &\ll \left\lceil |\mathcal{A}|^2 / \varphi(p^{r+d}) \right\rceil. \end{aligned} \tag{4.12}$$

Proof. Let $r_0 = \lceil \log_p |\mathcal{A}| - 1 \rceil$, if $r \leq r_0$, by the uniformity of set $\mathcal{A}$, then

$$\begin{aligned} \Theta(r) &\ll \varphi(p^r) \cdot \left(|\mathcal{A}| / \varphi(p^r) \right)^2 = |\mathcal{A}|^2 / \varphi(p^r), \text{ and} \\ \Theta(r)|_\delta &\ll \left\lceil |\mathcal{A}|^2 / \varphi(p^{r+d}) \right\rceil. \end{aligned}$$

So, assume that $r > r_0$. Suppose that $|\mathcal{A}| \doteq s \cdot p^{r_0}, \bar{q} = p^{r_0}$, and by the definition, without loss the generality, let

$$\mathcal{A}_{\bar{q}} = \{\lambda_i\}_1^s, \quad \lambda_i = \sum_{t \geq 0} c_{i,t} p^t, \quad 0 \leq c_{i,t} < p, \quad 1 \leq i \leq s.$$

For a integer $l, 1 \leq l \leq r$, let $\Omega_l = \{x \mid 0 < x < p^l, (x, p) = 1\}$, considering the equation

$$\begin{cases} f(x) + f(y) \equiv N \pmod{p^l}, & x, y \in \Omega_l \\ bf'(x) - f'(y) \equiv 0 \pmod{p^l}. \end{cases} \tag{4.13}$$

Denote by $\mathfrak{M}_l$ the set of solutions of equation (4.13). For a pair $(x, y) \in \mathfrak{M}_l$, by Lemma above,

we know that there is a bi-variables polynomial $F(u, v)$ such that $F(f(y), f'(y)) = 0$, then

$$F(N - f(x), bf'(x)) \equiv 0 \pmod{p^l}$$

This means that there are

$$[x]_l = \phi_1([b]_l, [N]_l), \quad [y]_l = \phi_2([b]_l, [N]_l). \tag{4.14}$$

But the function relations are possibly implicit.

Moreover, for a integer $h, 0 \leq h < p^{r-l}$, define

$$\mathfrak{M}_l(h, r) = \left\{ (x, y) \mid (x, y) \in \mathfrak{M}_l, f(x) + f(y) \equiv [N]_l + h \cdot p^l \pmod{p^r} \right\}.$$

It is clear that

$$\mathfrak{M}_l = \bigcup_{0 \leq h < p^{r-l}} \mathfrak{M}_l(h, r). \quad (4.15)$$

Clearly,

$$[x]_l = [x]_{r_0} + \sum_{r_0 \leq i < l} x_i p^i, \quad [y]_l = [y]_{r_0} + \sum_{r_0 \leq i < l} y_i p^i, \quad \text{for } l \geq r_0. \quad (4.16)$$

Furthermore, for a pair $(i, j), 1 \leq i, j \leq s$, in the expression (4.16), let $x_{r_0+t} = c_{i,t}, y_{r_0+t} = c_{j,t}$,

$0 \leq t < r - r_0$. i.e.

$$[x]_l = [x]_{r_0} + \sum_{0 \leq t < l - r_0} c_{i,t} p^{r_0+t}, \quad [y]_l = [y]_{r_0} + \sum_{0 \leq t < l - r_0} c_{j,t} p^{r_0+t}. \quad (4.17)$$

With (4.14), it has

$$[x]_l = \phi_1([b]_l, [N]_l) + \sum_{0 \leq t < l - r_0} c_{i,t} p^{r_0+t}, \quad [y]_l = \phi_2([b]_l, [N]_l) + \sum_{0 \leq t < l - r_0} c_{j,t} p^{r_0+t}. \quad (4.18)$$

Substituting (4.18) into the first equation of (4.13), and taking $l = r_0 + 1, r_0 + 2, \dots, r$ respectively, by Taylor expansion, then in each set $\mathfrak{M}_{r_0}(h, r), b_i, i = 0, 1, \dots, (r - r_0 - 1)$, will be determined in turn. Moreover, let $d = r - r_0$, by the induction on the second expression of (4.12), roughly speaking, the sizes of sets in $\left\{ \mathfrak{M}_{r_0}(h, r) \Big|_{\delta} \mid 0 \leq \delta < p^d \right\}$ are about equal. Therefore, there are at most p^{2r_0-r} solutions for the variables b_i 's.

For there are s^2 selections of pair $(\lambda_i, \lambda_j), 1 \leq i, j \leq s$. Hence

$$\Theta(r) \ll s^2 \cdot p^{2r_0-r} \ll \left\lceil |\mathcal{A}|^2 / p^r \right\rceil.$$

□

The rest proof of Theorem 2 is almost same as the one of Theorem 1, so is omitted.

Finally, maybe it should be mentioned that, in practical applying, we find that Theorem 2 seems more accurate in the cases that constant $a_n \equiv 0 \pmod{p}$. The first interpretation for the phenomenon is that equation $f(x) + f(y) \equiv N \pmod{q}$ can be transferred as $(f(x) - a_n) +$

$(f(y) - a_n) \equiv (N - 2a_n) \pmod{q}$, where $f(x) = \sum_{0 \leq i \leq n} a_i x^{n-i}$, for $N - 2a_n$ will be out of Ω ,

as N takes over Ω , if $a_n \not\equiv 0 \pmod{p}$.

References

1. G. H. Hardy and J. E. Littlewood, Some problems of "Partitio Numerorum": IV. The singular series in Waring's Problem and the value of the number $G(k)$, Math. Z. 12 (1922), 161-188.
2. _____, Some problems of "Partitio Numerorum" (VI): Further researches in Waring's problem, Math. Z. 23 (1925), 1-37.
3. _____, Some problems of "Partitio Numerorum" (VIII): The number $\Gamma(k)$ in Waring's problem, Proc. London Math. Soc. (2) 28 (1928), 518-542.
4. Vaughan and T.D. Wooley, Waring's problem: A survey, Number Theory for the Millennium III, A. K. Peters, 2002, 301-340.