arXiv:2004.09245v2 [math.NT] 22 Apr 2020

A power APN function CCZ-equivalent to Kasami function in even dimension

A. Musukwa[1] and M. Sala[2]

[1]Mzuzu University, P/Bag 201, Mzuzu 2, Malawi
augustinemusukwa@gmail.com

[2]University of Trento, Via Sommarive, 14, 38123 Povo, Trento, Italy
maxsalacodes@gmail.com

Abstract

Let n be an even number such that $n \equiv 0 \pmod 4$. We show that a power function x^d, with $d = 2^{\frac{n+2}{2}} + 2^{\frac{n-2}{2}} - 1$, on $\mathbb{F}_{2^n}$ is an APN function of degree $n/2$ which is CCZ-equivalent to Kasami functions of degrees $n/2$ and $(n+4)/2$.

Keywords: Power function; APN function; CCZ-equivalence

MSC 2010: 06E30, 94A60, 14G50

1 Introduction

A mapping F from $\mathbb{F}_{2^n}$ to itself is represented by a univariate polynomial which is given by $F(x) = \sum_{i=0}^{2^n-1} \delta_i x^i$, with $\delta_i \in \mathbb{F}_{2^n}$. Two functions F and H from $\mathbb{F}_{2^n}$ to itself are called *Carlet-Charpin-Zinoviev equivalent* (CCZ-equivalent) if, for some affine permutation $\mathcal{L}$ of $\mathbb{F}_{2^n} \times \mathbb{F}_{2^n}$, the image of the graph F is the graph H, that is, $\mathcal{L}(F) = H$, where $G_F = \{(x, F(x)) : x \in \mathbb{F}_{2^n}\}$ and $G_H = \{(x, H(x)) : x \in \mathbb{F}_{2^n}\}$.

A function F is called an *almost perfect nonlinear (APN)* if each equation given by $F(t) + F(t+a) = b$, with $a \in \mathbb{F}_{2^n} \setminus \{0\}$ and $b \in \mathbb{F}_{2^n}$, has at most two solutions t in $\mathbb{F}_{2^n}$. If F is defined by x^d then it is called a *power function*. If x^d is APN then $x^{d'}$ is

also APN for $d' \equiv 2^i d \pmod{2^n - 1}$, with $0 \leq i < n$, and for $d' = d^{-1} \pmod{2^n - 1}$ in case $\gcd(d, 2^n - 1) = 1$ (this happens only when n is odd) (see [6]).

Power APN functions have been extensively studied and discovered; the last one recorded was by Dobbertin in 2001 (see [7]). Many authors, including Beth and Ding in [1] and Canteaut, Charpin and Dobbertin in [2], have studied the application of these functions to cryptography. It has been and is still a big question as to whether all power APN functions discovered so far are complete. Dobbertin asked in [7] whether Table 1 is complete up to equivalence. We know from computational results done by Anne Canteaut that Table 1 is (up equivalence) complete for $n \leq 25$.

In this paper, we shall prove that if n is an even number such that $n \equiv 0 \pmod 4$, then a power function on $\mathbb{F}_{2^n}$ with the exponent $d = 2^{\frac{n+2}{2}} + 2^{\frac{n-2}{2}} - 1$ is an APN function of degree $n/2$ and it is CCZ-equivalent to Kasami functions of degrees $n/2$ and $(n+4)/2$.

It was observed in [3, 6] that we can show that a power function x^d in $\mathbb{F}_{2^n}[x]$ is APN by simply determining that the polynomial

$$\Phi(t) = t^d + (t+1)^d$$

defines a two-to-one mapping on $\mathbb{F}_{2^n}$.

In the table below we list all the known power APN functions.

Family	Exponent	Condition	Algebraic degree	Proven in
Gold	$2^i + 1$	$(i, n) = 1$	3	[8]
Kasami	$2^{2i} - 2^i + 1$	$(i, n) = 1$	$i + 1$	[5, 9]
Welch	$2^i + 3$	$n = 2i + 1$	3	[5]
Niho	$2^i + 2^{\frac{i}{2}} - 1, i$ even $2^i + 2^{\frac{3i+1}{2}} - 1, i$ odd	$n = 2i + 1$	$(i+2)/2$ $i + 1$	[6]
Inverse	$2^{2i} - 1$	$n = 2i + 1$	$n - 1$	[10]
Dobbertin	$2^{4i} + 2^{3i} + 2^{2i} + 2^i - 1$	$n = 5i$	$i + 3$	[7]

Table 1: Known power APN functions

Observe that if Table 1 is complete (up equivalence), then any power APN function in even dimension must be equivalent to Dobbertin, Gold or Kasami.

2 A power function with exponent $d = 2^{\frac{n+2}{2}} + 2^{\frac{n-2}{2}} - 1$

Let n be an even number such that $n \equiv 0 \pmod 4$ and let $d = 2^{\frac{n+2}{2}} + 2^{\frac{n-2}{2}} - 1$ be an exponent of a power function on $\mathbb{F}_{2^n}$. If we write $n = 2m$, then it is clear that m is even and d can as well be written as $d = 2^{m+1} + 2^{m-1} - 1$. The latter description of d is used in the rest of the paper. In the result that follows, it is shown that d is an exponent of a power APN function.

Theorem 1. *Let $d = 2^{m+1} + 2^{m-1} - 1$, with m even. Then a power function x^d on $\mathbb{F}_{2^{2m}}$ is an APN function of degree m.*

Proof. First observe that we can write

$$d = 2^{m+1} + 2^{m-1} - 1 = 2^{m+1} + 2^{m-2} + 2^{m-3} + \cdots + 2 + 1 \tag{2.1}$$

so that it is directly deduced from (2.1) that the power function x^d is of degree m.

Next we simply need to show that the polynomial

$$\Phi(t) = t^d + (t+1)^d \tag{2.2}$$

defines a two-to-one mapping on $\mathbb{F}_{2^{2m}}$.

Let $\varphi(t) = \sum_{i=0}^{e} t^i$ and $\phi(t) = \sum_{i=0}^{e-1} t^i = (t^e+1)/(t+1)$, with $e = 2^{m-1}-1$. Observe that we can write $\varphi(t) = t\phi(t) + 1$ or $\varphi(t) = t^e + \phi(t)$. So we have

$$\begin{aligned}
\Phi(t) &= t^{2^{m+1}+2^{m-1}-1} + (t+1)^{2^{m+1}+2^{m-1}-1} \\
&= t^{2^{m+1}+2^{m-1}-1} + (t+1)^{2^{m+1}}(t+1)^{2^{m-1}-1} \\
&= t^{2^{m+1}+2^{m-1}-1} + (t^{2^{m+1}}+1)(t+1)^{2^{m-2}+2^{m-3}+\cdots+2+1} \\
&= t^{2^{m+1}+2^{m-1}-1} + (t^{2^{m+1}}+1)\prod_{i=0}^{m-2}(t^{2^i}+1) \\
&= t^{2^{m+1}+2^{m-1}-1} + (t^{2^{m+1}}+1)\varphi(t) \\
&= t^{2^{m+1}+2^{m-1}-1} + (t^{2^{m+1}}+1)\left(t^e + \phi(t)\right) \\
&= t^{2^{m+1}+2^{m-1}-1} + t^{2^{m+1}+e} + t^{2^{m+1}}\phi(t) + (t^e + \phi(t)) \\
&= t^{2^{m+1}}\phi(t) + \varphi(t) = t^{2^{m+1}}\phi(t) + t\phi(t) + 1 \\
&= \phi(t)(t^{2^{m+1}} + t) + 1 = \frac{(t^e+1)(t^{2^{m+1}}+t)}{(t+1)} + 1
\end{aligned}$$

$$= \frac{(t^{2^{m-1}-1}+1)(t^{2^{m+1}}+t)}{(t+1)} + 1$$
$$= \frac{(t^{2^{m-1}}+t)(t^{2^{m+1}}+t)}{(t^2+t)} + 1. \tag{2.3}$$

Since $(m-1, 2m) = (m+1, 2m) = 1$, then we conclude from (2.3) that $\Phi(\alpha) = \Phi(\beta)$ if and only if $\beta = \alpha + c$, for $\alpha, \beta \in \mathbb{F}_{2^{2m}}$ and $c \in \mathbb{F}_2$. Hence $\Phi(t)$ is a two-to-one mapping. □

3 CCZ-equivalent to Kasami functions

In this section, we shall show that the power function which was shown to be APN in Section 2 is CCZ-equivalent to Kasami functions.

It is well-known from recent work of Dempwolff in [4] and Yoshiara in [11] who independently proved that if any two power APN functions are CCZ-equivalent then they are cyclotomic equivalent (that is, their powers are either in the same cyclotomic cosets or in the cosets of their inverses). In our case, since we are dealing with even dimension, and power APN functions, there are no permutations. So we simply have to check whether for a power APN function x^d on $\mathbb{F}_{2^n}$, we have $d \cdot 2^j \pmod{2^n - 1} \equiv d'$, for some $0 \leq j \leq n-1$, and d' is Kasami exponent.

Theorem 2. *The power function $x^{2^{m+1}+2^{m-1}-1}$ of degree m on $\mathbb{F}_{2^{2m}}$, with m even, is CCZ-equivalent to Kasami functions of degrees m and $m+2$.*

Proof. In Theorem 1, we established that the power function $x^{2^{m+1}+2^{m-1}-1}$ is an APN function of degree m. Let $d = 2^{m+1} + 2^{m-1} - 1$. To show that it is CCZ-equivalent to Kasami functions, we shall simply show, as noted above, that $d \cdot 2^j \pmod{2^{2m} - 1} \equiv d'$, for some $0 \leq j \leq 2m-1$, where d' is a Kasami exponent. Let $j = m - 1$. So we have

$$\begin{aligned} d \cdot 2^j &= 2^j(2^{m+1} + 2^{m-1} - 1) \\ &= 2^{m+j+1} + 2^{m+j-1} - 2^j \\ &= (2^{2m} + 2^{2m-2} - 2^{m-1}) \pmod{2^{2m} - 1} \\ &\equiv 2^{2m-2} - 2^{m-1} + 1 \\ &= 2^{2j} - 2^j + 1 = d'. \end{aligned}$$

Since we have $(j, 2m) = 1$, so clearly d' is a Kasami exponent and $x^{d'}$ is of degree $j + 1 = m$.

To show that it is also equivalent to Kasami function of degree $m + 2$, we now let $j = m + 1$ and, in similar manner, we have

$$\begin{aligned} d \cdot 2^j &= 2^j(2^{m+1} + 2^{m-1} - 1) = (2^{2m+2} + 2^{2m} - 2^{m+1}) \pmod{2^{2m} - 1} \\ &\equiv 2^{2j} - 2^j + 1 = d''. \end{aligned}$$

And since $(j, 2m) = 1$, then clearly $x^{d''}$ is a Kasami function of degree $j + 1 = m + 2$. □

4 Conclusion

In this paper, we have shown that a power function on $\mathbb{F}_{2^{2m}}$, with the exponent $2^{m+1} + 2^{m-1} + 1$ and m even, is an APN function of degree m and it is CCZ-equivalent to Kasami functions of degrees m and $m + 2$.

References

[1] Beth, T., Ding, C.: On almost perfect nonlinear permutations. *Advances in Cryptology-EUROCRYPT '93,* T. Helleseth (ed.), Lecture Notes in Computer Science, vol. 765, pp. 65-76, Springer-Verlag, 1994.

[2] Canteaut, A., Charpin, P., Dobbertin, H.: Weight Divisibility of Cyclic Codes, Highly Nonlinear Functions on $\mathbb{F}_{2^n}$, and Crosscorrelation of Maximum-Length Sequences. *SIAM J. Discrete Math.*, 13(1), (2000), 105138.

[3] Carlet, C.: Vectorial Boolean Functions for Cryptography. *Boolean models and methods in mathematics, computer science and engineering*, Cambridge Univ. Press, 2,(2010), 398-470.

[4] Dempwolff, U.: CCZ-equivalence of power functions. *Designs, Codes and Cryptography*, 86 (3) (2018), 665692

[5] Dobbertin, H.: Almost perfect nonlinear power functions on $GF(2^n)$: the Welch case. *IEEE Trans. Inf. Theory*, 45 (4) (1999), 1271-1275.

[6] Dobbertin, H.: Almost perfect nonlinear power functions on $GF(2^n)$: the Niho case. *Inf. Comput.*, 151 (1-2) (1999), 57-72.

[7] Dobbertin, H.: Almost perfect nonlinear power functions on $GF(2^n)$: a new case for n divisible by 5. In Jungnickel, D.Niederreiter, H. (Eds.): *Proceedings of the Conference on Finite Fields and Applications*. Augsburg, 1999, Springer-Verlag, Berlin (2001), pp. 113-121.

[8] Gold, R.: Maximal recursive sequences with 3-valued recursive cross-correlation functions. *IEEE Trans. Inf. Theory*, 14 (1968), 377-385.

[9] Kasami,T.: The weight enumerators for several classes of subcodes of the 2nd order binary ReedMuller codes. *Inf. Control*, 18 (1971), 369-394.

[10] Nyberg, K.: Differentially uniform mappings for cryptography. In Helleseth, T. (Ed.): *Advances in Cryptology - EUROCRYPT'93*, LNCS 765. Springer-Verlag Berlin Heidelberg, (1994), pp. 55-64.

[11] Yoshiara, S.: Equivalences of power APN functions with power or quadratic APN functions. *Journal of Algebraic Combinatorics*, 44 (3) (2016), 561-585.