

On Sombor index of graphs with a given number of cut-vertices

Sakander Hayat

Faculty of Engineering Sciences,
GIK Institute of Engineering Sciences and Technology,
Topi, Swabi, 23460, Pakistan
e-mail: sakander1566@gmail.com

Ansar Rehman

Faculty of Engineering Sciences,
GIK Institute of Engineering Sciences and Technology,
Topi, Swabi, 23460, Pakistan
e-mail: ges2115@giki.edu.pk

Yubin Zhong

School of Mathematics and Information Sciences,
Guangzhou University, Guangzhou,
Guangdong 510006 P.R. China
e-mail: zhong_yb@163.com

Abstract

Introduced by Gutman in 2021, the Sombor index is a novel graph-theoretic topological descriptor possessing potential applications in the modeling of thermodynamic properties of compounds. Let $\mathbb{G}_n^k$ be the set of all n -vertex connected graphs with k cut-vertices. In this paper, we present minimum Sombor indices of graphs in $\mathbb{G}_n^k$. The corresponding extremal graphs have been characterized as well.

2020 Mathematics Subject Classification: 05C92, 05C09, 05C35

Keywords: Sombor index; Graph; Cut-vertex; Block; Pendant tree; Extremal values

1 Introduction

We consider finite, connected and simple graphs only.

In modern chemistry, it is a cornerstone idea that the structure (molecular) of a chemical compound comprises information on the physicochemical characteristics of the respective compound. Topological indices are numerical quantities which provide efficient tools to retrieve this structural information of compounds. They have diverse applications in materials science, pharmacology, chemistry. among other, [12, 29]. Recently in 2021, Gutman [9] introduced a novel degree-based topological descriptors called the Sombor index. For a graph Γ , it is defined as follows:

$$SO(\Gamma) = \sum_{xy \in E(\Gamma)} \sqrt{\deg_{\Gamma}(x)^2 + \deg_{\Gamma}(y)^2},$$

where $\deg_{\Gamma}(x)$ (resp. $\deg_{\Gamma}(y)$) is the degree of the vertex x (resp. y) in Γ .

Within a short span of time, the Sombor index has attracted a significant amount of attention among researchers from mathematics and theoretical chemistry. Redžepović [25] investigated predictive potential of the Sombor index for statistical modeling of enthalpy of vaporization and entropy for alkanes. Successful predictive potential of the Sombor index has also been employed for simulating the thermodynamic characteristics of organic compounds. The reader is referred to [1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 13, 16, 17, 18, 19, 20, 21, 22, 23, 24, 26, 27, 30], for a detailed mathematical treatment of the Sombor index of graphs.

A graph Γ is an ordered pair $\Gamma = (V, E)$, where V is the set of point called vertices and $E \subseteq \binom{V}{2}$ is the set of lines called edges. The cardinality $n = |V|$ (resp. $\varepsilon = |E|$) of E is called the order (resp. size) of Γ . Two vertices $y, z \in V(\Gamma)$ are said to be adjacent or neighbors if $yz \in E(\Gamma)$. The set $N_{\Gamma}(x) = \{y \in V(\Gamma) \mid xy \in E(\Gamma)\}$ or $N(x)$ for short, is called the neighborhood of the vertex x in Γ . The number $\deg_{\Gamma}(x) = |N_{\Gamma}(x)|$ is said to be the degree of x . For an edge $xy \in E(\Gamma)$, the graph $\Gamma - xy$ is obtained by deleting $xy \in E(\Gamma)$ from Γ . On other hand, the subgraph $\Gamma - x$ for $x \in V(\Gamma)$ is obtained by deleting x and its incident edges from Γ . We denote the n -vertex path by P_n .

Investigating mathematical behavior of topological indices is a contemporary research topic these days. Determining extremal values of Sombor indices of graphs with a given property has been started recently. For instance, Sun & Du [28] studied extremal values of the Sombor index of graphs with given domination number. Zhou et al. [33, 34] investigated the extremal Sombor index of trees & unicyclic graphs with given matching number and maximum degree. On the other hand, studying extremal topological indices of graphs with given number of cut-vertices has also been studied frequently. For instance, Hua & Zhang [14] studied extremal Merrifield-Simmons index of graphs with given number of cut-vertices. Xu et al. [32] studied minimum Kirchhoff index of graphs with a given number of cut vertices. Ji & Wang recently studied minimum multiplicative Zagreb indices of graphs with given number of cut-vertices. In this paper, we study the minimum Sombor index of graphs with a given number of cut-vertices. Corresponding extremal graphs have also been characterized.

The organization of the paper is as follows: Section 2 (resp. 3) studied minimum trees (resp. non-tree graphs) with given number of cut-vertices.

2 The case of trees

Let $\mathbb{T}_n^k$ be the set of n -vertex trees having k number of cut-vertices. We find a sharp lower bound on the Sombor index of a tree in $\mathbb{T}_n^k$. Moreover, we characterize the case where the lower bound is achieved.

The following is an important result in graph theory connecting the concepts of a bridge and a cut-vertex.

Theorem 1 [31] *Let x be a vertex incident with a bridge in a connected graph Γ . Then x is a cut-vertex of Γ if and only if $\deg_{\Gamma}(x) \geq 2$.*

A vertex of degree 1 in Γ is said to be an end-vertex of Γ . Since every edge in a tree is a bridge, the following is an immediate corollary of Theorem 1.

Corollary 1 *If a vertex x of a nontrivial tree T is not an end-vertex of T , then x is a cut-vertex of T .*

The following remark follows from Corollary 1.

Remark 1 *Corollary 1 implies that for a tree $T \in \mathbb{T}_n^k$, if $k = n - 2$, then $T \cong P_n$.*

Here we first explain the τ -operation on a tree $T \in \mathbb{T}_n^k$. Let $T \in \mathbb{T}_n^k$ be a tree with vertices $u_1, v, w_1, x_1, y_1, w_k \in V(T)$ such that $\deg_T(v) \geq 3$, $u_1v, vw_1, vy_1, vx_1 \in E(T)$, and T_0 (resp. T_1) is a subtree attached with x_1 (resp. y_1). See Figure 1. Let $T_\tau = T - \{vx_1\} + \{w_kx_1\}$. Then, we say that T_τ is the τ -switched tree obtained from T by applying τ -operation.




Figure 1: The τ -operation used in Theorem 1.

The following lemma compares the Sombor index of a tree $T \in \mathbb{T}_n^k$ and T_τ .

Lemma 1 *Let T_τ be the τ -switched tree of a tree $T \in \mathbb{T}_n^k$ described in Figure 1. Then*

$$SO(T_\tau) < SO(T).$$

Proof: The structure of T in Figure 1 assumes vertices $u_1, v, w_1, x_1, y_1, w_k \in V(T)$ such that $\deg_T(v) \geq 3$, $u_1v, vw_1, vy_1, vx_1 \in E(T)$ and $\deg_T(w_k) = 1$. Moreover, let T_0 (resp. T_1) be a subtree identified to x_1 (resp. y_1). Let $N_T(v) = \{u_1, w_1, x_1, y_1\}$ be the neighborhood set for v in T (see Figure 1).

The graph T_τ is constructed by deleting $vx_1 \in E(T)$ and adding $w_kx_1 \in E(T)$. This operation would shift the tree T_0 associated with x_1 to the vertex w_k . Taking this shifting into account, the following relation between Sombor indices of T and T_τ emerges:

$$\begin{aligned} SO(T) - SO(T_\tau) &= \sqrt{\deg_T(v)^2 + \deg_T(u_1)^2} + \sqrt{\deg_T(v)^2 + \deg_T(w_1)^2} + \\ &\quad \sqrt{\deg_T(v)^2 + \deg_T(y_1)^2} + \sqrt{\deg_T(v)^2 + \deg_T(x_1)^2} + \\ &\quad \sqrt{\deg_T(w_{k-1})^2 + \deg_T(w_k)^2} - \sqrt{\deg_{T_\tau}(v)^2 + \deg_{T_\tau}(u_1)^2} - \\ &\quad \sqrt{\deg_{T_\tau}(v)^2 + \deg_{T_\tau}(w_1)^2} - \sqrt{\deg_{T_\tau}(v)^2 + \deg_{T_\tau}(y_1)^2} - \\ &\quad \sqrt{\deg_{T_\tau}(w_{k-1})^2 + \deg_{T_\tau}(w_k)^2} - \sqrt{\deg_{T_\tau}(w_k)^2 + \deg_{T_\tau}(x_1)^2}, \\ &\geq \sqrt{4^2 + 2^2} + \sqrt{4^2 + 2^2} + \sqrt{4^2 + 2^2} + \sqrt{4^2 + 2^2} + \sqrt{2^2 + 1^2} - \\ &\quad \sqrt{3^2 + 2^2} - \sqrt{3^2 + 2^2} - \sqrt{3^2 + 2^2} - \sqrt{2^2 + 2^2} - \sqrt{2^2 + 2^2}, \\ &= 4\sqrt{20} + \sqrt{5} - 3\sqrt{13} - 2\sqrt{8} > 0. \end{aligned}$$

This completes the proof.

Using Lemma 1, we show the following main result of this section.

Theorem 2 *Let T be a tree in $\mathbb{T}_n^k$. Then,*

$$SO(T) \geq 2(n-3)\sqrt{2} + 2\sqrt{5},$$

where the equality holds if and only if $T \cong \mathbb{T}_n^{n-2} \cong P_n$.

Proof: Let $T \in \mathbb{T}_n^k$ be a tree attaining the minimum Sombor index. Assume S is a set of k cut-vertices in T . Let $B_1, \dots, B_r$ be the blocks of T corresponding to k cut-vertices. Notice that, for some ℓ , $|B_\ell| = 2$. Since T is a tree, each B_i is K_2 . First, we prove the following claim.

Claim 1 For any vertex $u \in T$ in T , we have such that $1 \leq \deg_T(u) \leq 2$.

Proof of Claim 1. On contrary, we assume that there exists a vertex $z \in T$ such that $\deg_T(z) \geq 3$. Considering $z = v$ in T_τ in Figure 1 and successively applying Lemma 1, we obtain a tree T' such that $SO(T') < SO(T)$ is satisfied. This, in turn, contradicts with the choice of T having the minimum Sombor index. ■

By Claim 1 and Corollary 1, we obtain that T has $k = n - 2$. By Remark 1, we obtain that $T \cong \mathbb{T}_n^{n-2} \cong P_n$. Routine calculations show that $SO(P_n) = 2(n - 3)\sqrt{2} + 2\sqrt{5}$. This completes the proof and shows the theorem. ■

Next, we focus on n -vertex non-tree graphs with k cut-vertices.

3 The case of non-tree graphs

Next we assume $\mathbb{H}_n^k$ to be the family of graphs on order n and k number of cut-vertices having at least one cycle. First, we present some preliminary results which will be used in our characterization.

3.1 Auxiliary results

Regarding the Sombor index of a graph, we start with an elementary lemma below.

Lemma 2 Let Γ be a graph. If $xy \in E(\Gamma)$, then $SO(\Gamma - xy) < SO(\Gamma)$.

For the Sombor index, the following lemma presents an important property of maximal 2-connected blocks.

Lemma 3 Let B be a maximal 2-connected block of a graph $\Gamma \in \mathbb{H}_n^k$, where Γ has the smallest Sombor index. If $|B| \geq 3$, then B is a cycle.

Proof: The case of $|B| = 3$ follows immediately that $B \cong C_t$ for some $t \leq n$. Next, we assume that $|B| \geq 4$. On contrary, we suppose that the block B , comprising no cut-vertices, is not isomorphic to $B \cong C_t$ for some $t \leq n$. This assumption ensures existence of an $f = xy \in E(B)$ such that there are no cut-vertices in $B - \{f\}$. This implies that $\Gamma - xy \in \mathbb{H}_n^k$. By Lemma 2, $SO(\Gamma - xy) < SO(\Gamma)$, which arises a contradiction that Γ has the smallest Sombor index. This shows the lemma. ■

Next, we present four crucial auxiliary operations playing a key role in our characterization.

Here we first explain the α -operation on a graph $\Gamma \in \mathbb{H}_n^k$. Let $\Gamma \in \mathbb{H}_n^k$ be a graph with $y, y_1, x_1, x_2 \in \Gamma$ such that $\deg_\Gamma(y) \geq 3$, $\deg_\Gamma(y_1) = 1$, and x_1x_2 is an edge lying on a cycle of Γ . See Figure 2. Let $\Gamma_\alpha = \Gamma - \{x_1x_2, y_1y\} + \{x_1y_1, x_2y_1\}$. Then, we say that Γ_α is the α -switched graph obtained from Γ by α -operation.




Figure 2: The α -operation used in Lemma 4.

Based on the α -operation, we obtain the following lemma.

Lemma 4 Let Γ_α be the α -switched graph of a graph $\Gamma \in \mathbb{H}_n^k$ described in Figure 2. Then

$$SO(\Gamma_\alpha) < SO(\Gamma).$$

Proof: Following the structure of Γ in Figure 2, assume $y, x_1, x_2 \in \Gamma$ such that $\deg_\Gamma(y) \geq 3$ and x_1x_2 is an edge lying on a cycle of Γ , where $\deg_\Gamma(x_1), \deg_\Gamma(x_2) \geq 2$. Furthermore, let Γ comprises at least one pendent vertex y_1 . Let $N_\Gamma(y) = \{y_1, y_2, \dots, y_l\}$ be the neighborhood set for y with $l \geq 3$ (see Figure 2).

Let C be an arbitrary cycle of Γ . If $y \notin C$, then the α -switched graph Γ_α is constructed by removing the edges yy_1, x_1x_2 and adding x_1y_1, x_2y_1 . By taking this into account, the following relation between Sombor indices of Γ and Γ_α emerges.

$$\begin{aligned} SO(\Gamma) - SO(\Gamma_\alpha) &= \sum_{j=1}^l \sqrt{\deg_\Gamma(y)^2 + \deg_\Gamma(y_j)^2} + \sqrt{\deg_\Gamma(x_1)^2 + \deg_\Gamma(x_2)^2} - \\ &\quad \sum_{j=2}^l \sqrt{\deg_{\Gamma_\alpha}(y)^2 + \deg_{\Gamma_\alpha}(y_j)^2} - \sqrt{\deg_{\Gamma_\alpha}(x_1)^2 + \deg_{\Gamma_\alpha}(y_1)^2} - \\ &\quad \sqrt{\deg_{\Gamma_\alpha}(x_2)^2 + \deg_{\Gamma_\alpha}(y_1)^2} \\ &= \sqrt{\deg_\Gamma(y)^2 + \deg_\Gamma(y_1)^2} + \sqrt{\deg_\Gamma(x_1)^2 + \deg_\Gamma(x_2)^2} - \\ &\quad \sqrt{\deg_{\Gamma_\alpha}(x_1)^2 + \deg_{\Gamma_\alpha}(y_1)^2} - \sqrt{\deg_{\Gamma_\alpha}(x_2)^2 + \deg_{\Gamma_\alpha}(y_1)^2} \\ &\geq \sqrt{3^2 + 1^2} + \sqrt{2^2 + 2^2} - \sqrt{2^2 + 2^2} - \sqrt{2^2 + 2^2}. \end{aligned}$$

This implies that

$$SO(\Gamma) - SO(\Gamma_\alpha) = 7\sqrt{10} - \sqrt{8} > 0.$$

Thus, we are left with the remaining case, where $y \in C$ for some cycle C of Γ . In that case, if the vertex y_1 is the unique pendent vertex of Γ , then the case is settled. Next, we assume Γ contains another pendent vertex, say z_1 , and then $\Gamma_\alpha = \Gamma - yy_1 + y_1z_1$. Thus, the conclusion is verified as well. Repeating this process would need similar arguments, thus, they have been omitted.

Therefore, the proof is finished. $\blacksquare$

Next, we provide the β -operation on a graph $\Gamma \in \mathbb{H}_n^k$. Let $\Gamma \in \mathbb{H}_n^k$ be a graph with $y, z_1, z_2 \in \Gamma$ such that $\deg_\Gamma(y) \geq 3$ and z_1z_2 is an edge lying on C , where C is a cycle of Γ . See Figure 3. Let $\Gamma_\beta = \Gamma - \{yy_2, x_{21}y_2, z_1z_2\} + \{y_2z_1, y_2z_2, x_{21}x_{l_{t_1}}\}$. Then, we say that Γ_β is the β -switched graph obtained from Γ by β -operation.




Figure 3: The α -operation used in Lemma 5.

The following lemma compares the Sombor indices of Γ and Γ_β .

Lemma 5 Let Γ_β be the β -switched graph of a graph $\Gamma \in \mathbb{H}_n^k$ described in Figure 3. Then

$$SO(\Gamma_\beta) < SO(\Gamma).$$

Proof: Following the structure of Γ in Figure 3, assume $y, z_1, z_2 \in \Gamma$ such that $\deg_\Gamma(y) \geq 3$ and $z_1 z_2$ is an edge lying on C , where C is an arbitrary cycle of Γ . Let $N_\Gamma(y) = \{y_1, y_2, \dots, y_l\}$ be the neighborhood set for y with $l \geq 3$ (see Figure 2). Moreover, if Γ comprises at least one pendent vertex with y , then the β -operation can be reduced to α -operation, and Lemma 4 is applicable. Thus, we assume that y is associated with pendent paths only with length at least 2. Give that, if y is associated to a unique pendent path, then the case is settled.

Next, assume the case in which y is associated with at least two pendent paths, say, $P_2(y_2 x_{21} \dots x_{2t_2})$ and $P_l(y_l \dots x_{l1} x_{lt_l})$, with $t_2, t_l \geq 1$. Let $\Gamma_\beta = \Gamma - \{y y_2, x_{21} y_2, z_1 z_2\} + \{y_2 z_1, y_2 z_2, x_{21} x_{lt_l}\}$. By taking this into account, the following relation between Sombor indices of Γ and Γ_β emerges.

$$\begin{aligned} SO(\Gamma) - SO(\Gamma_\beta) &= \sum_{j=1}^l \sqrt{\deg_\Gamma(y)^2 + \deg_\Gamma(y_j)^2} + \sqrt{\deg_\Gamma(z_1)^2 + \deg_\Gamma(z_2)^2} + \\ &\quad \sqrt{\deg_\Gamma(x_{21})^2 + \deg_\Gamma(y_2)^2} + \sqrt{\deg_\Gamma(x_{lt_{l-1}})^2 + \deg_\Gamma(x_{lt_l})^2} - \\ &\quad \sum_{j=1, j \neq 2}^l \sqrt{\deg_{\Gamma_\beta}(y)^2 + \deg_{\Gamma_\beta}(y_j)^2} - \sqrt{\deg_{\Gamma_\beta}(y_2)^2 + \deg_{\Gamma_\beta}(z_1)^2} - \\ &\quad \sqrt{\deg_{\Gamma_\beta}(y_2)^2 + \deg_{\Gamma_\beta}(z_2)^2} - \sqrt{\deg_{\Gamma_\beta}(x_{lt_{l-1}})^2 + \deg_{\Gamma_\beta}(x_{lt_l})^2} - \\ &\quad \sqrt{\deg_{\Gamma_\beta}(x_{21})^2 + \deg_{\Gamma_\beta}(x_{lt_l})^2} \\ &= \sqrt{\deg_\Gamma(y)^2 + \deg_\Gamma(y_2)^2} + \sqrt{8} + \sqrt{8} + \sqrt{5} - \sqrt{8} - \sqrt{8} - \\ &\quad \sqrt{8} - \sqrt{8} \\ &\geq \sqrt{13} + \sqrt{5} - 2\sqrt{8} > 0 \end{aligned}$$

This completes the proof. ■

Next, we provide the third operation which we call the γ -operation on a graph $\Gamma \in \mathbb{H}_n^k$. Following the structure of Γ in Figure 4, let Γ_0 be a graph with $|\Gamma_0| \geq 2$ and $x, z \in \Gamma_0$. Assume that Γ_1 is the graph comprising C_1 , where C_1 is a cycle. Next, from Γ_0 , we construct an n -vertex graph $\Gamma \in \mathbb{H}_n^k$ with $n \geq 6$ by identifying a vertex $c_1 \in C_1$ (resp. $c_2 \in C_2$) with x (resp. z).

We construct $\Gamma_\gamma = \Gamma - \{y_1 z, y_0 y_2, x_1 x_2\} + \{x_1 y_0, x_2 y_1\}$ and we say that Γ_γ is the γ -switched graph obtained from Γ by applying the γ -operation.




Figure 4: The γ -operation used in Lemma 6.

The following lemma compares the Sombor indices of $\Gamma \in \mathbb{H}_n^k$ and Γ_γ .

Lemma 6 Let Γ_γ be the γ -switched graph of a graph $\Gamma \in \mathbb{H}_n^k$ described in Figure 4. Then $SO(\Gamma_\gamma) < SO(\Gamma)$.

Proof: Following the structure of Γ in Figure 4, assume that $x, z \in \Gamma$ (resp. C_1, C_2) be two cut-vertices (resp. cycles) in Γ such that C_2 is an endblock. Let $x_1x_2 \in E(C_1)$ and $y_1z, y_0y_2, y_2z \in E(C_2)$ be the edges of C_1 and C_2 such that $\deg_\Gamma(x_1), \deg_\Gamma(x_2) \geq 2$. Considering $\Gamma_\gamma = \Gamma - \{y_1z, y_0y_2, x_1x_2\} + \{x_1y_0, x_2y_1\}$, we obtain the following relation between Sombor indices of Γ and Γ_γ :

$$\begin{aligned}
SO(\Gamma) - SO(\Gamma_\gamma) &= \sum_{j=1}^l \sqrt{\deg_\Gamma(z)^2 + \deg_\Gamma(y_j)^2} + \sqrt{\deg_\Gamma(y_2)^2 + \deg_\Gamma(y_0)^2} + \\
&\quad \sqrt{\deg_\Gamma(x_1)^2 + \deg_\Gamma(x_2)^2} - \sum_{j=2}^l \sqrt{\deg_{\Gamma_\gamma}(z)^2 + \deg_{\Gamma_\gamma}(y_j)^2} - \\
&\quad \sqrt{\deg_{\Gamma_\gamma}(x_1)^2 + \deg_{\Gamma_\gamma}(y_0)^2} - \sqrt{\deg_{\Gamma_\gamma}(x_2)^2 + \deg_{\Gamma_\gamma}(y_1)^2}, \\
&= \sqrt{\deg_\Gamma(z)^2 + \deg_\Gamma(y_1)^2} + \sqrt{8} + \sqrt{8} - \sqrt{8} - \sqrt{8}, \\
&\geq \sqrt{13} > 0.
\end{aligned}$$

This shows the lemma. ■

Finally, we provide the fourth operation called the δ -operation on a graph $\Gamma \in \mathbb{H}_n^k$. Following the structure of Γ in Figure 5, let Γ_0 be a graph with $|\Gamma_0| \geq 2$ and $y \in \Gamma_0$ is a vertex of Γ_0 . Assume that Γ_1 is the graph comprising C_1 , where C_1 is a cycle. Next, from Γ_0 , we construct an n -vertex graph $\Gamma \in \mathbb{H}_n^k$ by attaching a vertex $c_1 \in C_1$ and $c_2 \in C_2$ with y . This implies that C_2 is an endblock of Γ . Considering $\Gamma_\delta = \Gamma - \{yy_2, y_0y_1\} + y_0y_2$, we say that Γ_δ is constructed from Γ by applying the δ -operation.




Figure 5: The δ -operation used in Lemma 7.

The following lemma compares the Sombor indices of $\Gamma \in \mathbb{H}_n^k$ and Γ_δ .

Lemma 7 Let Γ_δ be the δ -switched graph of a graph $\Gamma \in \mathbb{H}_n^k$ as described in Figure 5. Then $SO(\Gamma_\delta) < SO(\Gamma)$.

Proof: Following the structure of Γ in Figure 5, assume that C_1, C_2 (resp. $y \in \Gamma$) be two cycles (resp. common vertex of C_1, C_2) in Γ such that $\deg_\Gamma(y) \geq 4$ and C_2 is an endblock. Assume $N_\Gamma(y) = \{y_1, y_2, \dots, y_t\}$ with $t \geq 4$ be the set denoting neighbors of $y \in \Gamma$.

Considering Γ_δ to be constructed from Γ by removing edges yy_2, y_0y_1 , and joining y_2 to y_0 , we obtain

the following relation between Sombor indices of Γ and Γ_δ :

$$\begin{aligned}
SO(\Gamma) - SO(\Gamma_\delta) &= \sum_{j=1}^t \sqrt{\deg_\Gamma(y)^2 + \deg_\Gamma(y_j)^2} + \sqrt{\deg_\Gamma(y_0)^2 + \deg_\Gamma(y_1)^2} - \\
&\quad \sum_{j=3}^t \sqrt{\deg_{\Gamma_\delta}(y)^2 + \deg_{\Gamma_\delta}(y_j)^2} - \sqrt{\deg_{\Gamma_\delta}(y)^2 + \deg_{\Gamma_\delta}(y_1)^2} - \\
&\quad \sqrt{\deg_{\Gamma_\delta}(y_0)^2 + \deg_{\Gamma_\delta}(y_2)^2}, \\
&= \sqrt{\deg_\Gamma(y)^2 + \deg_\Gamma(y_1)^2} + \sqrt{\deg_\Gamma(y)^2 + \deg_\Gamma(y_2)^2} + \\
&\quad \sqrt{8} - \sqrt{\deg_{\Gamma_\delta}(y)^2 + \deg_{\Gamma_\delta}(y_1)^2} - \sqrt{8}, \\
&\geq \sqrt{4^2 + 2^2} + \sqrt{4^2 + 2^2} - \sqrt{3^2 + 2^2} > 0.
\end{aligned}$$

This verifies the conclusion. $\blacksquare$

Next, we consider Ω to be a graph satisfying $|E(\Omega)| - |V(\Omega)| \geq 0$ and two vertices $x, y \in \Omega$ lying on a cycle of Ω . By identifying a path P_r (resp. P_s) to x (resp. y) in Ω , we construct the graph $\Omega(r, s)$.

Lemma 8 *For $x, y \in \Omega(r, s)$ such that $\deg_{\Omega(r, s)}(x), \deg_{\Omega(r, s)}(y) \geq 3$, we have*

$$SO(\Omega(r, s)) \geq SO(\Omega(1, r + s - 1)).$$

Proof: We have that $\deg_{\Omega(r, s)}(x), \deg_{\Omega(r, s)}(y) \geq 3$, as $x, y \in \Omega$ lie on a cycle of Ω . One may assume, without loss of generality, that $\deg_{\Omega(r, s)}(x) \geq \deg_{\Omega(r, s)}(y)$. Let $xx_1x_2 \dots x_{r-1}$ (resp. $yy_1y_2 \dots y_{s-1}$) be the vertices of the path P_r (resp. P_s). Assume that $N_{\Omega(r, s)}(x) = \{x_1, z_1, z_2, \dots, z_t\}$, such that $t \geq 2$. Then, we construct the graph $\Omega(1, r + s - 1)$ from $\Omega(r, s)$ by removing $xx_1 \in E(\Omega(r, s))$ and adjoining x_1 with y_{s-1} . For the sake of simplicity, let $\Omega_0 = \Omega(r, s)$ and $\Omega'_0 = \Omega(1, r + s - 1)$. Then, we deduce the following relation between the Sombor indices of Ω_0 and Ω'_0 .

$$\begin{aligned}
SO(\Omega_0) - SO(\Omega'_0) &= \sum_{j=1}^t \sqrt{\deg_{\Omega_0}(x)^2 + \deg_{\Omega_0}(z_j)^2} + \sqrt{\deg_{\Omega_0}(x_0)^2 + \deg_{\Omega_0}(x_1)^2} + \\
&\quad \sqrt{\deg_{\Omega_0}(y_{s-2})^2 + \deg_{\Omega_0}(y_{s-1})^2} - \sum_{j=1}^t \sqrt{\deg_{\Omega'_0}(x)^2 + \deg_{\Omega'_0}(z_j)^2} - \\
&\quad \sqrt{\deg_{\Omega'_0}(y_{s-2})^2 + \deg_{\Omega'_0}(y_{s-1})^2} - \sqrt{\deg_{\Omega'_0}(y_{s-1})^2 + \deg_{\Omega'_0}(x_1)^2}, \\
&\geq \sqrt{3^2 + 2^2} + \sqrt{3^2 + 2^2} + \sqrt{3^2 + 2^2} + \sqrt{2^2 + 1^2} - \\
&\quad \sqrt{3^2 + 2^2} - \sqrt{3^2 + 2^2} - \sqrt{2^2 + 1^2} - \sqrt{2^2 + 1^2}, \\
&= \sqrt{13} + \sqrt{5} - 2\sqrt{8} > 0.
\end{aligned}$$

Thus, this verifies the proof. $\blacksquare$

If we identify the two vertices x, y in $\Omega(r, s)$, then using a similar way as we did in Lemma 8, we obtain a graph $\Omega(2, r + s - 2)$ implying that $SO(\Omega(r, s)) \geq SO(\Omega(r', s'))$ such that $r' = 2, s' = r + s - 2$. Therefore, this implies that x_1 is a pendant vertex and $P_{r'} = xx_1$. In that case, we employ Lemma 4, which suggests that there exists Ω' with $| \Omega' | = | \Omega | + 1$ which is constructed from Ω by applying subdivision on edge z_1z_2 lying on some cycle and denoting this vertex as x_1 such that $SO(\Omega(r, s)) \geq SO(\Omega'(1, r + s - 2))$ is verified. This suggests the following corollary:

Corollary 2 *The identification of two vertices x, y in $\Omega(r, s)$ suggests the existence of a graph Ω' on $| \Omega | + 1$ vertices for which $SO(\Omega(r, s)) \geq SO(\Omega'(1, r + s - 2))$ is satisfied.*

3.2 Main result

This section provides a sharp lower bound on the Sombor index of graphs in $\mathbb{H}_n^k$. The corresponding graphs have been characterized as well. For $n, k \geq 2$, let C_{n-k} be a cycle of order $n - k$ and $x \in C_{n-k}$ be a vertex of C_{n-k} . Then, the family $C_{n,k}$ is obtained by adjoining a path P_{k+2} of order $k + 2$ with the vertex x of C_{n-k} . This makes $C_{n,k}$ a graph of $\mathbb{H}_n^k$ on order n and k cut-vertices.

Next, we show the main result of this paper.

Theorem 3 *Let Γ be a graph in $\mathbb{H}_n^k$. Then*

$$SO(\Gamma) \geq 2(n-4)\sqrt{2} + 3\sqrt{13} + \sqrt{5},$$

where equality holds if and only if $\Gamma \cong C_{n,k}$.

Proof: Let $\Gamma \in \mathbb{H}_n^k$ be a graph attaining the minimal Sombor index. Assume S is a set of k cut-vertices in Γ . Let $B_1, \dots, B_r$ be the blocks of Γ corresponding to k cut-vertices. Notice that, for some ℓ , either $|B_\ell| = 2$ or $|B_\ell| \geq 3$. First, we prove the following claim.

Claim 2 *There exists exactly one pendent tree P_t in Γ . Moreover, P_t is a path.*

Proof of Claim 2. Among graphs in $\mathbb{H}_n^k$, since Γ is assumed to attain minimum value of $SO(\Gamma)$. We claim that Γ comprises one or more pendent trees, as otherwise, we would have constructed Γ' from Γ by Lemmas 6 & 7 for which $SO(\Gamma') < SO(\Gamma)$ holds which then arises a contradiction to the choice of Γ . Furthermore, each of the pendant tree has to be a path, as otherwise, Lemma 5 ensures construction of Γ'' from Γ which $SO(\Gamma'') < SO(\Gamma)$, which arises a contradiction again. Thus, we obtain that there exist at least one pendent tree in Γ and, in fact, every pendent tree is a path.

Next, we show that Γ comprises precisely one pendent tree i.e. a path. One contrary, assume that there exists at least two pendent trees in Γ . Then, Lemma 8 and Corollary 2 ensures existence of graph Γ_1 such that $SO(\Gamma_1) < SO(\Gamma)$ is satisfied. This contradict with the selection of Γ .

It is important to notice that the cardinality $|B|$ of B is not effected under operations of Lemmas 5, 6, 7 & 8. This shows the claim. ■

By employing Lemma 3, we obtain that Γ is a block graph with K_2 or cycles as its blocks. Moreover, Claim 2 implies that we obtain that Γ contains P_t as its unique pendent path. If Γ comprises only one block which is a cycle, then $\Gamma \cong C_{n,k}$ and the case is settled. Thus, we assume that Γ contains two or more cycles.

Next, we show that, except for K_2 of P_t , all endblocks are cycles. As otherwise, Γ would contain at least two pendent trees which contradicts Claim 2. Next, we divide our discussion into two possible cases.

Case 1. There exists precisely two endblocks in Γ .

By using our earlier claim, one of the endblocks is a cycle C_1 and the other is K_2 . By assumption, Γ comprises another cycle, say, C_2 . Given that, Lemmas 6 & 7 ensure existence of a graph Γ' such that $SO(\Gamma') < SO(\Gamma)$ holds. We arrive at a contradiction to the choice of Γ .

Case 2. The graph Γ contains at least two endblocks.

Earlier discussion implies that Γ comprises at least two endblocks which are cycles. Once again, by Lemmas 6 & 7, there exists a graph Γ'' satisfying $SO(\Gamma'') < SO(\Gamma)$, which is a contradiction.

Both Cases 1 & 2 arise contradiction to the minimality of Γ . Thus, Γ contains exactly one cycle C_r . Since Γ lies in $\mathbb{H}_n^k$, we conclude that $C_r \cong C_{n-k}$ and $P_t \cong P_{k+2}$. This implies that $\Gamma \cong C_{n,k}$. By some

routine calculations, we arrive at $SO(C_{n,k}) = 2(n-4)\sqrt{2} + 3\sqrt{13} + \sqrt{5}$. This completes the proof and shows the theorem. ■

Let $\mathbb{G}_n^k$ be the set of all n -vertex graphs having k cut-vertices. By combining Theorem 2 & 3, show the following result.

Corollary 3 *Let G be a graph in $\mathbb{G}_n^k$. Then,*

$$SO(G) \geq 2(n-3)\sqrt{2} + 2\sqrt{5},$$

where the equality holds if and only if $G \cong P_n \cong \mathbb{T}_n^{n-2}$.

References

- [1] S. Alikhani, N. Ghanbari, Sombor index of polymers, *MATCH Commun. Math. Comput. Chem.*, 86 (2021) 715–728.
- [2] H. Chen, W. Li, J. Wang, Extremal values on the Sombor index of trees, *MATCH Commun. Math. Comput. Chem.*, 87 (2022) 23–49.
- [3] R. Cruz, I. Gutman, J. Rada, Sombor index of chemical graphs, *Appl. Math. Comput.*, 399 (2021) 126018.
- [4] R. Cruz, J. Rada, Extremal values of the Sombor index in unicyclic and bicyclic graphs, *J. Math. Chem.*, 59 (2021) 1098–1116.
- [5] K.C. Das, A.S. Cevik, I.N. Cangul, Y. Shang, On Sombor index, *Symmetry (Basel)*, 13 (2021) 140.
- [6] K.C. Das, I. Gutman, On Sombor index of trees, *Appl. Math. Comput.*, 412 (2022) 126575.
- [7] X. Fang, L. You, H. Liu, The expected values of Sombor indices in random hexagonal chains, phenylene chains and Sombor indices of some chemical graphs, 2021, arXiv: 2103.07172.
- [8] S. Filipovski, Relations between Sombor index and some degree-based topological indices, *Iranian J. Math. Chem.*, 12 (2021) 19–26.
- [9] I. Gutman, Geometric approach to degree-based topological indices: Sombor indices, *MATCH Commun. Math. Comput. Chem.*, 86 (2021) 11–16.
- [10] Gutman, I. On Sombor index of trees, *Appl. Math. Comput.*, in press, (2022).
- [11] I. Gutman, Some basic properties of Sombor indices, *Open J. Discrete Appl. Math.*, 4 (2021) 1–3.
- [12] I. Gutman, B. Furtula, *Novel Molecular Structure Descriptors – Theory and Applications I*, Univ. Kragujevac, Kragujevac, 2010.
- [13] B. Horoldagva, C. Xu, On Sombor index of graphs, *MATCH Commun. Math. Comput. Chem.*, 86 (2021) 703–713.
- [14] H. Hua, S. Zhang, Graphs with given number of cut vertices and extremal Merrifield-Simmons index, *Discrete Appl. Math.*, 159(10) (2011) 971–980.

- [15] S. Ji, S. Wang, On the sharp lower bounds of Zagreb indices of graphs with given number of cut vertices, *J. Math. Anal. Appl.*, 458(1) (2018) 21–29.
- [16] S. Li, Z. Wang, M. Zhang, On the extremal Sombor index of trees with a given diameter, *Appl. Math. Comput.*, 416 (2022) 126731.
- [17] Z. Lin, On the spectral radius, energy and Estrada index of the Sombor matrix of graphs, 2021, arXiv: 2102.03960.
- [18] H. Liu, Ordering chemical graphs by their Sombor indices, 2021a, arXiv: 2103.05995.
- [19] H. Liu, Maximum Sombor index among cacti, 2021b, arXiv: 2103.07924.
- [20] H. Liu, L. You, Y. Huang, Ordering chemical graphs by Sombor indices and its applications, *MATCH Commun. Math. Comput. Chem.*, 87 (2022).
- [21] H. Liu, L. You, Z. Tang, J.B. Liu, On the reduced Sombor index and its applications, *MATCH Commun. Math. Comput. Chem.*, 86 (2021) 729–753.
- [22] I. Milovanović, E. Milovanović, A. Ali, M. Matejić, Some results on the Sombor indices of graphs, *Contrib. Math.*, 3 (2021) 59–67.
- [23] I. Milovanović, E. Milovanović, M. Matejić, On some mathematical properties of Sombor indices, *Bull. Int. Math. Virtual Inst.*, 11 (2021) 341–353.
- [24] W. Ning, Y. Song, K. Wang, More on Sombor index of graphs, *Mathematics (Basel)*, 10(3) (2022) 301.
- [25] I. Redžepović, Chemical applicability of Sombor indices, *Serb. Chem. Soc.*, 86 (2021) 445–457.
- [26] T. Réti, T. Došlić, A. Ali, On the Sombor index of graphs, *Contrib. Math.*, 3 (2021) 11–18.
- [27] Y. Shang, Sombor index and degree-related properties of simplicial networks, *Appl. Math. Comput.*, 419 (2022) 126881.
- [28] X. Sun, J. Du, On Sombor index of trees with fixed domination number, *Appl. Math. Comput.*, 421 (2022) 126946.
- [29] R. Todeschini, V. Consonni, *Handbook of Molecular Descriptors*, Wiley-VCH, Weinheim, 2000.
- [30] Z. Wang, Y. Mao, Y. Li, B. Furtula, On relations between Sombor and other degree-based indices, *J. Appl. Math. Comput.*, 68(1) (2022) 1–17.
- [31] D.B. West, *Introduction to Graph Theory*, Prentice Hall, Upper Saddle River, (2001).
- [32] K. Xu, H. Liu, Y. Yang, K.C. Das, The minimal Kirchhoff index of graphs with a given number of cut vertices, *Filomat*, 30(13) (2016) 3451–3463.
- [33] T. Zhou, Z. Lin, L. Miao, The Sombor index of trees and unicyclic graphs with given matching number, 2021a, arXiv: 2103.04645.
- [34] T. Zhou, Z. Lin, L. Miao, The Sombor index of trees and unicyclic graphs with given maximum degree, 2021b, arXiv: 2103.07947.