arXiv:2305.19969v1 [math.NT] 31 May 2023

On the p-isogenies of elliptic curves with multiplicative reduction over quadratic fields

George C. Ţurcaş

Abstract Let $q > 5$ be a prime and K a quadratic number field. In this article we extend a previous result of Najman and the author and prove that if E/K is an elliptic curve with potentially multiplicative reduction at all primes $\mathfrak{q} \mid q$, then E does not have prime isogenies of degree greater than 71 and different from q. As an application to our main result, we present a new variant of the asymptotic version of Fermat's Last Theorem over quadratic imaginary fields of class number one.

1 Introduction

An isogeny $\varphi : E \to E'$ between elliptic curves is a non-constant morphism that preserves the group structure. In this note, we will consider elliptic curves E and E' defined over a number field number field K. If, in addition, $\varphi^\tau = \varphi$ for any $\tau \in G_K = \mathrm{Gal}(\overline{K}/K)$, we say that φ is a K rational isogeny. In particular, this means that φ can be represented by a rational map whose coefficients lie in K. Suppose that p is a rational prime.

In this case, the degree of an isogeny is equal to the cardinality of its kernel. If p is a rational prime and $\varphi : E \to E'$ is of degree p and defined over K, we will say that E admits a rational p-isogeny. As we will see in Section 3, the condition that E has (or does not have) a p-isogeny can be formulated in the language of residual Galois representations.

Mazur [12, Theorem 1] proved that for $p > 163$, the elliptic curves E defined over $\mathbb{Q}$ do not have p-isogenies. For a general number field K, a direct analogue to Mazur's theorem is not true. Indeed, suppose E is an elliptic curve that has complex multiplication by an order $\mathcal{O}$ in some imaginary quadratic field and that p splits in $\mathcal{O}$. If K contains the Hilbert class field of $\mathcal{O}$, then E admits an endomorphism of

George C. Ţurcaş
Babeş-Bolyai University, Cluj-Napoca, Romania and "Simion Stoilow" Institute of Mathematics of the Romanian Academy, Bucharest, Romania e-mail: george.turcas@ubbcluj.ro

degree p defined over K. In such a situation, E admits isogenies of arbitrary large degree.

Momose [18] showed that for any number field K, there exists a constant $C(K)$, such that if E is an elliptic curve defined over K admitting a K-rational p-isogeny, then the isogeny falls into three types according to its isogeny character (see Section 3). We refer to the aforementioned work of Momose for a precise description of types I, II, and III of isogeny characters. We mention that for large p, isogenies of type I are often ruled out using formal immersion criteria such as the result due to Kamienny [9, Proposition 3.2], which we employ in Section 2. Momose also studied conditions that can, in some situations, rule out isogenies of type II. More recently, these were generalised by Banwait and Derickx [2]. In *loc. cit.* the authors produced an algorithm that can, in theory, be used to find an upper bound for the possible degrees of p-isogenies of Type I. Moreover, assuming the Generalised Riemann Hypothesis, the authors prove an explicit upper bound for the possible degrees of p-isogenies of type II. About the type III p-isogenies, Momose [18] writes that he expects them to appear only in the case of elliptic curves with complex multiplication.

Finally, we note that in recent work Banwait [1] described an algorithm which finds the (known to be finitely many) primes p for which there exists an elliptic curve over K, admitting a K-rational p-isogeny, when K is a quadratic field which is not imaginary of class number one. As we shall see, Theorem 2, which is the main result in this paper, asserts the existence of a bound for the degree of p-isogenies for infinite families of elliptic curves defined over quadratic fields K, including the imaginary quadratic fields of class number one.

One of the key results that Mazur uses to classify prime isogenies in [12] and then, in a different article [11], prime torsion is the following.

Theorem 1 (Corollary 4.4 in [12]) *Let $E/\mathbb{Q}$ be an elliptic curve and let $p > 19$ be a prime for which E admits a rational p-isogeny. Then E has potentially good reduction at any prime $q \notin \{2, p\}$.*

We are now ready to present our main result.

Theorem 2 *Let K be a quadratic field and let $q > 5$ be a rational prime that is unramified in K. Suppose that an elliptic curve E/K has potentially multiplicative reduction at all primes $\mathfrak{q}$ of K above q. For $p > 71$, the elliptic curve E does not have p-isogenies if the pair (q, p) satisfies one of the following two conditions:*

- *q is inert in K;*
- *q splits in K and $q \neq p$.*

We point out that the result above extends [20, Theorem 1.2], which covers the case when q is inert in K. In the present work, we focus on the case where q splits in K. Applying a formal *immersion criterion* (see Section 2) in this case was more difficult due to the possibility that a non-cuspidal point in $y \in X_0(p)(K)$ reduces to different cusps modulo the two distinct primes above q. Dealing with this possibility constitutes the main contribution of the present work and is what makes it complementary to [20, Theorem 1.2] and to [16, Proposition 4.11]. We would like

to point out that the proof of our theorem makes use, in an essential way, of a result due to Michaud-Jacobs [14, Theorem 4.6].

We note that for our application to Asymptotic Fermat (Theorem 3), one really uses the fact that the lower bound 71 for p does not depend on q, as one applies the main theorem to a (Frey) elliptic curve for which the prime q depends on unknown solutions to some Diophantine equation. In our case, we will show that q and hence also the solutions to the Diophantine equations do not exist.

Given an elliptic curve E over a number field K, which has a certain type of reduction at a fixed prime $\mathfrak{q}$, it is sometimes possible to find a bound $B_{K,\mathfrak{q}}$ such that for $p > B_{K,\mathfrak{q}}$ the elliptic curve E does not have p-isogenies. An example of such a result is [21, Proposition 6.1], which asserts the existence of $B_{K,\mathfrak{q}}$ if K is Galois, E has potentially multiplicative reduction at $\mathfrak{q}$ and E is semistable at all primes $\mathfrak{p} \mid p$. In section 3 we give a brief account of some recent results of this type.

Recall that an elliptic curve E/K has potentially good reduction at a prime ideal $\mathfrak{q}$ of K if $v_{\mathfrak{q}}(j(E)) \geq 0$, where $v_{\mathfrak{q}}$ is the $\mathfrak{q}$-adic valuation. If $v_{\mathfrak{q}}(j(E)) < 0$, then E has potentially multiplicative reduction at $\mathfrak{q}$. We will see that, when used for (Frey) elliptic curves within the modular method, our main theorem can give some information about the primes $\mathfrak{q}$ that divide $j(E)$. As the j-invariants of these (Frey) elliptic curves depend on the solutions to some Diophantine equation, this information can be used to show that the latter solutions do not exist.

In support of our our last point above, we show that Theorem 2 can be used to prove the following version of Asymptotic Fermat's Last Theorem for half of the prime exponents p.

Theorem 3 *Let K be a quadratic imaginary field of class number one and suppose that Conjecture 1 holds for K. Fix S, a finite set of rational primes containing 2, 3 and 5. There exists a constant C_S, depending only on S, such that for all $p > C_S$, if the following hold:*

1. $p \equiv 1 \pmod 3$*;*
2. $a, b, c \in \mathcal{O}_K \setminus \{0\}$ *are coprime such that* $a^p + b^p + c^p = 0$*;*
3. *If $q \notin S$ is a rational prime that divides* $\mathrm{Norm}(abc)$*, then all primes above q divide abc;*
4. *If p splits in K, then at most one of the primes above p divides abc in $\mathcal{O}_K$.*

then $K = \mathbb{Q}(\sqrt{-3})$ and the triple (a, b, c) is, up to reordering, $(1, \epsilon, \epsilon^2)$, where ϵ is a non-trivial third root of unity.

The reader familiar with Theorem 1.5 in [20] might note a strong similarity between that result and the theorem stated above. We point out that the third condition above is more relaxed than the corresponding one in Theorem 1.5, allowing for more possible primes in the support of abc.

This result is aligned with a more general Fermat Conjecture. We refer the interested reader to the discussion on page 2 of [6], where the authors note that the Fermat Conjecture follows from the abc-conjecture for number fields. In the same direction, assuming a different conjecture about the surjectivity of mod p Galois representations of non-CM elliptic curves, in previous work the author proved a

version of the Fermat Conjecture over quadratic imaginary fields of class number one (see Conjecture 5.2 and Theorem 5.3 in [25]).

Acknowledgements The author is indebted to Mugurel Barcău and Vicenţiu Paşol for helpful suggestions regarding quotients of arithmetic surfaces and the proof of Propostion 1. This work is supported by a grant from the Romanian Ministry of Education and Research, CNCS-UEFISCDI, project number PN-III-P4-ID-PCE-2020-2498.

2 Formal immersions and the proof of Theorem 2

Throughout, we will denote by p and q rational primes and by $\mathfrak{p}$ and $\mathfrak{q}$, respectively, prime ideals in O_K, the ring of integers of K. We write $X_0(p)$ for the classical modular curve of level p, whose non-cuspidal K-rational points represent isomorphism classes of pairs (E, C) of elliptic curves E together with a K-rational subgroup C of order p. The curve $X_0(p)$ has two cusps, both defined over $\mathbb{Q}$. We follow the convention in [4], where ∞ is the cusp that is unramified under the j-map $X_0(p) \to X_0(1) \simeq \mathbb{P}^1$. The j-map is ramified of degree p at the cusp 0. Under this convention, 0 represents a a Néron p-gon whereas ∞ corresponds to a Néron 1 -gon.

The symmetric square $X_0^{(2)}(p)$

Let S_2 be the symmetric group with two elements. This group acts on $X_0(p) \times X_0(p)$ by permuting the factors. We write $X_0^{(2)}(p)$ for the quotient of $X_0(p) \times X_0(p)$ under this action. This is called the symmetric square of $X_0(p)$ and we will denote its points by (unordered) pairs (y, z).

We write τ for the generator of $\mathrm{Gal}(K/\mathbb{Q})$, which can also be viewed as an element of $\mathrm{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$ by choosing an automorphism of $\overline{\mathbb{Q}}$ that extends τ. In this case, a quadratic point $x \in X_0(p)(K)$ gives rise to a rational point $(x, x^\tau) \in X_0^{(2)}(p)(\mathbb{Q})$. As $X_0(p)$ can be seen as a smooth scheme over $\operatorname{Spec} \mathbb{Z}[1/p]$, it follows that $X_0^{(2)}(p)$ can also be seen as a smooth arithmetic scheme over $\operatorname{Spec} \mathbb{Z}[1/p]$. This is true because one can parameterise the local rings of $X_0^{(2)}(p)$ by elementary symmetry functions, as is explained, for instance, in the proof of [17, Proposition 3.2].

The surface $X_0^w(p)$

It is known that the Fricke involution w_p acts on $X_0(p)$. We define a new action of the group S_2 on the product $X_0(p) \times X_0(p)$, as follows. The non-trivial element of S_2 sends a pair of points (y, z) to $(w_p(z), w_p(y))$.

The existence of this quotient of $X_0(p) \times X_0(p)$ as a scheme over $\operatorname{Spec}\mathbb{Z}\left[\frac{1}{p}\right]$ follows from [SGA I, Exposé V, Proposition 1.8]. We shall denote it by $X_0^w(p)$. Our next result shows that this scheme $X_0^w(p)$ is smooth.

Proposition 1 *The arithmetic surface $X_0^w(p)$ is smooth over* $\operatorname{Spec}\mathbb{Z}\left[\frac{1}{p}\right]$.

Proof We follow the classical arguments used in the proof of the smoothness of the symmetric power $X_0^{(2)}(p)$.

Let $\pi : X_0(p)\times X_0(p) \to X_0^w(p)$ be the projection map. The fact that the structure morphism $X_0^w(p) \to \operatorname{Spec}\mathbb{Z}[1/p]$ is flat follows immediately from [8, Proposition 9.7]. Following [19, Theorem 3' on pp. 221], it is enough to show that the fiber over any geometric point $\operatorname{Spec}\Omega \to \operatorname{Spec}\mathbb{Z}[1/p]$ of this structure morphism is regular (where Ω is an algebraically closed field).

If $Q \in X_0^w(p)$ is a point such that $\pi^{-1}(Q)$ consists of two distinct points P_1, P_2, then π is étale at P_1 and induces an isomorphism $\hat{O}_Q \to \hat{O}_{P_1}$ of complete local rings. Since P_1 is a regular point, it follows that Q is a regular point of $X_0^w(p)$.

If $Q \in X_0^w(p)$ is such that $\pi^{-1}(Q)$ consists of a single point $P = (y, w_p(y))$ (a fixed point of the S_2 action), then notice the following. The completed local ring $\hat{O}_P \simeq \Omega[[t, s]]$, where t is a local parameter at y and $s = t \circ w_p$ is a local parameter at $w_p(y)$. Notice that the induced action of S_2 on the pair (t, s) is trivial, therefore we have the ring isomorphisms $\hat{O}_Q \simeq \Omega[[t, s]]^{S_2} = \Omega[[t, s]] \simeq \hat{O}_P$, which imply that Q is a regular point of $X_0^w(p)$. □

Moreover, we have the following commutative diagram


$$\begin{array}{ccc} (y,z) & \longmapsto & (w_p(y),z) \\ & & \\ X_0(p)\times X_0(p) & \xrightarrow{\sim} & X_0(p)\times X_0(p) \\ \downarrow \text{proj} & & \downarrow \pi \\ X_0^{(2)}(p) & \longrightarrow & X_0^w(p) \\ & & \\ (y,z) & \longmapsto & (w_p(z),y) \end{array} \tag{1}$$

Since the top horizontal map is an isomorphism, proj and π have the same degree, we can deduce that the quotient surfaces $X_0^w(p)$ and $X_0^{(2)}(p)$ are in fact isomorphic over $\operatorname{Spec}\mathbb{Z}[1/p]$.

We write $J_0(p)$ for the Jacobian of $X_0(p)$ and let $h : X_0^{(2)}(p)(\mathbb{Q}) \to J_0(p)(\mathbb{Q})$ be such that $h(y, z) = [y + z - 2\infty]$. This map is injective if the genus $X_0(p)$ is greater than one and if $X_0(p)$ is not hyperelliptic. Thus, h is injective if $p > 71$ (see the discussion preceding Proposition 3. 2 in [9]).

Let J_p be the Eisenstein quotient of $J_0(p)$ (see [11] for the definition). Mazur [10, Theorem 4] proved that the rank of $J_p(\mathbb{Q})$ is zero.

We point out the existence of the following commutative diagram over $\operatorname{Spec}\mathbb{Z}\left[\frac{1}{p}\right]$.



(2)

The top left isomorphism above is given by $(y, z) \mapsto (w_p(z), y)$ and $h_w : X_0^w(p) \to J_0(p)$ is defined as

$$h_w(y, z) = [y + w_p(z) - 2\infty].$$

Note that the injectivity of h_w follows from that of h and the commutativity of the diagram.

Write $f : X_0^{(2)}(p) \to J_p$ for the composition between h and the natural projection from $J_0(p)$ to J_p on the diagram above. A result of Kamienny [9, Proposition 3.2] states that for $q > 5$ and $p > 71$, the map f is a formal immersion at $(\infty, \infty)_{\mathbb{F}_\mathfrak{q}}$.

Note that the isomorphism $X_0^w(p) \to X_0^{(2)}(p)$ sending $(y, z) \mapsto (w_p(z), y)$ maps $(\infty, 0) \in X_0^w(p)$ to $(\infty, \infty) \in X_0^{(2)}(p)$. Now, if we similarly denote by $f_w : X_0^w(p) \to J_p$ the composition between h_w and the natural projection proj on the diagram (2), the proof of the following result follows directly.

Proposition 2 *For $q > 5$ and $p > 71$, the map*

$$f_w : X_0^w(p)_{/\operatorname{Spec}\mathbb{Z}[1/p]} \to J_{p/\operatorname{Spec}\mathbb{Z}[1/p]}$$

is a formal immersion at $(\infty, 0)_{\mathbb{F}_q}$.

We are now ready to present the proof of the main result.

Proof (of Theorem 2) To prove our claims, we use the following standard argument. Suppose that there is an elliptic curve E, defined over K, that satisfies the hypotheses of the theorem. Suppose that the pair (q, p) satisfies either of the two conditions stated in the theorem and that E admits a p-isogeny for $p > 71$. The elliptc curve E together with the kernel of the isogeny gives rise to a non-cuspidal point $x \in X_0(p)(K)$. As explained in the beginning of the section, this gives rise to a rational point $y = (x, x^\tau) \in X_0^{(2)}(p)(\mathbb{Q})$, where $\tau \in \operatorname{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$ is an extension to $\overline{\mathbb{Q}}$ of the generator of $\operatorname{Gal}(K/\mathbb{Q})$. The fact that E has potentially multiplicative reduction at all primes $\mathfrak{q} \mid q$ implies that x reduces modulo every such $\mathfrak{q}$ to one of the two cusps of $X_0(p)(\mathbb{F}_\mathfrak{q})$. We therefore deduce that $y = (x, x^\tau) \in X_0^{(2)}(p)(\mathbb{Q})$ reduces modulo q to

$$(\infty, \infty)_{\mathbb{F}_q}, (0, 0)_{\mathbb{F}_q} \text{ or } (\infty, 0)_{\mathbb{F}_q}.$$

If $(x, x^\tau)_{\mathbb{F}_q} = (\infty, \infty)_{\mathbb{F}_q}$ or $(x, x^\tau)_{\mathbb{F}_q} = (0, 0)_{\mathbb{F}_q}$, the theorem follows from a result of Michaud-Jacobs [16, Proposition 4.11]. This always happens when q is inert in O_K, a case that was already settled in previous work by Najman and the author [20, Theorem 1.2].

Suppose now that the prime q splits in K and that $y = (x, x^\tau) \in X_0^{(2)}(\mathbb{Q})$ reduces to $(\infty, 0)_{\mathbb{F}_q}$ modulo q. In this situation, using Siksek's *relative symmetric Chabauty criterion*, Michaud-Jacobs [14, Theorem 4.6] proved that the action of the Fricke involution w_p on $x \in X_0(p)(K)$ coincides with the action of τ on the same point. To be precise, from the aforementioned result it follows that $w_p(x) = x^\tau$ on $X_0(p)$ and hence $y = (x, x^\tau)$ can also be regarded as a rational point (i.e. stable under $\mathrm{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$) on $X_0^w(p)$. Indeed, the action on τ on the quotient $(x, x^\tau) \in X_0^w(p)(K)$ can be written as

$$(x, x^\tau)^\tau = (x^\tau, x) = (w_p(x), x) = (x, w_p(x)) = (x, x^\tau),$$

where the third equality follows from the mere definition of $X_0^w(p)$. We would like to emphasise that the restriction $q \neq p$ is imposed by the hypotheses of Theorem 4.6 in [14] and the application of the latter result is the only place where we used this restriction.

Proposition 2 above asserts that f_w is a formal immersion at $(\infty, 0)_{\mathbb{F}_q}$. An important consequence of this fact is that there exists at most one point on $X_0^w(p)(\mathbb{Q})$ reducing to $(\infty, 0)_{\mathbb{F}_q}$ (see, for instance, [15, Lemma 3.1]). Since we know that $(\infty, 0) \in X_0^w(p)(\mathbb{Q})$ reduces to $(\infty, 0)_{\mathbb{F}_q}$, it follows that $(x, x^\tau) = (\infty, 0) \in X_0^w(p)(\mathbb{Q})$. This contradicts the hypothesis that $x \in X_0(p)(K)$ is non-cuspidal and hence finishes our proof. □

? Questions

1. Can one adapt the techniques above and prove a statement analogous to Theorem 2 for a general number field K of degree $d > 2$? Assuming that E has multiplicative reduction at all primes $\mathfrak{q} \mid q$, one has to be able to handle cases in which the corresponding point on $X_0^{(d)}(p)(\mathbb{Q})$ reduces to $(\infty, \infty, 0)$. Proving a result similar to [5, Lemma 1.7] for $X_0^{(d)}(p)$ might help.
2. We expect that the conclusion of Theorem 2 holds if E has potentially multiplicative reduction at one prime $\mathfrak{q} \mid q$. Furthermore, we believe that the restriction $q \neq p$ is not necessary. Can any of these conditions be relaxed using similar techniques?

3 Elliptic curves with bad reduction at a known fixed prime $\mathfrak{q}$

In this section, we will restrict our attention to elliptic curves E over K that are semistable at all primes $\mathfrak{p} \mid p$. This last condition is satisfied by the (Frey) elliptic curves used in the modular method, where p is an unknown exponent appearing

in some Diophantine equation. Throughout, we assume that $p \geq 17$ and that p is unramified in K.

Suppose that E admits a p-isogeny φ defined over K. Then, its kernel is a K-rational cyclic subgroup of $E[p]$ and, hence, the mod p Galois representation $\overline{\rho}_{E,p} : G_K \to \mathrm{Aut}(E[p]) \cong \mathrm{GL}_2(\mathbb{F}_p)$ of E is reducible and

$$\overline{\rho}_{E,p} \sim \begin{pmatrix} \theta & * \\ 0 & \theta' \end{pmatrix},$$

where $\theta, \theta' : G_K \to \mathbb{F}_p^\times$ are characters whose product $\theta\theta'$ is equal to χ_p, the mod p cyclotomic character. The character θ is called the *isogeny character* of $(E, \ker\varphi)$ and dictates the action of G_K on $\ker\varphi$. To be precise, if $P \in \ker\varphi$, then

$$P^\sigma = \theta(\sigma)P, \text{ for all } \sigma \in G_K.$$

Let us now fix $\mathfrak{q}$, a prime ideal of K.

Proposition 6.1 in [21] asserts that for E as above, if one knows that $\mathfrak{q}$ is a prime of potentially multiplicative reduction, then there exists a constant $B_{K,\mathfrak{q}}$ such that if E has a p-isogeny, then $p \leq B_{K,\mathfrak{q}}$. This result is proved using a careful study of the isogeny character θ. The authors show that the 12-th powers of θ or θ' are everywhere unramified, a fact that implies that E or an elliptic curve that is isogenous to E admits a p-torsion point defined over a fixed degree extension of K. The upper bound on p is then obtained applying Merel's uniform boundedness theorem for torsion points [13].

Now assume that $\mathfrak{q}$ is a prime of potentially good reduction for E. It is known that the character θ^{12} is unramified at the infinite places of K and at all finite places of K that are coprime to p. Moreover, if $\mathfrak{p} \mid p$, restricting to $I_\mathfrak{p} < G_K$ the inertia subgroup at $\mathfrak{p}$, we have that $\theta^{12}|_{I_\mathfrak{p}} = \chi_p^{s_\mathfrak{p}}|_{I_\mathfrak{p}}$, where $s_\mathfrak{p} \in \{0, 12\}$. Recall that τ is the generator of $\mathrm{Gal}(K/\mathbb{Q})$, so $\tau(\mathfrak{p})$ is also a prime ideal above p. The tuple $(s_\mathfrak{p}, s_{\tau(\mathfrak{p})})$ is called the *isogeny signature* of the character θ and is equal to one of

$$(0,0), (12,12), (12,0) \text{ or } (0,12).$$

If we add the assumption that p is inert in K, then the signatures $(12,0)$ and $(0,12)$ are not possible. In this case, the isogeny signature is $(0,0)$ or $(12,12)$. It follows from Lemma 34 in [16] that, replacing E with a p-isogenous elliptic curve if necessary, one can assume that the isogeny signature is $(0,0)$. The existence of a bound $B_{K,\mathfrak{q}}$ such that $p < B_{K,\mathfrak{q}}$ is guaranteed by [16, Corollary 4.7].

The discussion above can be summarised in the following proposition.

Proposition 3 *Let E be an elliptic curve which has bad reduction at some fixed prime $\mathfrak{q}$ and is semistable at all primes $\mathfrak{p}$ above some rational prime p. Suppose the pair $(\mathfrak{q}, p)$ satisfies one of the following conditions*

- *$\mathfrak{q}$ is of potentially multiplicative reduction;*
- *$\mathfrak{q}$ is of potentially good reduction and p is inert in K.*

Then there exists a bound $B_{K,\mathfrak{q}}$ such that for $p > B_{K,\mathfrak{q}}$, the elliptic curve E does not have p-isogenies.

In the next section, we shall present the proof of the following Diophantine application of this result.

Theorem 4 *Let K be a quadratic imaginary field of class number one and suppose that Conjecture 1 holds for K. Then, there is an effective constant B_K such that for every prime $p > B_K$, $p \equiv 1 \pmod 3$ and p inert in K the only solutions $a, b, c \in K \setminus \{0\}$ to the equation*

$$a^p + b^p + c^p = 0 \tag{3}$$

are, up to reordering, $(1, \epsilon, \epsilon^2)$, where ϵ is a non-trivial third root of unity.

4 The Asymptotic Fermat's Last Theorem over some quadratic imaginary fields

Most of the literature discussing the resolution of certain Diophantine equations using the modular method is restricted to the case where K is totally real. This is mostly due to the progress in proving modularity lifting theorems over such fields. On the other hand, modularity results over number fields with complex embeddings are mostly conjectural, with a few notable exceptions such as those presented in the recent extraordinary work of Caraiani and Newton [3].

In this section, we will restrict to the case in which K is a quadratic imaginary field of class number one and $p \geq 5$ a rational prime that does not ramify in K. For notation, relevant definitions, and a detailed discussion concerning complex and mod p eigenforms for GL_2 over K, we refer the reader to [24, Section 2].

The results concerning the Fermat equation presented in this work are conjectural upon the veracity of the following version of Serre's modularity conjecture [25, Conjecture 2.2].

Conjecture 1 Let $p > 3$ be a prime and E an elliptic curve defined over K. Suppose that the mod p Galois representation $\overline{\rho}_E : G_K \to \mathrm{GL}_2(\overline{\mathbb{F}}_p)$, induced by the action of G_K on the p-torsion of E, is absolutely irreducible, continuous with Serre conductor $\mathcal{N}$ (prime-to-p part of its Artin conductor). Assume that $\overline{\rho}_E|_{G_{K_\mathfrak{p}}}$ arises from a finite-flat group scheme over $\mathcal{O}_{K_\mathfrak{p}}$ for every prime $\mathfrak{p}|p$. Then there is a mod p eigenform $c \in H^1(Y_0(\mathcal{N}), \overline{\mathbb{F}}_p)$ such that for all prime ideals $(\pi) \subseteq \mathcal{O}_K$, coprime to $p\mathcal{N}$

$$T_\pi(c) = \mathrm{Trace}(\overline{\rho}_E(\mathrm{Frob}_{(\pi)}))$$

Suppose that $a, b, c \in K \setminus \{0\}$ are such that $a^p + b^p + c^p = 0$. Let

$$E = E_{a,b,c} : Y^2 = X(X - a^p)(X + b^p). \tag{4}$$

As we are working in quadratic imaginary fields of class number one, one can assume that $a, b, c \in \mathcal{O}_K$, where $\mathcal{O}_K$ is the ring of integers of K, and that their are coprime.

From the proof of Theorem 1.3 in [25] it follows easily that if K is quadratic imaginary of class number one and $p \geq 19$, assuming Conjecture 1 holds for K, there are no solutions to the Fermat equation $a^p + b^p + c^p = 0$ for which $\overline{\rho}_{E,p}$ is absolutely irreducible. Furthermore, the author shows that if $2 \mid \mathrm{Norm}_{K/\mathbb{Q}}(abc)$, then for $p \geq 19$, the representation $\overline{\rho}_{E,p}$ is absolutely irreducible. The latter implies that for such K and p, the Fermat equation

$$a^p + b^p + c^p = 0,$$

does not have solutions in coprime $a, b, c \in \mathcal{O}_K \setminus \{0\}$ such that $2 \mid abc$.

We start proving Theorems 3 and 4 in parallel, following the same strategy as in [20, Section 3]. Recall that the quadratic imaginary fields of class number one are $K = \mathbb{Q}(\sqrt{-d})$, where d is one of 1, 2, 3, 7, 11, 19, 43, 67 or 163. The cases $d \in \{1, 2, 7\}$ follow by more general results in [24]. From now on we will assume that K is one of the six remaining fields. In each of the latter quadratic fields, the prime 2 is inert. From the proof of [25, Theorem 1.3] it is easy to deduce that the following hold simultaneously:

- The prime ideal $2\mathcal{O}_K$ does not divide abc.
- If $E : Y^2 = X(X - a^p)(X + b^p)$ is the Frey elliptic curve attached to this solution, then E has additive potentially good reduction at the prime $2\mathcal{O}_K$ and E is semistable at every prime ideal $\mathfrak{q} \neq 2\mathcal{O}_K$.
- The mod p Galois representation $\overline{\rho}_{E,p}$ is absolutely reducible and unramified outside the primes above $2p$.

If we assume $p \geq 53$, it follows from [23, Lemme 18'] that the image of $\overline{\rho}_{E,p}(G_K)$ under the projection $\mathrm{GL}_2(\mathbb{F}_p) \to \mathrm{PGL}_2(\mathbb{F}_p)$ contains an element whose order is at least 13. We point out that the validity of the proof for Lemme 18' in loc. cit. extends beyond the case $K = \mathbb{Q}$, as long as p does not ramify in K. The existence of such an element ensures that the image of $\overline{\rho}_{E,p}(G_K)$ in $\mathrm{PGL}_2(\mathbb{F}_p)$ is not contained in any of its exceptional subgroups A_4, S_4 or A_5 (see [22, Subsection 2.5] for more details). From the discussion in [22, Section2] we deduce that if $\overline{\rho}_{E,p} \otimes_{\mathbb{F}_p} \overline{\mathbb{F}}_p$ is diagonalisable, then the image $\overline{\rho}_{E,p}(G_K)$ is contained in a Cartan subgroup of $\mathrm{GL}_2(\mathbb{F}_p)$.

Assume additionally that $p \equiv 1 \pmod 3$, as it is specified in the hypotheses of both theorems 3 and 4.

We can distinguish the following two cases.

Suppose first that $\overline{\rho}_{E,p}$ is irreducible but absolutely reducible. In this situation, it follows that the image $\overline{\rho}_{E,p}(G_K)$ is contained in a Cartan non-split subgroup. We have that $\overline{\rho}_{E,p} \otimes_{\mathbb{F}_p} \overline{\mathbb{F}}_p \sim \begin{pmatrix} \theta & 0 \\ 0 & \theta^p \end{pmatrix}$, where $\theta : G_K \to \mathbb{F}_{p^2}^{\times}$ is a character. The latter is not $\mathbb{F}_p$-valued and $\theta^{p+1} = \chi_p$, where $\chi_p : G_K \to \mathbb{F}_p^{\times}$ is the mod p cyclotomic character.

As $2\mathcal{O}_K$ is a prime of potentially good reduction for E, one knows that $v_{2\mathcal{O}_K}(j(E)) \geq 0$. Using [7, Lemma 3.7], we deduce that 3 divides $\#\overline{\rho}_{E,p}(I_{2\mathcal{O}_K})$, where $I_{2\mathcal{O}_K} \subseteq G_K$ is the inertia subgroup at $2\mathcal{O}_K$. By Cauchy's Theorem, there exists an element $g \in I_{2\mathcal{O}_K}$ such that $\overline{\rho}_{E,p}(g)$ and $\theta(g)$ have both order 3 in $\mathrm{GL}_2(\mathbb{F}_{p^2})$. As $p > 2$, we know that χ_p is unramified at places above 2. It follows that $\chi_p(g) = \theta^{p+1}(g) = 1$, and hence $3 \mid p+1$, a contradiction to the choice $p \equiv 1 \pmod 3$.

We are now left with the case that $\overline{\rho}_{E,p}$ is reducible, meaning that

$$\overline{\rho}_{E,p} \sim \begin{pmatrix} \theta & * \\ 0 & \theta' \end{pmatrix},$$

where $\theta, \theta' : G_K \to \mathbb{F}_p^{\times}$ are the isogeny characters. Now we discuss the proofs of theorems 3 and 4 separately.

4.1 Proof of Theorem 3 as an application to Theorem 2

Suppose that a, b, c are as above. Since

$$j(E) = \frac{2^4(b^{2p} - a^p c^p)}{(abc)^{2p}},$$

we see that if $\mathfrak{q}$ is a prime that divides abc, then $v_{\mathfrak{q}}(j(E)) < 0$. This means that E has potentially multiplicative reduction at $\mathfrak{q}$.

Recall that S is a finite set of rational primes containing 2, 3 and 5. The third condition in the hypothesis of Theorem 3 guarantees that if $q \notin S$ is a prime that divides $\mathrm{Norm}_{K/\mathbb{Q}}(abc)$, then E has multiplicative reduction at all primes $\mathfrak{q}$ of $\mathcal{O}_K$ above q. At the same time, from the formula for the discriminant $\Delta(E) = 2^4(abc)^{2p}$ we note that if $q \notin S$ and $q \nmid \mathrm{Norm}_{K/\mathbb{Q}}(abc)$, then E has good reduction at all primes $\mathfrak{q}$ above q.

If we ensure that $C_S > 71$, from Theorem 2 it follows that E cannot have multiplicative reduction at the primes above q if $q \notin S$. Notice that here we use the fourth hypothesis of the theorem, namely that if p splits in K, then at most one of the primes above p divides abc. This is to exclude the possibility that $q = p$ when q splits in K, as that case is not covered by Theorem 2. Therefore, E has good

reduction at such primes and, therefore, $j(E)$ is integral outside of S. In particular, abc is supported only on primes that lie above those in S.

Writing the Fermat equation as $(-a/c)^p + (-b/c)^p = 1$, we note that $(-a/c)^p$ and $(-b/c)^p$ are solutions to the S-unit equation $x + y = 1$, where $x, y \in O_{K,S}^{\times}$.

Using Siegel's theorem, we deduce that every such S-unit equation has finitely many solutions. Suppose that one of a, b or c is not a unit in O_K. Without losing generality, we can assume that a is divisible by some prime ideal $\mathfrak{q}$ of K. As the equation $x + y = 1$ has finitely many solutions $x, y \in O_{K,S}^{\times}$, the possible valuations of $v_{\mathfrak{q}}(x)$ belong to a finite set. A contradiction can be reached by increasing the exponent p. This proves that there is a bound $B_{K,S}$ such that if $p > B_{K,S}$ then a, b, c are units in O_K.

Let us now take C_S to be larger than all the constants $B_{K,S}$ arising from all the six possibilities for K. For each of the number fields K, there are finitely many units in O_K. Verifying all the possibilities, we find that if p and a, b, c are as in the hypothesis of Theorem 3, the only possible solutions arise when $K = \mathbb{Q}(\sqrt{-3})$ and these are $(a, b, c) = (1, \epsilon, \epsilon^2)$, up to reordering.

4.2 Proof of Theorem 4 as an application to Proposition 3

Suppose that $a, b, c \in O_K \setminus \{0\}$ are coprime and satisfy the Fermat equation $a^p + b^p + c^p = 0$. Following carefully the discussion at the beginning of this section, one can see that the conclusion of Theorem 4 follows if $\overline{\rho}_{E,p}$ is irreducible. Hence, for $p > C_K$ (a constant that depends on the field K), if $a, b, c \in O_K \setminus \{0\}$ is a solution such that $\{a, b, c\} \neq \{1, \epsilon, \epsilon^2\}$, then $\overline{\rho}_{E,p}$ must be reducible, that is, E has a p-isogeny.

We remarked previously that $2O_K$ is a prime of potentially good reduction for E. Recall that we are under the assumption that p is inert in K. Applying Proposition 3, it follows that there exists a constant $B_{K,2}$ such that $p \leq B_{K,2}$. The conclusion of Theorem 4 follows if we set $B_K := \max(C_K, B_{K,2})$.

Remark 1 We note that if $K = \mathbb{Q}(\sqrt{-3})$, the prime p splits in K if and only if $p \equiv 1 \pmod 3$. In that case, Theorem 4 is stronger than Theorem 3. However, for the other five quadratic imaginary fields discussed here, namely $K = \mathbb{Q}(\sqrt{-d})$ where $d \in \{11, 19, 43, 67, 163\}$ the set $A = \{p \text{ is prime } \mid p \equiv 1 \pmod 3\}$ has strictly larger (Dirichlet) density than the set $B = \{p \text{ is prime } \mid p \equiv 1 \pmod 3 \text{ and } p \text{ is inert in } K\}$. Hence, for these latter choices of K, Theorem 3 asserts that a version of the Asymptotic Fermat's Last Theorem holds for a larger set of exponents p than Theorem 4.

References

1. Banwait, B.S.: Explicit isogenies of prime degree over quadratic fields. Int. Math. Res. Not (2022). Rnac134
2. Banwait, B.S., Derickx, M.: Explicit isogenies of prime degree over number fields. arXiv e-prints (2022)
3. Caraiani, A., Newton, J.: On the modularity of elliptic curves over imaginary quadratic fields. arXiv e-prints (2023)
4. Deligne, P., Rapoport, M.: Les schémas de modules de courbes elliptiques. In: Modular functions of one variable, II (Proc. Internat. Summer School, Univ. Antwerp, Antwerp, 1972), *Lecture Notes in Math.*, vol. 349, pp. 143–316. Springer, Berlin, Germany (1973)
5. Derickx, M., Kamienny, S., Stein, W., Stoll, M.: Torsion points on elliptic curves over number fields of small degree. Algebra Number Theory **17**(2), 267–308 (2023)
6. Freitas, N., Kraus, A., Siksek, S.: Class field theory, Diophantine analysis and the asymptotic Fermat's Last Theorem. Adv. Math. **363**, 106964 (2020)
7. Freitas, N., Siksek, S.: An asymptotic Fermat's Last Theorem over five-sixths of real quadratic fields. Compos. Math. **151**, 1395–1415 (2015)
8. Hartshorne, R.: Algebraic Geometry, *Graduate Texts in Mathematics*, vol. 52. Springer New York, NY (1977)
9. Kamienny, S.: Torsion points on elliptic curves and q-coefficients of modular forms. Invent. Math. **109**(1), 221–229 (1992)
10. Mazur, B.: Rational points of abelian varieties with values in towers of number fields. Invent. Math. **18**, 183–266 (1972)
11. Mazur, B.: Modular curves and the Eisenstein ideal. Inst. Hautes Études Sci. Publ. Math. pp. 33–186 (1978)
12. Mazur, B.: Rational isogenies of prime degree. Invent. Math. **44**, 129–162 (1978)
13. Merel, L.: Bornes pour la torsion des courbes elliptiques sur les corps de nombres. Invent. Math. **124**, 437–449 (1996)
14. Michaud-Jacobs, P.: Fermat's last theorem and modular curves over real quadratic fields. Acta Arith. **203**, 319–351 (2022)
15. Michaud-Jacobs, P.: Mazur's isogeny theorem. ArXiv e-prints (2022)
16. Michaud-Jacobs, P.: On elliptic curves with p-isogenies over quadratic fields. Canad. J. Math. **75**(3), 945–964 (2022)
17. Milne, J.S.: Jacobian Varieties. In: Cornell, G. Silvermann, J. (eds.) Arithmetic Geometry, pp. 197–212. Springer, New York, NY (1986)
18. Momose, F.: Isogenies of prime degree over number fields. Compos. Math. **95**(3), 329–348 (1995)
19. Mumford D., B.: The Red Book on Varieties and Schemes, second edn. Lecture Notes in Mathematics. Springer Berlin, Heidelberg (1999)
20. Najman, F., Ţurcaş, G.C.: Irreducibility of mod p Galois representations of elliptic curves with multiplicative reduction over number fields. Int. J. Number Theory **17**(08), 1729–1738 (2021)
21. Şengün, M.H., Siksek, S.: On the asymptotic Fermat's Last Theorem over number fields. Commentarii Matematici Helvetici **93**, 359–372 (2018)
22. Serre, J.P.: Propriétés galoisiennes des points d'ordre fini des courbes elliptiques. Invent. Math. **15**, 259–331 (1971/72)
23. Serre, J.P.: Quelques applications du théorème de densité de Chebotarev. Publ. Math. Inst. Hautes Études Sci. **54**, 323–401 (1981)
24. Ţurcaş, G.C.: On Fermat's equation over some quadratic imaginary number fields. Research in Number Theory **4**(2), 1–24 (2018)
25. Ţurcaş, G.C.: On Serre's modularity conjecture and Fermat's equation over quadratic imaginary fields of class number one. J. Number Theory **209**, 516–530 (2020)