arXiv:2310.14878v1 [math.PR] 23 Oct 2023

A note on the Hitczenko-Kwapien conjecture about a Rademacher sequence

Shi-Zhen Liu, Ze-Yu Tao, Ze-Chun Hu*

College of Mathematics, Sichuan University, Chengdu 610065, China

Abstract

Let (ϵ_i) be a Rademacher sequence, i.e., a sequence of independent and identically distributed random variables satisfying $P(\epsilon_i = 1) = P(\epsilon_i = -1) = 1/2$. Set $S_n = a_1\epsilon_1 + \cdots + a_n\epsilon_n$ for $a = (a_1, \ldots, a_n) \in \mathbb{R}^n$. The Hitczenko-Kwapien conjecture says that $P(|S_n| \geq \|a\|) \geq 7/32$ for all $a \in \mathbb{R}^n$ and $n \in \mathbb{N}$. Up to now, we know that it holds when $n \leq 7$. In this note, we show that it holds when $n = 8$.

Keywords: Rademacher sequence; Hitczenko-Kwapien conjecture; probability inequality

1 Introduction and main result

Throughout this note, we let $\{\epsilon_i, i = 1, 2, \ldots\}$ be a Rademacher sequence, i.e., a sequence of independent random variables with $P(\epsilon_i = -1) = P(\epsilon_i = 1) = 1/2, \forall i \geq 1$. Let $n \in \mathbb{N}$ and $\epsilon = (\epsilon_1, \ldots, \epsilon_n)$. For $a = (a_1, \ldots, a_n) \in \mathbb{R}^n$, denote $a \cdot \epsilon := a_1\epsilon_1 + \cdots + a_n\epsilon_n$. The Hitczenko-Kwapien conjecture can be formulated as

$$P(|a \cdot \epsilon| \geq \|a\|) \geq \frac{7}{32}.$$

Hereafter $\|a\| := (\sum_{i=1}^n {a_i}^2)^{1/2}$.

Consider the non-increasing sequence of constants

$$G_n = \inf\{P(|a \cdot \epsilon| \geq \|a\|) : a \in \mathbb{R}^n\}, \quad n \in \mathbb{N}.$$

$G = \inf_n\{G_n\}$ is known as Hitczenko and Kwapień's constant. This constant has been applied to Kolmogorov-type lower estimates for independent random variables and other problems (cf. [1, 3]). Equivalently, the Hitczenko-Kwapien conjecture says that $G = \frac{7}{32}$.

*Corresponding author: zchu@scu.edu.cn

Burkholder [1] showed that $G > 0$. Hitczenko and Kwapień [3] showed that $G > e^{-4}/4$. Oleszkiewicz [7] showed that $G > \frac{1}{10}$, pointed out that G might be equal to $\frac{7}{32}$ and proved that $G_6 = \frac{7}{32}$. It can be checked that $G_1 = 1,\ G_2 = \frac{1}{2},\ G_3 = G_4 = G_5 = \frac{1}{4}$. Hu et al. [4] proved that $G_7 = \frac{7}{32}$. Recently, Dvorak and Klein [2] proved that $G_n \geq \frac{6}{32}$ for any $n \in \mathbb{N}$. The goal of this note is to prove

Theorem 1.1. $G_8 = \frac{7}{32}$.

The rest of this note is organized as follows. In Section 2, we give the proof of Theorem 1.1. In Section 3, we give some remarks.

2 Proof of Theorem 1.1

In this section, we will give the proof of Theorem 1.1. For convenience, we give some notations at first. By the symmetry of (ϵ_i), we can assume without loss of generality that $a_1 \geq a_2 \geq \cdots \geq a_8 \geq 0$. Following [4], we define

$$R_d^{(8)} := \{(a_1, \cdots, a_8) \in \mathbb{R}^8 : a_1 \geq a_2 \geq \cdots \geq a_8 \geq 0\}.$$

For any sign vector $s = (s_1, \cdots, s_8) \in \{-1, 1\}^8$ and $a = (a_1, \cdots, a_8) \in R_d^{(8)}$, set $a \cdot s = a_1 s_1 + \cdots + a_n s_n$. Define

$$V_{sd}(a) := \{s \in \{-1, 1\}^8 : a \cdot s \geq \|a\|\}.$$

To finish the proof, we can assume without loss of generality that $\|a\| = 1$ and will estimate the cardinality of the set $V_{sd}(a)$.

For $J \subset I_8 = \{1, \cdots, 8\}$, let $(J)_8$ denote the sign vector $s = (s_1, \cdots, s_8) \in \{-1, 1\}^8$ such that $s_j = -1$ for $j \in J$ and $s_j = 1$ otherwise. Obviously, $J \mapsto (J)_8$ sets up a one-to-one correspondence between 2^{I_8} and $\{-1, 1\}^8$. For example, $(4)_8$ and $(5, 7)_8$ denote $(1, 1, 1, -1, 1, 1, 1, 1)$ and $(1, 1, 1, 1, -1, 1, -1, 1)$, respectively. In the case that $J = \varnothing$, we just write $()_8$ for $(1, 1, 1, 1, 1, 1, 1, 1)$.

A subset A of $V_{sd}(a)$ is called a *good set* if there exists at least one element in A belonging to $V_{sd}(a)$.

In order to complete the proof of Theorem 1.1, we prove three lemmas at first.

Lemma 2.1. *Define*

$$\begin{array}{llll} A_1 := \{(4, 7)_8\}, & A_2 := \{(4, 8)_8\}, & A_3 := \{(5, 7)_8\}, & A_4 := \{(5, 8)_8\}, \\ A_5 := \{(6, 7)_8\}, & A_6 := \{(6, 8)_8\}, & A_7 := \{(7, 8)_8\}, & A_8 := \{(4)_8\}, \\ A_9 := \{(5)_8\}, & A_{10} := \{(6)_8\}, & A_{11} := \{(7)_8\}, & A_{12} := \{(8)_8\}, \\ A_{13} := \{()_8\}. & & & \end{array}$$

Then $A_1, \cdots, A_{13}$ are good sets.

Proof. By the assumption that $a_1 \geq a_2 \geq \cdots \geq a_8 \geq 0$, it is enough to show that $(4, 7)_8 \in V_{sd}(a)$. By the definition of $V_{sd}(a)$ and the assumptions on $(a_1, \cdots, a_8)$, it is enough to show that

$$a_1 + a_2 + a_3 - a_4 + a_5 + a_6 - a_7 + a_8 \geq 1. \tag{2.1}$$

We have

$$\begin{aligned}
&(a_1+a_2+a_3-a_4+a_5+a_6-a_7+a_8)^2-1\\
=&(a_1+a_2+a_3-a_4+a_5+a_6-a_7+a_8)^2-\sum_{i=1}^{8}a_i^2\\
\geq&(a_1+a_2+a_3-a_4+a_5+a_6-a_7)^2+a_8^2-\sum_{i=1}^{8}a_i^2\\
=&2[a_1a_2+a_1a_3-a_1a_4+a_1a_5+a_1a_6-a_1a_7\\
&+a_2a_3-a_2a_4+a_2a_5+a_2a_6-a_2a_7\\
&-a_3a_4+a_3a_5+a_3a_6-a_3a_7\\
&-a_4a_5-a_4a_6+a_4a_7\\
&+a_5a_6-a_5a_7-a_6a_7]\\
=&2[a_1(a_3-a_4)+a_1a_5+a_1(a_6-a_7)+a_2(a_3-a_4)+a_2(a_6-a_7)\\
&+(a_1a_2-a_3a_4)+a_3(a_6-a_7)+(a_3-a_4)a_5+(a_2a_5-a_4a_6)\\
&+a_5(a_6-a_7)+(a_4-a_6)a_7]\\
\geq&0,
\end{aligned}$$

which together with $a_1+a_2+(a_3-a_4)+a_5+(a_6-a_7)+a_8\geq 0$ implies that

$$a_1+a_2+a_3-a_4+a_5+a_6-a_7+a_8\geq 1,$$

i.e. (2.1) holds. The proof is complete. □

Define $b_8=a_8$ and $b_i=a_i-a_{i+1}$ for $i=1,\cdots,7$. Then by the assumptions on $(a_1,\cdots,a_8)$, we know that $b_i\geq 0,\forall i=1,\cdots,8$, and

$$a_i=b_i+\cdots+b_8,\forall i=1,\cdots,8. \tag{2.2}$$

Lemma 2.2. *Define*

$$\begin{aligned}
&A_{14}:=\{(5,6)_8,(2,7,8)_8\}, \quad A_{15}:=\{(2,7)_8,(5,6,8)_8\}, \quad A_{16}:=\{(2,3)_8,(6,7,8)_8\},\\
&A_{17}:=\{(2,8)_8,(4,6,8)_8\}, \quad A_{18}:=\{(3,4)_8,(5,7,8)_8\}, \quad A_{19}:=\{(3,5)_8,(4,7,8)_8\},\\
&A_{20}:=\{(3)_8,(4,5,6,7,8)_8\}.
\end{aligned}$$

Then $A_{14},\cdots,A_{20}$ are good sets.

Proof. (i) By (2.2) and symbolic calculation (by mathematical software), we get

$$\begin{aligned}
&(a_1+a_2+a_3+a_4-a_5-a_6+a_7+a_8)^2+(a_1-a_2+a_3+a_4+a_5+a_6-a_7-a_8)^2-2\\
=&(a_1+a_2+a_3+a_4-a_5-a_6+a_7+a_8)^2+(a_1-a_2+a_3+a_4+a_5+a_6-a_7-a_8)^2-2\sum_{i=1}^{8}a_i^2\\
=&4b_1b_3+4b_2b_3+4b_3^2+8b_1b_4+8b_2b_4+16b_3b_4+12b_4^2+8b_1b_5+4b_2b_5+12b_3b_5
\end{aligned}$$

$$
\begin{aligned}
&+20b_4b_5+8b_5^2+8b_1b_6+8b_3b_6+16b_4b_6+16b_5b_6+8b_6^2+8b_1b_7+4b_2b_7+12b_3b_7+20b_4b_7\\
&+16b_5b_7+12b_6b_7+4b_7^2+8b_1b_8+8b_2b_8+16b_3b_8+24b_4b_8+16b_5b_8+8b_6b_8+8b_7b_8+4b_8^2\\
\geq&0,
\end{aligned}
$$

which together with

$$
\begin{aligned}
a_1+a_2+a_3+a_4-a_5-a_6+a_7+a_8&=a_1+a_2+(a_4-a_5)+(a_3-a_6)+a_7+a_8\geq 0,\\
a_1-a_2+a_3+a_4+a_5+a_6-a_7-a_8&=(a_1-a_2)+a_3+a_4+(a_6-a_7)+(a_5-a_8)\geq 0,
\end{aligned}
$$

implies that either

$$a_1+a_2+a_3+a_4-a_5-a_6+a_7+a_8\geq 1.$$

or

$$a_1-a_2+a_3+a_4+a_5+a_6-a_7-a_8\geq 1,$$

It follows that A_{14} is a good set.

(ii) By (2.2) and symbolic calculation, we get

$$
\begin{aligned}
&(a_1-a_2+a_3+a_4+a_5+a_6-a_7+a_8)^2+(a_1+a_2+a_3+a_4-a_5-a_6+a_7-a_8)^2-2\\
=&(a_1-a_2+a_3+a_4+a_5+a_6-a_7+a_8)^2+(a_1+a_2+a_3+a_4-a_5-a_6+a_7-a_8)^2-2\sum_{i=1}^{8}a_i^2\\
=&4b_1b_3+4b_2b_3+4b_3^2+8b_1b_4+8b_2b_4+16b_3b_4+12b_4^2+8b_1b_5+4b_2b_5+12b_3b_5\\
&+20b_4b_5+8b_5^2+8b_1b_6+8b_3b_6+16b_4b_6+16b_5b_6+8b_6^2+8b_1b_7+4b_2b_7+12b_3b_7\\
&+20b_4b_7+16b_5b_7+12b_6b_7+4b_7^2+8b_1b_8+8b_3b_8+16b_4b_8+16b_5b_8+16b_6b_8+8b_7b_8+4b_8^2\\
\geq&0,
\end{aligned}
$$

which together with

$$
\begin{aligned}
a_1-a_2+a_3+a_4+a_5+a_6-a_7+a_8&=(a_1-a_2)+a_3+a_4+a_5+(a_6-a_7)+a_8\geq 0,\\
a_1+a_2+a_3+a_4-a_5-a_6+a_7-a_8&=a_1+a_2+(a_4-a_5)+(a_3-a_6)+(a_7-a_8)\geq 0,
\end{aligned}
$$

implies that either

$$a_1-a_2+a_3+a_4+a_5+a_6-a_7+a_8\geq 1.$$

or

$$a_1+a_2+a_3+a_4-a_5-a_6+a_7-a_8\geq 1,$$

It follows that A_{15} is a good set.

(iii) By (2.2) and symbolic calculation, we get

$$
\begin{aligned}
&(a_1-a_2-a_3+a_4+a_5+a_6+a_7+a_8)^2+(a_1+a_2+a_3+a_4+a_5-a_6-a_7-a_8)^2-2\\
=&(a_1-a_2-a_3+a_4+a_5+a_6+a_7+a_8)^2+(a_1+a_2+a_3+a_4+a_5-a_6-a_7-a_8)^2-2\sum_{i=1}^{8}a_i^2\\
=&4b_2b_3+4b_3^2+4b_1b_4+8b_2b_4+12b_3b_4+8b_4^2+8b_1b_5+12b_2b_5+16b_3b_5+24b_4b_5+16b_5^2
\end{aligned}
$$

$$
\begin{aligned}
&+8b_1b_6+8b_2b_6+8b_3b_6+16b_4b_6+24b_5b_6+8b_6^2+8b_1b_7+4b_2b_7+8b_4b_7+16b_5b_7+12b_6b_7\\
&+4b_7^2+8b_1b_8-8b_3b_8+8b_5b_8+8b_6b_8+8b_7b_8+4b_8^2\\
\geq&4b_3^2+4b_8^2-8b_3b_8=4(b_3-b_8)^2\geq 0,
\end{aligned}
$$

which implies that either

$$(a_1-a_2-a_3+a_4+a_5+a_6+a_7+a_8)^2\geq 1, \tag{2.3}$$

or

$$(a_1+a_2+a_3+a_4+a_5-a_6-a_7-a_8)^2\geq 1. \tag{2.4}$$

(iii.1) Assume that (2.4) holds. By the fact that $a_1+a_2+a_3+a_4+a_5-a_6-a_7-a_8=a_1+a_2+(a_5-a_6)+(a_4-a_7)+(a_3-a_8)\geq 0$, we get that

$$a_1+a_2+a_3+a_4+a_5-a_6-a_7-a_8\geq 1.$$

Thus $(5,6,7)_8\in V_{sd}(a)$.

(iii.2) Assume (2.3) holds. If $a_1-a_2-a_3+a_4+a_5+a_6+a_7+a_8\geq 0$, then by (2.3), we have $a_1-a_2-a_3+a_4+a_5+a_6+a_7+a_8\geq 1$, and thus $(2,3)_8\in V_{sd}(a)$. If $a_1-a_2-a_3+a_4+a_5+a_6+a_7+a_8<0$, then by (2.3), we have $a_1-a_2-a_3+a_4+a_5+a_6+a_7+a_8\leq -1$, which implies that

$$-(a_1-a_2-a_3+a_4+a_5+a_6+a_7+a_8)\geq 1.$$

It follows that

$$
\begin{aligned}
&a_1+a_2+a_3+a_4+a_5-a_6-a_7-a_8\\
=&-(a_1-a_2-a_3+a_4+a_5+a_6+a_7+a_8)+2(a_1+a_4+a_5)\\
\geq&1+2(a_1+a_4+a_5)>1.
\end{aligned}
$$

And thus $(6,7,8)_8\in V_{sd}(a)$.

In a word, A_{16} is always a good set.

(iv) By (2.2) and symbolic calculation, we get

$$
\begin{aligned}
&(a_1-a_2+a_3+a_4+a_5+a_6+a_7-a_8)^2+(a_1+a_2+a_3-a_4+a_5-a_6+a_7-a_8)^2-2\\
=&(a_1-a_2+a_3+a_4+a_5+a_6+a_7-a_8)^2+(a_1+a_2+a_3-a_4+a_5-a_6+a_7-a_8)^2-2\sum_{i=1}^{8}a_i^2\\
=&4b_1b_3+4b_2b_3+4b_3^2+4b_1b_4+4b_3b_4+8b_1b_5+4b_2b_5+12b_3b_5+8b_4b_5+8b_5^2+8b_1b_6\\
&+8b_3b_6+8b_4b_6+16b_5b_6+8b_6^2+12b_1b_7+4b_2b_7+16b_3b_7+16b_4b_7+28b_5b_7+28b_6b_7\\
&+20b_7^2+8b_1b_8+8b_3b_8+8b_4b_8+16b_5b_8+16b_6b_8+24b_7b_8+4b_8^2\\
\geq&0,
\end{aligned}
$$

which together with

$$a_1-a_2+a_3+a_4+a_5+a_6+a_7-a_8=(a_1-a_2)+a_3+a_4+a_5+a_6+(a_7-a_8)\geq 0,$$

$$a_1+a_2+a_3-a_4+a_5-a_6+a_7-a_8=a_1+a_2+(a_3-a_4)+(a_5-a_6)+(a_7-a_8)\geq 0,$$

implies that either

$$a_1-a_2+a_3+a_4+a_5+a_6+a_7-a_8\geq 1.$$

or

$$a_1+a_2+a_3-a_4+a_5-a_6+a_7-a_8\geq 1,$$

It follows that A_{17} is a good set.

(v) By (2.2) and symbolic calculation, we get

$$\begin{aligned}
&(a_1+a_2-a_3-a_4+a_5+a_6+a_7+a_8)^2+(a_1+a_2+a_3+a_4-a_5+a_6-a_7-a_8)^2-2\\
=&(a_1+a_2-a_3-a_4+a_5+a_6+a_7+a_8)^2+(a_1+a_2+a_3+a_4-a_5+a_6-a_7-a_8)^2-2\sum_{i=1}^{8}a_i^2\\
=&4b_1b_2+4b_2^2+4b_1b_3+8b_2b_3+4b_3^2+4b_1b_4+8b_2b_4+12b_3b_4+8b_4^2+4b_1b_5+8b_2b_5\\
&+8b_3b_5+8b_4b_5+8b_1b_6+16b_2b_6+16b_3b_6+16b_4b_6+8b_5b_6+8b_6^2+8b_1b_7+16b_2b_7\\
&+12b_3b_7+8b_4b_7+4b_5b_7+12b_6b_7+4b_7^2+8b_1b_8+16b_2b_8+8b_3b_8+8b_6b_8+8b_7b_8+4b_8^2\\
\geq&0,
\end{aligned}$$

which together with

$$\begin{aligned}
&a_1+a_2-a_3-a_4+a_5+a_6+a_7+a_8=(a_2-a_3)+(a_1-a_4)+a_5+a_6+a_7+a_8\geq 0,\\
&a_1+a_2+a_3+a_4-a_5+a_6-a_7-a_8=a_1+a_2+(a_4-a_5)+(a_6-a_7)+(a_3-a_8)\geq 0,
\end{aligned}$$

implies that either

$$a_1+a_2-a_3-a_4+a_5+a_6+a_7+a_8\geq 1.$$

or

$$a_1+a_2+a_3+a_4-a_5+a_6-a_7-a_8\geq 1,$$

It follows that A_{18} is a good set.

(vi) By (2.2) and symbolic calculation, we get

$$\begin{aligned}
&(a_1+a_2-a_3+a_4-a_5+a_6+a_7+a_8)^2+(a_1+a_2+a_3-a_4+a_5+a_6-a_7-a_8)^2-2\\
=&(a_1+a_2-a_3+a_4-a_5+a_6+a_7+a_8)^2+(a_1+a_2+a_3-a_4+a_5+a_6-a_7-a_8)^2-2\sum_{i=1}^{8}a_i^2\\
=&4b_1b_2+4b_2^2+4b_1b_3+8b_2b_3+4b_3^2+4b_1b_4+8b_2b_4+4b_3b_4+4b_1b_5+8b_2b_5+8b_3b_5\\
&+8b_1b_6+16b_2b_6+16b_3b_6+8b_4b_6+8b_5b_6+8b_6^2+8b_1b_7+16b_2b_7+12b_3b_7+8b_4b_7+4b_5b_7\\
&+12b_6b_7+4b_7^2+8b_1b_8+16b_2b_8+8b_3b_8+8b_4b_8+8b_6b_8+8b_7b_8+4b_8^2\\
\geq&0,
\end{aligned}$$

which together with

$$a_1+a_2-a_3+a_4-a_5+a_6+a_7+a_8=a_1+(a_2-a_3)+(a_4-a_5)+a_6+a_7+a_8\geq 0,$$

$$a_1+a_2+a_3-a_4+a_5+a_6-a_7-a_8=a_1+a_2+(a_3-a_4)+(a_6-a_7)+(a_5-a_8)\geq 0,$$

implies that either

$$a_1+a_2-a_3+a_4-a_5+a_6+a_7+a_8\geq 1.$$

or

$$a_1+a_2+a_3-a_4+a_5+a_6-a_7-a_8\geq 1,$$

It follows that A_{19} is a good set.

(vii) By (2.2) and symbolic calculation, we get

$$\begin{aligned}
&(a_1+a_2-a_3+a_4+a_5+a_6+a_7+a_8)^2+(a_1+a_2+a_3-a_4-a_5-a_6-a_7-a_8)^2-2\\
=&(a_1+a_2-a_3+a_4+a_5+a_6+a_7+a_8)^2+(a_1+a_2+a_3-a_4-a_5-a_6-a_7-a_8)^2-2\sum_{i=1}^{8}a_i^2\\
=&4b_1b_2+4b_2^2+4b_1b_3+8b_2b_3+4b_3^2+4b_1b_4+8b_2b_4+4b_3b_4+4b_1b_5+8b_2b_5+4b_1b_6+8b_2b_6\\
&-4b_3b_6+4b_5b_6+4b_6^2+4b_1b_7+8b_2b_7-8b_3b_7+8b_5b_7+16b_6b_7+12b_7^2+4b_1b_8+8b_2b_8\\
&-12b_3b_8+12b_5b_8+24b_6b_8+36b_7b_8+24b_8^2\\
\geq&4b_3^2-4b_3b_6+4b_6^2-8b_3b_7+16b_6b_7+12b_7^2-12b_3b_8+24b_6b_8+36b_7b_8+24b_8^2\\
=&4\left[b_3^2-(b_6+b_7+3b_8)b_3+(b_6^2+4b_6b_7+3b_7^2+6b_6b_89+b_7b_8+6b_8^2)\right]\\
:=&4f(b_3),
\end{aligned}$$

where $f(x)=x^2-(b_6+2b_7+3b_8)x+(b_6^2+4b_6b_7+3b_7^2+6b_6b_8+9b_7b_8+6b_8^2)$. Then we have

$$\begin{aligned}
4f(b_3)&\geq 4f\left(\frac{b_6}{2}+b_7+\frac{3b_8}{2}\right)\\
&=3b_6^2+12b_6b_7+8b_7^2+18b_6b_8+24b_7b_8+15b_8^2\\
&\geq 0.
\end{aligned}$$

Thus

$$(a_1+a_2-a_3+a_4+a_5+a_6+a_7+a_8)^2+(a_1+a_2+a_3-a_4-a_5-a_6-a_7-a_8)^2-2\geq 0,$$

which implies that either

$$(a_1+a_2-a_3+a_4+a_5+a_6+a_7+a_8)^2\geq 1, \tag{2.5}$$

or

$$(a_1+a_2+a_3-a_4-a_5-a_6-a_7-a_8)^2\geq 1. \tag{2.6}$$

(vii.1) Assume that (2.5) holds. By the fact that $a_1+a_2-a_3+a_4+a_5+a_6+a_7+a_8=a_1+(a_2-a_3)+a_4+a_5+a_6+a_7+a_8\geq 0$, we get

$$a_1+a_2-a_3+a_4+a_5+a_6+a_7+a_8\geq 1,$$

Thus $(3)_8 \in V_{sd}(a)$.

(vii.2) Assume that (2.6) holds. If $a_1+a_2+a_3-a_4-a_5-a_6-a_7-a_8 \geq 0$, then $a_1+a_2+a_3-a_4-a_5-a_6-a_7-a_8 \geq 1$, and thus $(4,5,6,7,8)_8 \in V_{sd}(a)$. If $a_1+a_2+a_3-a_4-a_5-a_6-a_7-a_8 < 0$, then by (2.6), we have $a_1+a_2+a_3-a_4-a_5-a_6-a_7-a_8 \leq -1$, which implies that

$$-(a_1+a_2+a_3-a_4-a_5-a_6-a_7-a_8) \geq 1.$$

It follows that

$$\begin{aligned} & a_1+a_2-a_3+a_4+a_5+a_6+a_7+a_8 \\ =& -(a_1+a_2+a_3-a_4-a_5-a_6-a_7-a_8)+2(a_1+a_2) \\ \geq& 1+2(a_1+a_2) > 1. \end{aligned}$$

And thus $(3)_8 \in V_{sd}(a)$.

In a word, A_{20} is always a good set. □

Lemma 2.3. *Define*

$A_{21} := \{(3,8)_8, (4,5,6,7)_8\}$, $A_{22} := \{(3,7)_8, (4,5,6,8)_8\}$,
$A_{23} := \{(2)_8, (4,6,7)_8\}$, $A_{24} := \{(4,6)_8, (1,8)_8\}$,
$A_{25} := \{(3,6)_8, (4,5,7,8)_8, (1,7)_8\}$, $A_{26} := \{(2,6)_8, (3,6,7)_8, (4,5,8)_8, (1,7,8)_8\}$,
$A_{27} := \{(4,5)_8, (1)_8, (5,6,7)_8, (3,6,8)_8\}$, $A_{28} := \{(3,7,8)_8, (4,5,7)_8, (1,6)_8, (2,5)_8, (5,6,7,8)_8\}$.

Then $A_{21}, \cdots, A_{28}$ are good sets.

Proof. Obviously, it is enough to prove that

(1) if $a_1+a_2+a_3-a_4-a_5-a_6-a_7+a_8 \leq 1$, then $a_1+a_2-a_3+a_4+a_5+a_6+a_7-a_8 \geq 1$;

(2) if $a_1+a_2-a_3+a_4+a_5+a_6-a_7+a_8 \leq 1$, then $a_1+a_2+a_3-a_4-a_5-a_6+a_7-a_8 \geq 1$;

(3) if $a_1-a_2+a_3+a_4+a_5+a_6+a_7+a_8 \leq 1$, then $a_1+a_2+a_3-a_4+a_5-a_6-a_7+a_8 \geq 1$;

(4) if $-a_1+a_2+a_3+a_4+a_5+a_6+a_7-a_8 \leq 1$, then $a_1+a_2+a_3-a_4+a_5-a_6+a_7+a_8 \geq 1$;

(5) if $a_1+a_2+a_3-a_4-a_5+a_6-a_7-a_8 \leq 1$ and $-a_1+a_2+a_3+a_4+a_5+a_6-a_7+a_8 \leq 1$, then $a_1+a_2-a_3+a_4+a_5-a_6+a_7+a_8 \geq 1$;

(6) if $a_1+a_2-a_3+a_4+a_5-a_6-a_7+a_8 \leq 1, a_1+a_2+a_3-a_4-a_5+a_6+a_7-a_8 \leq 1$ and $-a_1+a_2+a_3+a_4+a_5+a_6-a_7-a_8 \leq 1$, then $a_1-a_2+a_3+a_4+a_5-a_6+a_7+a_8 \geq 1$;

(7) if $-a_1+a_2+a_3+a_4+a_5+a_6+a_7+a_8 \leq 1, a_1+a_2+a_3+a_4-a_5-a_6-a_7+a_8 \leq 1$ and $a_1+a_2-a_3+a_4+a_5-a_6+a_7-a_8 \leq 1$, then $a_1+a_2+a_3-a_4-a_5+a_6+a_7+a_8 \geq 1$;

(8) if $a_1+a_2+a_3-a_4-a_5+a_6-a_7+a_8 \leq 1, -a_1+a_2+a_3+a_4+a_5-a_6+a_7+a_8 \leq 1, a_1-a_2+a_3+a_4-a_5+a_6+a_7+a_8 \leq 1$ and $a_1+a_2+a_3+a_4-a_5-a_6-a_7-a_8 \leq 1$, then $a_1+a_2-a_3+a_4+a_5+a_6-a_7-a_8 \geq 1$.

In the following, we give the proofs of (1)-(8).

(1) By (2.2) and symbolic calculation (by mathematical software), we have

$$
\begin{aligned}
&(a_1+a_2-a_3+a_4+a_5+a_6+a_7-a_8)^2+\frac{1}{3}(a_1+a_2+a_3-a_4-a_5-a_6-a_7+a_8)^2-\frac{4}{3}\\
=&(a_1+a_2-a_3+a_4+a_5+a_6+a_7-a_8)^2+\frac{1}{3}(a_1+a_2+a_3-a_4-a_5-a_6-a_7+a_8)^2-\frac{4}{3}\sum_{i=1}^{8}a_i^2\\
=&\frac{4}{3}\left(2b_1b_2+2b_2^2+b_1b_3+2b_2b_3+2b_1b_4+4b_2b_4\right)+\frac{8}{3}\left(b_4b_5+b_5^2+2b_1b_6+4b_2b_6+2b_4b_6+4b_5b_6\right)\\
&+4b_1b_5+8b_2b_5+8b_6^2+\frac{20}{3}(b_1b_7+2b_2b_7)+8b_4b_7+16b_5b_7+24b_6b_7+16b_7^2\\
&+\frac{16}{3}(b_1b_8+2b_2b_8+b_4b_8+2b_5b_8)+16b_6b_8+\frac{16}{3}(4b_7b_8+b_8^2)\\
\geq&0,
\end{aligned}
$$

which together with

$$
\begin{aligned}
1&\geq a_1+a_2+a_3-a_4-a_5-a_6-a_7+a_8\\
&=(a_1-a_4)+(a_2-a_5)+(a_3-a_6)-a_7+a_8\\
&\geq -a_7\geq -a_1\geq -1,
\end{aligned}
$$

implies that

$$
(a_1+a_2-a_3+a_4+a_5+a_6+a_7-a_8)^2\geq\frac{4}{3}-\frac{1}{3}(a_1+a_2+a_3-a_4-a_5-a_6-a_7+a_8)^2\geq 1.
$$

Then by $a_1+a_2-a_3+a_4+a_5+a_6+a_7-a_8=a_1+(a_2-a_3)+a_4+a_5+a_6+(a_7-a_8)\geq 0$, we get $a_1+a_2-a_3+a_4+a_5+a_6+a_7-a_8\geq 1$.

(2) By (2.2) and symbolic calculation, we have

$$
\begin{aligned}
&(a_1+a_2+a_3-a_4-a_5-a_6+a_7-a_8)^2+3(a_1+a_2-a_3+a_4+a_5+a_6-a_7+a_8)^2-4\\
=&(a_1+a_2+a_3-a_4-a_5-a_6+a_7-a_8)^2+3(a_1+a_2-a_3+a_4+a_5+a_6-a_7+a_8)^2-4\sum_{i=1}^{8}a_i^2\\
=&8b_1b_2+8b_2^2+4b_1b_3+8b_2b_3+8b_1b_4+16b_2b_4+12b_1b_5+24b_2b_5+8b_4b_5+8b_5^2+16b_1b_6\\
&+32b_2b_6+16b_4b_6+32b_5b_6+24b_6^2+12b_1b_7+24b_2b_7+8b_4b_7+16b_5b_7+24b_6b_7+16b_1b_8\\
&+32b_2b_8+16b_4b_8+32b_5b_8+48b_6b_8+16b_7b_8+16b_8^2\\
\geq&0,
\end{aligned}
$$

which together with

$$
1\geq a_1+a_2-a_3+a_4+a_5+a_6-a_7+a_8=a_1+(a_2-a_3)+a_4+a_5+(a_6-a_7)+a_8\geq 0,
$$

implies that

$$
(a_1+a_2+a_3-a_4-a_5-a_6+a_7-a_8)^2\geq 4-3(a_1+a_2-a_3+a_4+a_5+a_6-a_7+a_8)^2\geq 1.
$$

Then by $a_1+a_2+a_3-a_4-a_5-a_6+a_7-a_8=(a_1-a_4)+(a_2-a_5)+(a_3-a_6)+(a_7-a_8)\geq 0$, we get $a_1+a_2+a_3-a_4-a_5-a_6+a_7-a_8\geq 1$.

(3) By (2.2) and symbolic calculation, we have

$$\begin{aligned}
&(a_1+a_2+a_3-a_4+a_5-a_6-a_7+a_8)^2+\frac{1}{2}(a_1-a_2+a_3+a_4+a_5+a_6+a_7+a_8)^2-\frac{3}{2}\\
=&(a_1+a_2+a_3-a_4+a_5-a_6-a_7+a_8)^2+\frac{1}{2}(a_1-a_2+a_3+a_4+a_5+a_6+a_7+a_8)^2-\frac{3}{2}\sum_{i=1}^{8}a_i^2\\
=&b_1b_2+b_2^2+4b_1b_3+6b_2b_3+5b_3^2+3b_1b_4+2b_2b_4+5b_3b_4+6b_1b_5+6b_2b_5+12b_3b_5\\
&+6b_4b_5+6b_5^2+5b_1b_6+2b_2b_6+7b_3b_6+4b_4b_6+9b_5b_6+3b_6^2+4b_1b_7-2b_2b_7+2b_3b_7+2b_4b_7\\
&+6b_5b_7+6b_6b_7+3b_7^2+7b_1b_8+2b_2b_8+9b_3b_8+8b_4b_8+15b_5b_8+14b_6b_8+13b_7b_8+10b_8^2\\
\geq&b_2^2+3b_7^2-2b_2b_7\geq 0,
\end{aligned}$$

which together with

$$1\geq a_1-a_2+a_3+a_4+a_5+a_6+a_7+a_8=(a_1-a_2)+a_3+a_4+a_5+a_6+a_7+a_8\geq 0,$$

implies that

$$(a_1+a_2+a_3-a_4+a_5-a_6-a_7+a_8)^2\geq\frac{3}{2}-\frac{1}{2}(a_1-a_2+a_3+a_4+a_5+a_6+a_7+a_8)^2\geq 1.$$

Then by $a_1+a_2+a_3-a_4+a_5-a_6-a_7+a_8=a_1+(a_3-a_4)+(a_5-a_6)+(a_2-a_7)+a_8\geq 0$, we get $a_1+a_2+a_3-a_4+a_5-a_6-a_7+a_8\geq 1$.

(4) By (2.2) and symbolic calculation, we have

$$\begin{aligned}
&(a_1+a_2+a_3-a_4+a_5-a_6+a_7+a_8)^2+\frac{1}{5}(-a_1+a_2+a_3+a_4+a_5+a_6+a_7-a_8)^2-\frac{6}{5}\\
=&(a_1+a_2+a_3-a_4+a_5-a_6+a_7+a_8)^2+\frac{1}{5}(-a_1+a_2+a_3+a_4+a_5+a_6+a_7-a_8)^2-\frac{6}{5}\sum_{i=1}^{8}a_i^2\\
=&\frac{4}{5}\left(2b_1b_2+2b_2^2+4b_1b_3+9b_2b_3+7b_3^2+b_1b_4+4b_2b_4+7b_3b_4+3b_1b_5+9b_2b_5\right)+12b_3b_5\\
&+\frac{4}{5}\left(6b_4b_5+3b_5^2+4b_2b_6+8b_3b_6+2b_4b_6+6b_5b_6+b_1b_7+9b_2b_7+16b_3b_7+8b_4b_7\right)\\
&+12b_5b_7+4b_1b_8+8b_6b_8+\frac{4}{5}\left(7b_6b_7+7b_7^2+14b_2b_8+23b_3b_8+12b_4b_8+21b_5b_8+19b_7b_8+12b_8^2\right)\\
\geq&0,
\end{aligned}$$

which together with

$$\begin{aligned}
1&\geq -a_1+a_2+a_3+a_4+a_5+a_6+a_7-a_8\\
&=-a_1+a_2+a_3+a_4+a_5+a_6+(a_7-a_8)\\
&\geq -a_1\geq -1,
\end{aligned}$$

implies that

$$(a_1+a_2+a_3-a_4+a_5-a_6+a_7+a_8)^2 \geq \frac{6}{5}-\frac{1}{5}(-a_1+a_2+a_3+a_4+a_5+a_6+a_7-a_8)^2 \geq 1.$$

Then by $a_1+a_2+a_3-a_4+a_5-a_6+a_7+a_8 = a_1+a_2+(a_3-a_4)+(a_5-a_6)+a_7+a_8 \geq 0$, we get $a_1+a_2+a_3-a_4+a_5-a_6+a_7+a_8 \geq 1$.

(5) By (2.2) and symbolic calculation, we have

$$\begin{aligned}
&(a_1+a_2-a_3+a_4+a_5-a_6+a_7+a_8)^2+\frac{3}{7}(a_1+a_2+a_3-a_4-a_5+a_6-a_7-a_8)^2\\
&+\frac{2}{7}(-a_1+a_2+a_3+a_4+a_5+a_6-a_7+a_8)^2-\frac{12}{7}\\
=&(a_1+a_2-a_3+a_4+a_5-a_6+a_7+a_8)^2+\frac{3}{7}(a_1+a_2+a_3-a_4-a_5+a_6-a_7-a_8)^2\\
&+\frac{2}{7}(-a_1+a_2+a_3+a_4+a_5+a_6-a_7+a_8)^2-\frac{12}{7}\sum_{i=1}^{8}a_i^2\\
=&\frac{4}{7}\big(4b_1b_2+4b_2^2+b_1b_3+4b_2b_3+2b_1b_4+8b_2b_4+3b_1b_5+12b_2b_5+6b_4b_5\\
&+6b_5^2+8b_2b_6+2b_3b_6+4b_4b_6+6b_5b_6+3b_1b_7+12b_2b_7+6b_4b_7+12b_5b_7\\
&+4b_1b_8+16b_2b_8+12b_4b_8+24b_5b_8+8b_6b_8+12b_7b_8+12b_8^2\big)\\
\geq&0,
\end{aligned}$$

which together with

$$1 \geq a_1+a_2+a_3-a_4-a_5+a_6-a_7-a_8 = (a_1-a_4)+(a_2-a_5)+(a_3-a_7)+(a_6-a_8) \geq 0,$$

and

$$\begin{aligned}
1 &\geq -a_1+a_2+a_3+a_4+a_5+a_6-a_7+a_8\\
&= -a_1+a_2+a_3+a_4+a_5+(a_6-a_7)+a_8\\
&\geq -a_1 \geq -1,
\end{aligned}$$

implies that

$$\begin{aligned}
&(a_1+a_2-a_3+a_4+a_5-a_6+a_7+a_8)^2\\
\geq&\frac{12}{7}-\frac{3}{7}(a_1+a_2+a_3-a_4-a_5+a_6-a_7-a_8)^2-\frac{2}{7}(-a_1+a_2+a_3+a_4+a_5+a_6-a_7+a_8)^2\\
\geq&1.
\end{aligned}$$

Then by $a_1+a_2-a_3+a_4+a_5-a_6+a_7+a_8 = a_1+(a_2-a_3)+a_4+(a_5-a_6)+a_7+a_8 \geq 0$, we get $a_1+a_2-a_3+a_4+a_5-a_6+a_7+a_8 \geq 1$.

(6) By (2.2) and symbolic calculation, we have

$$(a_1-a_2+a_3+a_4+a_5-a_6+a_7+a_8)^2+\frac{3}{4}(a_1+a_2-a_3+a_4+a_5-a_6-a_7+a_8)^2-3$$

$$
\begin{aligned}
&+\frac{3}{4}(a_1+a_2+a_3-a_4-a_5+a_6+a_7-a_8)^2+\frac{1}{2}(-a_1+a_2+a_3+a_4+a_5+a_6-a_7-a_8)^2\\
=&(a_1-a_2+a_3+a_4+a_5-a_6+a_7+a_8)^2+\frac{3}{4}(a_1+a_2-a_3+a_4+a_5-a_6-a_7+a_8)^2-3\sum_{i=1}^{8}a_i^2\\
&+\frac{3}{4}(a_1+a_2+a_3-a_4-a_5+a_6+a_7-a_8)^2+\frac{1}{2}(-a_1+a_2+a_3+a_4+a_5+a_6-a_7-a_8)^2\\
=&b_1b_3+2b_1b_4+3b_1b_5+6b_4b_5+6b_5^2+2b_3b_6+4b_4b_6+6b_5b_6+3b_1b_7+6b_3b_7\\
&+6b_4b_7+6b_5b_7+6b_1b_8+4b_3b_8+8b_4b_8+12b_5b_8\\
\geq&0,
\end{aligned}
$$

which together with

$$
\begin{aligned}
&1\geq a_1+a_2-a_3+a_4+a_5-a_6-a_7+a_8=(a_2-a_3)+a_4+(a_5-a_6)+(a_1-a_7)+a_8\geq 0,\\
&1\geq a_1+a_2+a_3-a_4-a_5+a_6+a_7-a_8=a_1(a_3-a_4)+(a_2-a_5)+a_6+(a_7-a_8)\geq 0,\\
&1\geq -a_1+a_2+a_3+a_4+a_5+a_6-a_7-a_8=-a_1+a_2+a_3+a_4+(a_6-a_7)+(a_5-a_8)\geq -a_1\geq -1,
\end{aligned}
$$

implies that

$$
\begin{aligned}
&(a_1-a_2+a_3+a_4+a_5-a_6+a_7+a_8)^2\\
\geq&3-\frac{3}{4}(a_1+a_2-a_3+a_4+a_5-a_6-a_7+a_8)^2-\frac{3}{4}(a_1+a_2+a_3-a_4-a_5+a_6+a_7-a_8)^2\\
&-\frac{1}{2}(-a_1+a_2+a_3+a_4+a_5+a_6-a_7-a_8)^2\\
\geq&3-\frac{3}{4}-\frac{3}{4}-\frac{1}{2}=1.
\end{aligned}
$$

Then by $a_1-a_2+a_3+a_4+a_5-a_6+a_7+a_8=(a_1-a_2)+a_3+a_4+(a_5-a_6)+a_7+a_8\geq 0$, we get $a_1-a_2+a_3+a_4+a_5-a_6+a_7+a_8\geq 1$.

(7) By (2.2) and symbolic calculation, we have

$$
\begin{aligned}
&(a_1+a_2+a_3-a_4-a_5+a_6+a_7+a_8)^2+(-a_1+a_2+a_3+a_4+a_5+a_6+a_7+a_8)^2+\\
&+2(a_1+a_2+a_3+a_4-a_5-a_6-a_7+a_8)^2+2(a_1+a_2-a_3+a_4+a_5-a_6+a_7-a_8)^2-6\\
=&(a_1+a_2+a_3-a_4-a_5+a_6+a_7+a_8)^2+(-a_1+a_2+a_3+a_4+a_5+a_6+a_7+a_8)^2+\\
&+2(a_1+a_2+a_3+a_4-a_5-a_6-a_7+a_8)^2+2(a_1+a_2-a_3+a_4+a_5-a_6+a_7-a_8)^2-6\sum_{i=1}^{8}a_i^2\\
=&8b_1b_2+8b_2^2+8b_1b_3+20b_2b_3+12b_3^2+12b_1b_4+32b_2b_4+36b_3b_4+24b_4^2+8b_1b_5+\\
&28b_2b_5+24b_3b_5+40b_4b_5+16b_5^2+16b_2b_6+16b_3b_6+24b_4b_6+16b_5b_6+20b_2b_7+16b_3b_7+\\
&24b_4b_7+24b_5b_7+12b_6b_7+12b_7^2+24b_2b_8+32b_3b_8+40b_4b_8+32b_5b_8+24b_6b_8+32b_7b_8+20b_8^2\\
\geq&0,
\end{aligned}
$$

which together with

$$
1\geq -a_1+a_2+a_3+a_4+a_5+a_6+a_7+a_8\geq -a_1\geq -1,
$$

$$1 \geq a_1 + a_2 + a_3 + a_4 - a_5 - a_6 - a_7 + a_8 = a_1 + (a_2 - a_5) + (a_3 - a_6) + (a_4 - a_7) + a_8 \geq 0,$$
$$1 \geq a_1 + a_2 - a_3 + a_4 + a_5 - a_6 + a_7 - a_8 = a_1 + (a_2 - a_3) + a_4 + (a_5 - a_6) + (a_7 - a_8) \geq 0,$$

implies that

$$\begin{aligned}
&(a_1 + a_2 + a_3 - a_4 - a_5 + a_6 + a_7 + a_8)^2 \\
\geq& 6 - (-a_1 + a_2 + a_3 + a_4 + a_5 + a_6 + a_7 + a_8)^2 - 2(a_1 + a_2 + a_3 + a_4 - a_5 - a_6 - a_7 + a_8)^2 \\
&- 2(a_1 + a_2 - a_3 + a_4 + a_5 - a_6 + a_7 - a_8)^2 \\
\geq& 1.
\end{aligned}$$

Then by $a_1 + a_2 + a_3 - a_4 - a_5 + a_6 + a_7 + a_8 = a_1 + (a_3 - a_4) + (a_2 - a_5) + a_6 + a_7 + a_8 \geq 0$, we get $a_1 + a_2 + a_3 - a_4 - a_5 + a_6 + a_7 + a_8 \geq 1$.

(8) By (2.2) and symbolic calculation, we have

$$\begin{aligned}
&(a_1 + a_2 - a_3 + a_4 + a_5 + a_6 - a_7 - a_8)^2 + \frac{1}{2}(a_1 + a_2 + a_3 - a_4 - a_5 + a_6 - a_7 + a_8)^2 \\
&+ \frac{1}{2}(-a_1 + a_2 + a_3 + a_4 + a_5 - a_6 + a_7 + a_8)^2 + \frac{1}{2}(a_1 - a_2 + a_3 + a_4 - a_5 + a_6 + a_7 + a_8)^2 \\
&+ \frac{1}{6}(a_1 + a_2 + a_3 + a_4 - a_5 - a_6 - a_7 - a_8)^2 - \frac{8}{3} \\
=&(a_1 + a_2 - a_3 + a_4 + a_5 + a_6 - a_7 - a_8)^2 + \frac{1}{2}(a_1 + a_2 + a_3 - a_4 - a_5 + a_6 - a_7 + a_8)^2 \\
&+ \frac{1}{2}(-a_1 + a_2 + a_3 + a_4 + a_5 - a_6 + a_7 + a_8)^2 + \frac{1}{2}(a_1 - a_2 + a_3 + a_4 - a_5 + a_6 + a_7 + a_8)^2 \\
&+ \frac{1}{6}(a_1 + a_2 + a_3 + a_4 - a_5 - a_6 - a_7 - a_8)^2 - \frac{8}{3}\sum_{i=1}^{8} a_i^2 \\
=&\frac{2}{3}\left(2b_1b_2 + 2b_2^2 + b_1b_3 + 2b_2b_3 + b_1b_5 + 8b_2b_5\right) + 2b_1b_4 + 4b_2b_4 + 2b_3b_4 + 2b_4^2 + \\
&+ \frac{2}{3}\left(7b_4b_5 + 4b_5^2 + 8b_1b_6 + 16b_2b_6 + 14b_4b_6 + 14b_5b_6 10b_6^2\right) + 4b_3b_6 + 2b_1b_7 + 4b_2b_7 \\
&+ 6b_4b_7 + 2b_3b_8 + 4b_6b_8 + \frac{2}{3}\left(8b_5b_7 + 10b_6b_7 + b_1b_8 + 2b_2b_8 + 10b_4b_8 + 5b_5b_8 + b_7b_8 + b_8^2\right) \\
\geq& 0,
\end{aligned}$$

which together with

$$1 \geq a_1 + a_2 + a_3 - a_4 - a_5 + a_6 - a_7 + a_8 = a_1 + (a_3 - a_4) + (a_2 - a_5) + (a_6 - a_7) + a_8 \geq 0,$$
$$1 \geq -a_1 + a_2 + a_3 + a_4 + a_5 - a_6 + a_7 + a_8 = -a_1 + a_2 + a_3 + a_4 + (a_5 - a_6) + a_7 + a_8 \geq -a_1 \geq -1,$$
$$1 \geq a_1 - a_2 + a_3 + a_4 - a_5 + a_6 + a_7 + a_8 = (a_1 - a_2) + a_3 + (a_4 - a_5) + a_6 + a_7 + a_8 \geq 0,$$
$$1 \geq a_1 + a_2 + a_3 + a_4 - a_5 - a_6 - a_7 - a_8 = (a_1 - a_5) + (a_2 - a_6) + (a_3 - a_7) + (a_4 - a_8) \geq 0,$$

implies that

$$(a_1 + a_2 - a_3 + a_4 + a_5 + a_6 - a_7 - a_8)^2$$

$$\geq \frac{8}{3} - \frac{1}{2}(a_1+a_2+a_3-a_4-a_5+a_6-a_7+a_8)^2 - \frac{1}{2}(-a_1+a_2+a_3+a_4+a_5-a_6+a_7+a_8)^2$$
$$- \frac{1}{2}(a_1-a_2+a_3+a_4-a_5+a_6+a_7+a_8)^2 - \frac{1}{6}(a_1+a_2+a_3+a_4-a_5-a_6-a_7-a_8)^2$$
$$\geq 1.$$

Then by $a_1+a_2-a_3+a_4+a_5+a_6-a_7-a_8 = a_1+(a_2-a_3)+a_4+(a_6-a_7)+(a_5-a_8) \geq 0$, we get $a_1+a_2-a_3+a_4+a_5+a_6-a_7-a_8 \geq 1$.

The proof is complete. □

Proof of Theorem 1.1. By Lemmas 2.1-2.3, we know that $A_1, \cdots, A_{28}$ are good sets. Note that for any $i, j \in \{1, 2, \cdots, 28\}, i \neq j$, we have $A_i \cap A_j = \emptyset$. Hence for any $a = (a_1, \cdots, a_8) \in R_d^{(8)}$, the number of elements in the set $V_{sd}(a)$ is more than or equal to 28. It follows that

$$\begin{aligned} P\left(|a \cdot \epsilon| \geq \|a\|\right) &= P(\{s \in \{-1,1\}^8 : |a \cdot s| \geq \|a\|\}) \\ &= 2P(\{s \in \{-1,1\}^8 : a \cdot s \geq \|a\|\}) \\ &\geq 2 \cdot \frac{28}{2^8} = \frac{7}{32}, \end{aligned}$$

which implies that $G_8 \geq \frac{7}{32}$. By $G_8 \leq G_7 = \frac{7}{32}$, we obtain $G_8 = \frac{7}{32}$. The proof is complete. □

3 Some remarks

Remark 3.1. *As to Rademacher sequences, another famous inequality (Tomaszewski's conjecture) is*

$$P\left(|a \cdot \epsilon| \leq \|a\|\right) \geq \frac{1}{2},$$

which has been proved by Keller and Klein [5].

Remark 3.2. *In order to find 28 good sign vectors (especially on $A_{25}, \cdots, A_{28}$ defined in Lemma 2.3), we used the method of conditional minimization.*

Remark 3.3. *Our method can be applied to other similar problems. In fact, in a forthcoming paper [8], we will consider the following inequality associated with a Rademacher sequence:*

$$P\left(|a \cdot \epsilon| \geq \frac{1}{\sqrt{7}}\|a\|\right) \geq \frac{1}{2},$$

which was posed by Lowther [6].

Acknowledgments This work was supported by the National Natural Science Foundation of China (12171335) and the Science Development Project of Sichuan University (2020SCUNL201).

References

[1] D. L. Burkholder. Independent sequences with the Stein property. *Ann. Math. Statis.*, 39: 1282-1288, 1967.

[2] V. Dvorak, O. Klein. Probability mass of Rademacher sums beyond one standard deviation. *SIAM J. Discrere Math.*, 36: 2393-2410, 2022.

[3] P. Hitczenko, S. Kwapień. On the Rademacher series. In *Probability in Banach spaces, 9 (Sandjberg, 1994)*, Volume 35 of *Progr. Probab.*, 31-36. Birkhäuser Boston, Boston, MA, 1994.

[4] Z.-C. Hu, G.L. Lan, W. Sun. Some explorations on two conjectures about Rademacher Sequences. *Acta Math. Appl. Sin. (Engl. Ser.)*, 37: 1-16, 2021.

[5] N. Keller, O. Klein. Proof of Tomaszewski's conjecture on randomly signed sums. *Adv. Math.*, 407, Paper No. 108558, 2022.

[6] G. Lowther. https://mathoverflow.net/questions/366894/a-rademacher-root-7-anti-concentration- inequality.

[7] K. Oleszkiewicz. On the Stein property of Rademacher sequences. *Probab. Math. Stat.*, 16: 127-130, 1996.

[8] Z.-Y. Tao, S.-Z. Liu, Z.-C. Hu. An anti-concentration inequality beyond $\frac{1}{\sqrt{7}}$ standard deviation of Rademacher sums, In preparation.