A NOTE ON SCROLLS OF SMALLEST EMBEDDED CODIMENSION

Takao FUJITA

arXiv:alg-geom/9507003v2 11 Jul 1995

§0. Introduction

The situation considered in this note is as follows. Let M be an algebraic submanifold of $\mathbb{P}^N$ with $n = \dim M$. M is said to be a scroll over S if there is a surjective morphism $\pi : M \to S$ such that every fiber $F_x = \pi^{-1}(x)$ over $x \in S$ is a linear subspace in $\mathbb{P}^N$ of dimension $r - 1 = n - s$, where $s = \dim S$. This is equivalent to saying that $M \cong \mathbb{P}_S(\mathcal{E})$ for some vector bundle $\mathcal{E}$ of rank $r = n - s + 1$ and the tautological bundle $H(\mathcal{E})$ is the hyperplane section bundle of M.

We have $b_i(M) = b_i(\mathbb{P}^N)$ for $i \leq 2n - N$ by Barth-Lefschetz Theorem, hence $N \geq 2n - 1$ for scrolls, since otherwise $1 + b_2(S) = b_2(M) = b_2(\mathbb{P}^N) = 1$, contradiction. Thus we want to study scrolls such that $N = 2n - 1$.

The case $s = 1$ was studied by Lanteri-Turrini [**LT**], who showed that M is the Segre scroll over $\mathbb{P}^1$ in this case; namely $S \cong \mathbb{P}^1$, $\mathcal{E} \cong \mathcal{O}(1)^{\oplus n}$, $M \cong \mathbb{P}^1 \times \mathbb{P}^{n-1}$ and $M \subset \mathbb{P}^N$ is the Segre embedding. In this paper we are interested mainly in the case $s = 2$.

This problem was studied by Ottaviani [**Ot**] and Beltrametti-Schneider-Sommese [**BSS**], [**BS**] in case $n = 3$ and by Ionescu-Toma [**IT**] for general n. Their results are as follows.

Theorem. (cf. [**Ot**]). *Let $M \subset \mathbb{P}^5$ be a three-dimensional scroll over a surface S. Then one of the following conditions is satisfied.*
(1) (Segre scroll) $S \cong \mathbb{P}^2$, $\mathcal{E} \cong \mathcal{O}(1)^{\oplus 2}$, $d = \deg M = 3$.
(2) (Bordiga scroll) $S \cong \mathbb{P}^2$, $c_1(\mathcal{E}) = \mathcal{O}(4)$, $c_2(\mathcal{E}) = 10$ *and* $d = 6$.
(3) (Palatini scroll) *S is isomorphic to a cubic surface,* $c_1(\mathcal{E}) = \mathcal{O}_S(2)$, $c_2(\mathcal{E}) = 5$ *and* $d = 7$.
(4) (K3 scroll) *S is a K3-surface obtained as a linear section of the Grassman variety parametrizing lines in $\mathbb{P}^5$ embedded by Plücker, and $\mathcal{E}$ is the restriction of the tautological vector bundle.* $c_2(\mathcal{E}) = 5$ *and* $d = 9$.

All these four cases actually occur.

Theorem. (cf. [**IT**]). *Let $M \subset \mathbb{P}^N$ be a scroll over a surface S with $N = 2n - 1$, $n > 3$ and let $\mathcal{E}$ and $d = \deg M$ be as above. Then, M is one of the following types:*
(S_n) $S \cong \mathbb{P}^2$, $\mathcal{E} \cong \mathcal{O}(1)^{\oplus(n-1)}$, $M \cong \mathbb{P}^2 \times \mathbb{P}^{n-2}$, $d = n(n-1)/2$.
(B_n) $S \cong \mathbb{P}^2$, $c_1(\mathcal{E}) = \mathcal{O}(n+1)$, $c_2(\mathcal{E}) = (n+1)(n+2)/2$, $d = n(n+1)/2$.
(M_n) *S is a K3 surface,* $c_1(\mathcal{E})^2 = 2n^2 - 4$, $c_2(\mathcal{E}) = n^2 - 4$, $d = n^2$.
($?_n$) *S is a surface of general type.*

The existence of scrolls of the above type (S_n) is classical and is due to Segre. The case (B_n) is shown to exist for every n in [**IT**]. The existence of the type (M_n) is proved by Mukai ([**Mu**]). On the other hand, no example of type ($?_n$) is found: [**IT**] suspect rather that there is no such scroll. Any way, such scrolls must satisfy several numerical relations among their invariants. Here I propose a conjecture concerning further classifications derived from these relations, which is verified for $n \leq 11$ by hand and for $n \leq 1100$ by a computer. In particular, a scroll of the type ($?_n$) does not exist unless $n = 6, 10, 11, 12, 16, 18, 20, 24, 30, \cdots$. On the other hand, no example is known and the existence problem is unsettled for these n (and also for

Mathematical tools used here are almost the same as [**IT**], but we review them here for the convenience of the reader. When I started this study, I was not aware of this paper [**IT**]. I would like to express my hearty thanks to Professors Ottaviani who informed me of the result in [**IT**]. I thank Professor Mukai, who communicated to me the existence of scrolls of the type (M_n). I also thank Professors Lanteri and Schneider for their cooperations in e-mail correspondence during the preparation of this paper.

§1. Computing Chern classes

(1.1) Throughout this paper let M be a scroll in $\mathbb{P}^{2n-1}$ over S as in §0. Let $\mathcal{E}$ be the vector bundle on S of rank $r = n - s + 1$, $s = \dim S$, such that $M \cong \mathbb{P}_S(\mathcal{E})$ and the tautological bundle $H(\mathcal{E})$ is the hyperplane section bundle $\mathcal{O}_M(1)$, which will be denoted simply by H from now on. We put $d = \deg M = H^n\{M\}$.

(1.2) *Fact. Via the ring homomorphism $\pi^* : H^\cdot(S) \to H^\cdot(M)$, the cohomology ring of M becomes a free $H^\cdot(S)$-module generated by $1, h, h^2, \cdots, h^{r-1}$, where $h = c_1(H) \in H^2(M)$. Moreover $\sum_{i=0}^{r}(-h)^i e_{r-i} = 0$ in $H^\cdot(M)$, where $e_j = \pi^* c_j(\mathcal{E})$.*

This is well known for general vector bundle $\mathcal{E}$ of rank r. Thus, the ring structure of $H^\cdot(M)$ is determined by $H^\cdot(S)$ and Chern classes of $\mathcal{E}$.

(1.3) *Fact. Put $s_i(\mathcal{E}) = \pi_* h^{r-1+i} \in H^{2i}(S)$ and $s(\mathcal{E}) = \sum_{i=0}^{\infty} s_i(\mathcal{E}) \in H^\cdot(S)$. Then $s(\mathcal{E})c(\mathcal{E}^\vee) = 1$, where $c(\mathcal{E}^\vee)$ is the total Chern class of the dual bundle $\mathcal{E}^\vee$ of $\mathcal{E}$.*

This is also standard. $s_i(\mathcal{E})$ is called the i-th Segre class of $\mathcal{E}$ and $s(\mathcal{E})$ is called the total Sege class of $\mathcal{E}$. Moreover the following formulas are well known:

$$\begin{aligned} s_1(\mathcal{E}) &= c_1(\mathcal{E}), \\ s_2(\mathcal{E}) &= c_1(\mathcal{E})^2 - c_2(\mathcal{E}), \\ &\cdots \end{aligned}$$

(1.4) **Corollary.** *For any vector bundle E on X with $\mathrm{rank}E = r$ and for any line bundle L on X with $c_1(L) = \ell$, we have $s_i(E \otimes L) = \sum_j \binom{r-1+i}{j} s_{i-j}(E)\ell^j$.*

Indeed, $s_i(E \otimes L) = \pi_*((h+\ell)^{(r-1+i)}) = \sum_j \binom{r-1+i}{j} \pi_* h^{r-1+i-j} \cdot \ell^j$ for $\pi : \mathbb{P}(E) \to X$.

(1.5) Let $\pi : M = \mathbb{P}(\mathcal{E}) \to S$ be the projection. Let $\mathcal{A}$ be the kernel of the natural surjection $\pi^*\mathcal{E} \to \mathcal{O}[H]$. Then we have an exact sequence $0 \to \mathcal{O} \to \mathcal{E}^\vee \otimes H \to \mathcal{A}^\vee \otimes H \to 0$. This is identified with the relative Euler sequence, and so we have the exact sequence $0 \to \mathcal{A}^\vee \otimes H \to \Theta_M \to \pi^*\Theta_S \to 0$, where Θ_X denotes the tangent bundle of X.

From these exact sequences we obtain the following relation $c(\Theta_M) = \pi^* c(\Theta_S) c(\pi^*\mathcal{E}^\vee \otimes H)$ of total Chern classes.

(1.6) Let $\mathcal{N}$ be the normal bundle of M in $\mathbb{P}^{2n-1}$ and let Θ be the restriction of the tangent bundle of $\mathbb{P}^{2n-1}$ to M. Then we have $c(\Theta_M)c(\mathcal{N}) = c(\Theta) = (1+h)^{2n}$, where $h = c_1(H)$.

Combining (1.5), we get $c(\mathcal{N}) = (1+h)^{2n} c(\pi^*\mathcal{E}^\vee \otimes H)^{-1} \pi^* c(\Theta_S)^{-1} = (1+h)^{2n} s(\pi^*\mathcal{E} \otimes [-H]) \pi^* s(\Omega_S)$.

(1.7) In the following computation, $\pi^*\alpha$ will be denoted simply by α for $\alpha \in H^\cdot(S)$.

By (1.4), we have $s(\mathcal{E}_M \otimes [-H]) = \sum_{i=0}^{\infty} (\sum_{j=0}^{r-1+i} \binom{r-1+i}{j} s_{i-j}(\mathcal{E})(-h)^j)$. Therefore the component of $(1+h)^{2n} s(\mathcal{E}_M \otimes [-H])$ of degree $2k$ is $\sum_{i,j} \binom{2n}{k-i}\binom{r-1+i}{j}(-1)^j h^{k-i+j} s_{i-j}(\mathcal{E})$ $= \sum_\ell (\sum_j \binom{2n}{k-j-\ell}\binom{r-1+j+\ell}{j}(-1)^j) h^{k-\ell} s_\ell(\mathcal{E}) = \sum_\ell \binom{2n-r-\ell}{k-\ell} h^{k-\ell} s_\ell(\mathcal{E})$ by the following

(1.8) *Claim.* $\sum_{(a,b)|a+b=c} \binom{m-1+a}{a}(-1)^a \binom{p}{b} = \binom{p-m}{c}$.

To see this, use the Taylor expansion $(1+T)^{-m} = \sum_{a=0}^{\infty} \binom{m-1+a}{a}(-T)^a$ and compute the

(1.9) Combining (1.6) and (1.7), we get a formula for $c(\mathcal{N})$. On the other hand, we have $c_n(\mathcal{N}) = 0$ and $c_{n-1}(\mathcal{N}) = dh^{n-1}$ since M is of codimension $n-1$ in $\mathbb{P}^N$. This gives non-trivial relations among Chern classes of $\mathcal{E}$ and Θ_S. In the next section we analyse them precisely in case $s = \dim S = 2$ and $r = n-1$.

§2. Over a surface

From now on we assume $s = \dim S = 2$ and set $e_j = c_j(\mathcal{E})$, $\gamma_i = c_i(\Theta_S)$.

(2.1) From (1.6) and (1.7) we obtain
$c_{n-1}(\mathcal{N}) = \binom{n+1}{n-1}h^{n-1} + \binom{n}{n-2}h^{n-2}e_1 + \binom{n-1}{n-3}h^{n-3}(e_1^2 - e_2) - \binom{n+1}{n-2}h^{n-2}\gamma_1 - \binom{n}{n-3}h^{n-3}e_1\gamma_1 + \binom{n+1}{n-3}h^{n-3}(\gamma_1^2 - \gamma_2)$.
This is dh^{n-1}, while $h^{n-1} - h^{n-2}e_1 + h^{n-3}e_2 = 0$ is the unique relation in $H^{\cdot}(M)$ (cf. (1.2)). Hence, substituting $h^{n-1} = h^{n-2}e_1 - h^{n-3}e_2$ and comparing the coefficients of h^{n-2} and h^{n-3}, we get the following relations in $H^{\cdot}(S)$:

(i) $$(n^2 - d)e_1 = (n+1)n(n-1)\gamma_1/6, \qquad \text{and}$$

(ii) $$\tfrac{(n-1)(n-2)}{2}e_1^2 + (d - n^2 + n - 1)e_2 - \tfrac{n(n-1)(n-2)}{6}e_1\gamma_1 + \tfrac{(n+1)n(n-1)(n-2)}{24}(\gamma_1^2 - \gamma_2) = 0.$$

Next from $c_n(\mathcal{N}) = 0$ we get

(iii) $$3e_1^2 - 2e_2 - ne_1\gamma_1 + \tfrac{(n+1)(n-1)}{6}(\gamma_1^2 - \gamma_2) = 0.$$

(2.2) Eliminating γ_2 from (ii) and (iii), we get $-3(n^2-4)e_1^2 + 6(2d - n^2 - 2)e_2 + n(n-2)(n+2)e_1\gamma_1 = 0$. Eliminating γ_1 further using (i) and noting $d = (e_1^2 - e_2)\{S\}$, we obtain

(*) $$\{2(q+2)d - q(q+5)\}\{(q+2)(q-4)d - (q+2)^2e_2 + (q-1)(q-4)\} = -q(q-1)(q-4)(q+5)$$

for $q = n^2$.

(2.3) Thus, $2(q+2)d - q(q+5)$ is one of the finitely many divisors of the right hand side, so there are only finitely many numerical possibilities for d and e_2. Note that they are positive integers since $\mathcal{E}$ is ample.

(2.4) For each (d, e_2) satisfying (*), we examine the relation (i). By the result [**IT**], it suffices to consider the case where the canonical bundle of S is ample, or equivalently, $d > q = n^2$. Let $6(d-n^2)/(n+1)n(n-1) = a/b$ for coprime integers a, b. Then $ae_1 \sim -b\gamma_1$, so $e_1 \sim bA$ and $K_S \sim -\gamma_1 \sim aA$ for some ample line bundle A on S. Therefore

$$e_1^2 = d + e_2 \quad \text{is divided by} \quad b^2.$$

(2.5) In the above case, we have $\gamma_1^2 = a^2(d+e_2)/b^2$ and $e_1\gamma_1 = -a(d+e_2)/b$. Using (iii), we solve γ_2. The result has to satisfy the Noether relation:

$$\gamma_1^2 + \gamma_2 \equiv 0 \text{ modulo } 12.$$

(2.6) It is easy to produce a computer programm to enumerate pairs (n, d) satisfying the numerical conditions (2.2), (2.4) and (2.5). In view of the result of our experiment, we make the following

Conjecture. *Any pair (n, d) with $n > 3$ as above is one of the following types:*
(1) *$n \equiv 0, 2, 6, 12$ or 16 modulo 18 and $d = q(q+5)/6$ for $q = n^2$. Moreover $e_2 = (q-4)(q+3)/6$, $K_S \sim ne_1$, $K_S^2 = q(q^2+2q-6)/3$, $\gamma_2 = e(S) = (q^3+8q^2+24q+36)/3$ and $\chi(\mathcal{O}_S) =$*

(2) $n = 10$ *and* $d = 595$. *Moreover* $e_2 = 561$, $K_S \sim 3e_1$, $K_S^2 = 10404$, $e(S) = 12648$ *and* $\chi(\mathcal{O}_S) = 1921$.
(3) $n = 11$ *and* $d = 231$. *Moreover* $e_2 = 221$, $2K_S \sim e_1$, $K_S^2 = 113$, $e(S) = 283$ *and* $\chi(\mathcal{O}_S) = 33$.

(2.7) *Remarks.*
(1) The above conjecture is verified for $n \leq 11$ by my hand, and for $n \leq 1100$ by a personal computer. In particular, the case with smallest n is $(n, d) = (6, 246)$. [**IT**] apparently claims that such a case with $n = 6$ can be ruled out by "divisibility manipulations", but I cannot see how this can be done.
(2) The above condition for n of the type (1) is equivalent to $q \equiv 0$ or 4 modulo 18.
(3) It is perhaps a delicate problem whether scrolls of the type $(?_n)$ exist or not for a pair (n, d) in (2.6). I find no example at present. To settle the problem, we need some more geometric observations. I feel that the case (2.6.3) might be of particular interest, since this is the unique case with odd n and moreover the invariants are relatively small.
(4) The sectional genus $g = g(M, \mathcal{O}(1))$ can be computed by using the relation $2g - 2 = (K_S + e_1)e_1\{S\}$, and g is bounded by the Castelnuovo inequality. But it turns out that no pair (n, d) in (2.6) is thus ruled out.

References

[**BSS**] C. Beltrametti, M. Schneider and A. J. Sommese, *Threefolds of degree 9 and 10 in* $\mathbb{P}^5$, Math. Ann. **288** (1990), 613–644.

[**BS**] C. Beltrametti and A. J. Sommese, *New properties of special varieties arising from adjunction theory*, J. Math. Soc. Japan **43** (1991), 381–412.

[**FL**] W. Fulton and R. Lazarsfeld, *Positive polynomials for ample vector bundles*, Ann. of Math. **118** (1983), 35–60.

[**IT**] P. Ionescu and M.Toma, *Boundedness for some special families of embedded manifolds*, Contemporary Mathematics **162** (1994), 215–225.

[**LT**] A. Lanteri and C. Turrini, *Some formulas concerning nonsingular algebraic varieties embedded in some ambient variety*, Atti Accad. Sci. Torino **116** (1982), 463–474.

[**Mu**] N. Mukai, *K3 scrolls of dimension n in a projective space of dimension* $2n - 1$, letter, dated June 2, 1995.

[**Ott**] G. Ottaviani, *On 3-folds in* $\mathbb{P}^5$ *which are scrolls*, Annali d. Scuola Norm. Sup. Pisa **19** (1992), 451–471.

Takao FUJITA
Department of Mathematics
Tokyo Institute of Technology
Oh-okayama, Meguro, Tokyo
152 Japan
e-mail:fujita@math.titech.ac.jp