arXiv:alg-geom/9608034v2 30 Aug 1996

The ring of global sections of multiples of a line bundle on a toric variety

To appear in Proceedings of the AMS

E. Javier Elizondo*

July 14, 2011

Abstract

In this article we prove that for any complete toric variety, and for any Cartier divisor, the ring of global sections of multiples of the line bundle associated to the divisor is finitely generated.

1 Introduction

Let us start by stating the result and a few consequences. Let X be an algebraic variety which is complete over a field K, and let D be any effective Cartier divisor on X. We denote by $\mathcal{O}(nD)$ the line bundle associated to nD, with $n \geq 0$. It is a very interesting problem to know if the ring

$$R := \bigoplus_{n \geq 0} H^0\left(X, \mathcal{O}(nD)\right) \tag{1}$$

is a finitely generated K-algebra. It was O. Zariski who worked out the case of algebraic surfaces in order to solve the Riemann-Roch problem, and gave examples where R is not finitely generated, see [Zar62]. In this article we

*Supported in part by grant CONACYT 3936-E

prove that the ring R is finitely generated when X is a complete toric variety (perhaps singular).

The ring R appears in many interesting problems, for example, if X is a nonsingular projective variety, then the finite generation of R implies that the series

$$\sum_{n\geq 0} \dim H^0(X, \mathcal{O}(nD))\, t^n \tag{2}$$

is rational. It was asked by S. D. Cutkosky and V. Srinivas [CS93] if this series is rational when D is a nef divisor. Our result gives trivially a positive answer in the case of a complete toric variety, and D any effective Cartier divisor.

Furthermore, the rationality of the series also allows us to compute, at least theoretically, the dimension of $H^0(X, \mathcal{O}(nD))$ in terms of n. This is the Riemann-Roch problem, see for example [Zar62] and [CS93]. Observe that these dimensions are not given by the Riemann-Roch theorem since the variety X can be singular, and the line bundle $\mathcal{O}(nD)$ is not necessarily generated by its global sections. We would like to make a final remark. The ring R is a subring of the ring $S = \bigoplus_{D\geq 0} H^0(X, \mathcal{O}(D))$, which is finitely generated in the case of toric varieties. In fact, D. Cox proves in [Cox95] that the ring S is a polynomial ring graded by the the monoid of effective divisors classes in the Chow group $A_{k-1}(X)$ of X, where k is the dimension of X.

2 The ring of global sections

In this section we prove that the ring R, which was defined by equation (1), is a finitely generated K-algebra. Throughout this section X means a complete toric variety over a field K, and D a Cartier divisor in X. The construction of the cone C_R, in the proof of the Theorem, is a well know construction, see for example [Bat93]. We start by recalling an important lemma that will be

use in the proof of the theorem.

Lemma 2.1 (Gordan). *If σ is a strongly convex rational polyhedral cone, and $\sigma^\vee$ its dual, then $S_\sigma = \sigma^\vee \cap M$ is a finitely generated semigroup.*

Now, we are ready for the main result.

Theorem 2.2. *Let X be a complete toric variety, perhaps singular, and let D be an Cartier divisor in X. Then the ring*

$$R := \bigoplus_{n \geq 0} H^0(X, \mathcal{O}(nD))$$

is finitely generated as a K-algebra.

Proof. We first observe that we can consider our divisor D to be T-invariant. For the following sequence is exact, see [Dan78, page 116]


$$0 \longrightarrow M \longrightarrow \mathrm{Div_T}\ X \longrightarrow \mathrm{Pic}\,(X) \longrightarrow 0$$

where $\mathrm{Div_T}\ X$ is the group of T-invariant Cartier divisors and $M := \mathrm{Hom}_{\mathbb{Z}}(N, \mathbb{Z})$ is the dual lattice of $N \cong \mathbb{Z}^k$. Then we can write D as $D = \sum_{i=1}^{s} a_i D_i$ with $\{D_i\}$ the set of invariant divisors. We consider the convex rational polyhedron P_{nD} in $M_{\mathbb{Q}} := M \bigotimes_{\mathbb{Z}} \mathbb{Q}$ defined as

$$P_{nD} = \{u \in M_{\mathbb{Q}} \mid < u, v_i > \geq -na_i \text{ for all } i\}$$

where v_i is the the first element in the lattice appearing in the divisor D_i. We know that generators for the space $H^0(X, \mathcal{O}(nD))$ are given by the elements of $P_{nD} \cap M$, which is a finite set because X is a complete variety, this also implies that P_D is a rational convex polytope. Let us embed the k-dimensional $\mathbb{Q}$-vector space $M_{\mathbb{Q}}$ into the (k+1)-dimensional $\mathbb{Q}$-vector

space $M_{\mathbb{Q}}^{k+1} := \mathbb{Z}^{k+1} \bigotimes_{\mathbb{Z}} \mathbb{Q}$ as the hyperplane with equation $x_{k+1} = 1$, where $(x_1, \ldots, x_{k+1})$ are coordinates for $M_{\mathbb{Q}}^{k+1}$. Denote by C_R the $(k+1)$-dimensional cone in $M_{\mathbb{Q}}^{k+1}$ generated by the rays starting at the origin and passing through the vertices of P_D (see figure below).



The cone C_R

It follows from the definition that $P_{nD} = nP_D$, and this implies that the intersection of the hyperplane $x_{k+1} = n$ with the cone C_R is just the polyhedron P_{nD}. In other words, the cone C_R is the cone associated to the ring R, in the sense that any element of the ring R is a finite linear combination of integral points of $C_R \cap \mathbb{Z}^{k+1}$. The theorem follows since the semigroup $C_R \cap \mathbb{Z}^{k+1}$ is finitely generated by Lemma 2.1 (Gordan). □

Acknowledgement:. I thank very much V. Srinivas for suggesting me this problem and for many wonderful and fruitful conversations. I also would like to thank A. King for bringing to my attention the article of D. Cox, and to the referee for pointing out the article of V. Batyrev.

References

[Bat93] Victor V. Batyrev. Variations of the mixed hodge structure of affine hypersurfaces in algebraic tori. *Duke Math. J.*, 69(2):349–409, Feb 1993.

[Cox95] David A. Cox. The homogeneous coordinate ring of a toric variety. *J. Algebraic Geom.*, 4(3):17–50, 1995.

[CS93] S. D. Cutkosky and V. Srinivas. On a problem of Zariski on dimensions of linerar systems. *Ann. of Math.*, 137:531–559, 1993.

[Dan78] V. I. Danilov. The geometry of toric varieties. *Russian Math. Surveys*, 33(2):97–154, 1978.

[Zar62] O. Zariski. The theorem of Riemann-Roch for high multiples of an effective divisor on an algebraic surface. *Ann. of Math.*, 76:560–616, 1962.

Instituto de Matemáticas
Ciudad Universitaria, UNAM
México D.F. 04510
México

e-mail: javier@math.unam.mx