arXiv:alg-geom/9711005v1 4 Nov 1997

# SOME FUNDAMENTAL PROBLEMS ON REAL-ANALYTIC SETS

TOHSUKE URABE

## 1. Introduction

The category of real-analytic sets and real-analytic maps is the most important category in application. However, in spite of efforts by F. Bruhat, H. Cartan, H. Whitney et al., the basic theory of real-analytic category does not yet seem to be well-developed. In this article I would like to point out several basic problems. (Urabe [6].)

Let $Q$ denote a *connected* real-analytic *smooth* manifold, and $X \subset Q$ a subset of $Q$.

## 2. When is $X$ real-analytic?

We have two kinds of definitions.

**Definition 2.1.** We say that $X$ is real-analytic, if for every point $x \in X$ there are an open set $U$ with $x \in U \subset Q$ and a finite number of real-analytic functions $f_1, f_2, \ldots, f_k$ on $U$ such that $X \cap U$ coincides with the set of common zero-points of $f_1, f_2, \ldots, f_k$.

Under Definition 2.1 $X$ is not necessarily a closed subset of $Q$. It is locally closed, or there is an open subset $U$ of $Q$ containing $X$ such that $X$ is a closed subset of $U$.

**Definition 2.2.** We say that $X$ is real-analytic, if for every point $x \in Q$ there are an open set $U$ with $x \in U \subset Q$ and a finite number of real-analytic functions $f_1, f_2, \ldots, f_k$ on $U$ such that $X \cap U$ coincides with the set of common zero-points of $f_1, f_2, \ldots, f_k$.

Note that only the phrase "$x \in X$" has been replaced by "$x \in Q$". Under Definition 2.2 $X$ is closed in $Q$.

I myself prefer former Definition 2.1, because it gives us flexibility of the theory and it reduces non-essential descriptions in related subjects, in which real-analytic sets in Definition 2.1 naturally appear.

## 3. When is a point $x \in X$ smooth?

**Definition 3.1.** A real-analytic germ $(X, x)$ of sets is *smooth*, if $(X, x)$ is real-analytically isomorphic to $(\mathbf{R}^d, 0)$ where $d = \dim(X, x)$ and $0 \in \mathbf{R}^d$ is a point.

1991 *Mathematics Subject Classification.* Primary 32C07; Secondary 26-02.
*Key words and phrases.* real-analytic, smooth, irreducible.

**Example 3.2.** Let $k = 4$ or $5$, $X = \{(x, y) \in \mathbf{R}^2 \mid x^3 = y^k\}$ and $0 = (0, 0)$. Then, the germ $(X, 0)$ is *not* smooth under above Definition 3.1. However, $(X, 0)$ is isomorphic to $(\mathbf{R}, 0)$ in the category of germs of sets and maps of class $C^1$. Note that $|y|^{4/3}$ and $y|y|^{2/3}$ are $C^1$-functions on $\mathbf{R}$.

## 4. When can we define the global concept of irreducible decompositions?

**Definition 4.1.** We say that a real-analytic subset $X \subset Q$ is *global*, if we have a set $F$ of real-analytic functions defined globally on $Q$ such that $X$ coincides with the set of common zero-points of members in $F$.

**Definition 4.2.** We say that a global real-analytic subset $X \subset Q$ is *reducible*, if $X = Y \cup Z$ for some global real-analytic subsets $Y, Z \subset Q$ with $X \neq Y$ and $X \neq Z$. We say that $X$ is *irreducible*, if it is not reducible.

**Theorem 4.3.** *Any global irreducible real-analytic subset in $Q$ is connected.*

**Definition 4.4.** Let $X$ be a global real-analytic subset of $Q$. A family $S$ of global irreducible real-analytic subsets in $Q$ is called the *irreducible decomposition* of $X$, if the union of members in $S$ coincides with $X$ and if any two members in $S$ have no inclusion relation.

In *compact* case we can show the following:

**Theorem 4.5.** *Any compact global real-analytic set $X$ in $Q$ has a unique irreducible decomposition.*

**Example 4.6.** Consider an *irreducible* polynomial $f(x, y) = y^2 - x^2(x - 1)$ for example. The set $X = \{(x, y) \in \mathbf{R}^2 \mid f(x, y) = 0\}$ of zero-points of $f(x, y)$ is a global real-analytic set. Drawing the picture of $X$, we see that it is not connected. Thus, by Theorem 4.3 $X$ is reducible.

**Example 4.7.** Let $m$ be a positive integer and $X_m = \{(x, y) \in \mathbf{R}^2 \mid x^2 + y^2 = m^2\}$. The union $X = \bigcup_{m=1}^{\infty} X_m$ is a non-compact global real-analytic subset with countably many connected components.

**Problem 4.8.** Give generalization of Theorem 4.5 for real-analytic subsets which are not necessarily closed.

## References

[1] François Bruhat and Henri Cartan, *Sur la structure de sous-ensemble analytique-réelles*, C. R. Acad. Sci. Paris **244** (1957), 988–990.

[2] ———, *Sur les composantes irréducibles d'un sous-ensemble analytique-réelles*, C. R. Acad. Sci. Paris **244** (1957), 1123–1126.

[3] François Bruhat and Hassler Whitney, *Quelques propriétés fondamentales des ensembles analytiques-réels*, Comm. Math. Helv. **33** (1959), 132–160.
[4] Henri Cartan, *Variétés analytiques réelles et variétés analytiques complexes*, Bull. Soc. math. France **85** (1957), 77–99.
[5] Hans Grauert, *On Levi's problem and the imbedding of real-analytic manifolds*, Annals of Math. **68** (1958), 460–472.
[6] Tohsuke Urabe, *The Gauss map and the dual variety of real-analytic submanifolds in a sphere or in hyperbolic space*, preprint (1995).
[7] Hassler Whitney, *Elementary structure of real algebraic varieties*, Annals of Math. **66** (1957), no. 3, 545–556.

Department of Mathematics, Tokyo Metropolitan University, Minami-Ohsawa 1-1, Hachioji-shi, Tokyo, 192-03, Japan
*E-mail address*: urabe@comp.metro-u.ac.jp