

Version Date : 21th of July 1999

The broad Brillouin doublets and central peak of KTaO_3

E. Farhi^{a,b}, A.K. Tagantsev^c, B. Hehlen^a, R. Currat^b, L.A. Boatner^d, and E. Courtens^a

^a *Laboratoire des Verres, Université de Montpellier 2, F-34095 Montpellier, France*

^b *Institut Laue Langevin, BP 156, 38042 Grenoble Cedex 9, France*

^c *Laboratoire de Céramique, EPFL, CH-1015 Lausanne, Switzerland*

^d *Solid State Division, Oak Ridge Nat. Lab., Oak Ridge, TN 37831-6056, USA*

Abstract

The incipient ferroelectric KTaO_3 presents low- T Brillouin spectra anomalies, *e.g.* a broad central peak (CP), and some additional Brillouin doublets (BD), whose origin is interpreted in terms of phonon-density fluctuation processes. A parameterisation from new extensive high-resolution neutron-scattering measurements is used to show that hydrodynamic second sound from high damping (compared to BD frequency) TA phonons may exist in the crystal. Furthermore, low damping thermal phonons may scatter light through two-phonon difference processes and appear on the Brillouin spectra either as a sharp or a broader BD, depending on the phonon damping and group velocity. The comparison between computed anisotropies and experimental measurements favours the second process.

keywords : Computer modelling, Perovskite compounds, Phonons, Spectroscopy.

Corresponding Author : E. Farhi

Address : Institut Laue-Langevin, BP 156, F-38042 Grenoble Cedex 9, France

Fax : 33 (0)4 76 48 39 06

E-mail Address : farhi@ill.fr

PACS numbers : 77.84 Dy, 63.20.-e, 62.20.Dc, 78.70.Nx, 66.90.+r, 78.35.+c

Physica B Preprint

Incipient ferroelectrics SrTiO₃ or KTaO₃ are perovskite type crystals that remain paraelectric down to the lowest temperature, due to zero-point quantum fluctuations [1]. The dielectric constant saturates at a high value as the soft transverse optic (TO) zone-centre frequency stabilises to a low finite value. Since the case of SrTiO₃ is complicated by the structural transition at $T_a \sim 105$ [2], we study KTaO₃ which remains in the cubic phase down to the lowest temperatures.

Experimental evidence of anomalous behaviour in pure KTaO₃ when cooling into the quantum paraelectric (QPE) regime are numerous. In Particular, the low energy Brillouin spectra present unusual features. Hehlen *et al.* [3] found a new sound-like Brillouin doublet (BD) both in SrTiO₃ and KTaO₃ in addition to the quasi-elastic broad central peak (CP) first reported by Lyons and Fleury [4]. It was interpreted as the signature of second sound [3, 5]. Our purpose here is to investigate the origin of CP and BD in the frame of phonon-density fluctuation processes.

The anisotropy of anomalous Brillouin contributions was measured in KTaO₃ Ultra High Purity crystals along the three high symmetry axes by means of Brillouin light-backscattering, with a momentum transfer $|\vec{q}| = 3.55 \cdot 10^{-3}$ rlu. The observations of Hehlen *et al.* [2, 3] are confirmed for the $\vec{q} // \langle 001 \rangle$ symmetry axis. A sharp BD appears upon the broad CP [4] below $T \lesssim 22$ K. Its associated velocity is $v_{BD} \sim 1100$ m/s. In the $\langle 110 \rangle$ direction, the BD and CP contributions are also observed, as well as an additional broad BD (BBD) for temperatures between 30 and 100 K. While the CP and BBD contributions are also seen along the $\langle 111 \rangle$ direction, no sharp BD could be measured for temperatures as low as 5 K [6, 7].

In order to look for an explanation of CP, BD, and BBD contributions in terms of phonon-density fluctuation processes, we parameterised the low-energy phonon dispersion surfaces $\omega(\vec{q})$ below 100 K within a simple phenomenological quasi-harmonic anisotropic model due to Vaks [7, 8]. The precise determination of the necessary parameters required additional new inelastic neutron-scattering experiments to be carried out. Taking into account the 4-dimensional nature of the instrumental resolution ellipsoid and the anisotropy of the dispersion surfaces, an effective energy uncertainty as low as 0.1 meV was achieved. The resulting dispersion curves and their parameterisation are presented in Fig. 1 along the principal symmetry axes, showing the well-known TO-TA coupling [9]. The transverse phonon dispersion surfaces show deep 'valleys' along the $\langle 001 \rangle$ directions and for $\vec{q} // \langle 110 \rangle$ with $\vec{e} // \langle 001 \rangle$. Some 'uphill' regions are found elsewhere, specially for $\vec{q} // \langle 111 \rangle$,

and $\vec{q} // \langle 110 \rangle$ with $\vec{e} // \langle 1\bar{1}0 \rangle$. This anisotropic character was confirmed by off-principal-axis measurements.

We now analyse the origin of the anomalous Brillouin contributions in terms of classical phonon-kinetics processes. Second sound (2S) is the collective mode of phonon-density fluctuations induced by multiphonon processes [5, 10]. It may propagate at a frequency ω_{2S} in a pure crystal when the 'window' condition, $\Gamma_R < \omega_{2S} < \Gamma_N$ (1), is fulfilled. Here, Γ_N and Γ_R are the normal and resistive damping coefficients respectively. In KTaO_3 , Eq. (1) is satisfied since the proximity of the optic and acoustic modes enhances the relaxation processes. The computation of 2S velocity using Vaks model and the thermalized part of the phonon spectrum gives $v_{2S} = 1100 \pm 200$ m/s, in excellent agreement with measurements of BD velocity. However, it seems difficult to explain the directional dependence of BD in this framework.

When Eq. (1) is not satisfied, low-damping phonons might interact *via* two-phonon difference scattering (TPDS). Following Wehner and Klein [6, 10], we evaluated the corresponding Brillouin spectra for different anisotropic dispersion surfaces. It was found that TPDS from low- Γ_N *valley*-TA with small group velocity v_g appears as a sharp BD on low- T spectra, whereas a BBD is obtained by TPDS at slightly higher T from thermalized TA phonons around *uphill* regions, where Γ_N and v_g are higher. Finally, a broad CP originates from high- Γ_N TO phonon TPDS. Thus, the intrinsically anisotropic TPDS spectral shapes are in agreement with Brillouin experiments.

As a conclusion, we reported new Brillouin spectroscopy and high-resolution inelastic neutron-scattering results in pure KTaO_3 . On the basis of those experimental data, a simple phenomenological parameterisation of the phonon dispersion surfaces was used to study two different scenarios for the appearance of additional Brillouin contributions : two-phonon difference scattering processes (TPDS) and second sound (2S). The comparison between computed anisotropies and experimental measurements seems to favour the TPDS explanation.

References

- [1] K.A. Müller and H. Burkard, Phys. Rev. B 19 (1979) 3593.

- [2] E. Courtens, B. Hehlen, G. Coddens, B. Hennion, *Physica B : Cond. Matt.* 219 (1996) 577.
- [3] B. Hehlen, A-L. Pérou, E. Courtens, and R. Vacher, *Phys. Rev. Lett.* 75 (1995) 2416.
- [4] K.B. Lyons and P.A. Fleury, *Phys. Rev. Lett.* 37 (1976) 161.
- [5] V.L. Gurevich and A.K. Tagantsev, *Zh. Eksp. Teor. Fiz.* 94 (1988) 370 [*Sov. Phys. JETP* 67 (1988) 206].
- [6] E. Farhi, Ph.D. Thesis, Université Montpellier II (1998).
- [7] E. Farhi, A.K. Tagantsev, R. Currat, B. Hehlen, L.A. Boatner, and E. Courtens, submitted to *Phys. Rev. B* (1999).
- [8] V.G. Vaks, *Zh. Eksp. Teor. Fiz.* 54 (1968) 910 [*Sov. Phys. JETP* 27 (1968) 486].
- [9] C.H. Perry, R. Currat, H. Buhay, R.M. Migoni, W.G. Stirling, and J.D. Axe, *Phys. Rev. B* 39 (1989) 8666.
- [10] R.K. Wehner and R. Klein, *Physica (Utrecht)* 62 (1972) 161.

Figure 1: Low- T dispersion curves and parameterisation of KTaO_3 low energy phonons. The mean energy uncertainty is 0.1 meV. The LA mode is for $\vec{q} = \langle 0 \ 0 \ 2+\xi \rangle$.

