arXiv:cond-mat/9504121v1 30 Apr 1995

Theory of Brownian motion of Massive Particle in a Space with Curvature and Torsion and Crystals with Defects

H. Kleinert and S.V. Shabanov*
Institut für Theoretische Physik, Freie Universität Berlin, Arnimallee 14, D-14195 Berlin
(August 13, 2013)

We develop a theory of Brownian motion of a massive particle, including the effects of inertia (Kramers' problem), in spaces with curvature and torsion. This is done by invoking the recently discovered generalized equivalence principle, according to which the equations of motion of a point particle in such spaces can be obtained from the Newton equation in euclidean space by means of a nonholonomic mapping. By this principle, the known Langevin equation in euclidean space goes over into the correct Langevin equation in the Cartan space. This, in turn, serves to derive the Kubo and Fokker-Planck equations satisfied by the particle distribution as a function of time in such a space.

The theory can be applied to classical diffusion processes in crystals with defects.

1. The classical equation of a massive point particle in a thermal environment reads

$$m\ddot{x}_t^i = f_t^i + \bar{f}_t^i, \tag{1}$$

where f_t^i is an arbitrary time-dependent external force and $\bar{f}_t^i$ a stochastic force caused by the thermal fluctuations (we use subscripts for the time variable). The stochastic force may be modeled by a bath of harmonic oscillators of all frequencies ω as $\bar{f}_t^i = \int_0^\infty d\omega \lambda_\omega \dot{X}_{\omega t}^i$. The oscillator coordinates satisfy the equations of motion $\ddot{X}_{\omega t}^i + \omega^2 X_{\omega t}^i = \lambda_\omega \dot{x}_t^i$, the right-hand side arising from the back-reaction of the particle. Solving the latter equation with respect to $X_{\omega t}^i$ we find $\bar{f}_t^i$ as a functional of $\dot{x}_t^i$. Assuming an equal coupling of all oscillators, $\lambda_\omega \equiv \sqrt{2\gamma/\pi}$, we obtain the stochastic differential equation

$$m\ddot{x}_t^i + \gamma \dot{x}_t^i - f_t^i = \eta_t^i \ , \tag{2}$$

where η_t^i is called the *noise variable*. This has the decomposition

$$\eta_t^i = \int_0^\infty d\omega \lambda_\omega \left[\dot{X}_\omega^i \cos\omega t - X_\omega^i \sin\omega t \right]. \tag{3}$$

For any given phase space distribution $\rho^B = \rho^B(\dot{X}, X)$ of initial oscillator velocities $\dot{X}_\omega^i$ and coordinates X_ω^i, Eq.(2) becomes a classical Langevin equation with noise averages being defined as mean-values with respect to the distribution ρ^B. In thermal equilibrium, ρ^B follows the Boltzmann law $\sim \exp(-H^B/kT)$, where the bath Hamiltonian is a sum of the oscillator energies $H^B = \int d\omega (\dot{X}_\omega^2 + \omega^2 X_\omega^2)/2$. In this case, the noise is Gaussian and completely specified by its vanishing expectation and its two-point correlation function:

$$\langle \eta_t^i \rangle = 0 \ , \quad \langle \eta_t^i \eta_{t'}^j \rangle = 6\gamma k T \delta^{ij} \delta(t - t') \ . \tag{4}$$

2. By a nonholonomic mapping $dx^i = e^i{}_\mu(q) dq^\mu$ [1], [2] the euclidean space is carried into an arbitrary metric-affine space with curvature and torsion [3]. The geometry in this space is defined by the metric $g_{\mu\nu} = e^i{}_\mu e^i{}_\nu$ and the affine connection $\Gamma_{\mu\nu}{}^\lambda = e_i{}^\lambda \partial_\mu e^i{}_\nu$ [2]. The torsion is carried by the antisymmetric part of the affine connection, $S_{\mu\nu}{}^\lambda = (\Gamma_{\mu\nu}{}^\lambda - \Gamma_{\nu\mu}{}^\lambda)/2$, which is a tensor, in contrast to $\Gamma_{\mu\nu}{}^\lambda$ itself. The curvature tensor arises from noncommuting derivatives of $e^i{}_\lambda$: $R_{\mu\nu\lambda}{}^\kappa = e_i{}^\kappa (\partial_\mu \partial_\nu - \partial_\mu \partial_\nu) e^i{}_\lambda$. Thus, in a curved space, the nonholonomic quantities $e_i{}^\mu(q)$ fail to satisfy Schwarz' criterion and are no proper functions [1].

*Alexander von Humboldt fellow, on leave from Laboratory of Theoretical Physics, JINR, P.O.Box 79, Moscow, Russia

Freie Universität Berlin Preprint FUB–HEP/95–7
Email: kleinert@einstein.physik.fu-berlin.de; shabanov@axp3.physik.fu-berlin.de
World Wide Web homepage: http://www.physik.fu-berlin.de/∼kleinert

Recently it was found that the nonholonomic mapping $dx^i = e^i{}_\mu(q)dq^\mu$ transforms euclidean equations of motion, Schroedinger equations, and path integrals correctly into spaces with curvature and torsion [4]. This universal mapping rule was called *quantum equivalence principle* (QEP) [5].

Applying this principle to the equation of motion (2), we obtain the analogous equation in the general metric-affine space:

$$m\left(\ddot{q}^\mu_t + \Gamma_{\sigma\nu}{}^\mu \dot{q}^\sigma_t \dot{q}^\nu_t\right) + \gamma \dot{q}^\mu_t - f^\mu_t = e_i{}^\mu(q_t)\eta^i_t \ . \tag{5}$$

To obtain physical consequences, we must find equations in which the nonholonomic quantities $e_i{}^\mu(q_t)$ are eliminated in favor of the well-defined geometrical objects, the metric $g_{\mu\nu}(q_t)$ and the affine connection $\Gamma_{\mu\nu}{}^\lambda(q_t)$. This is possible by deriving from (5) Kubo's stochastic Liouville equation and the Fokker-Planck equation.

For this purpose we rewrite the Langevin equation as a system of two first-order differential equations

$$\dot{q}^\mu_t = \frac{1}{m} g^{\mu\nu}(q_t) p^t_\nu \ ; \tag{6}$$

$$\begin{aligned} \dot{p}^t_\mu &= -\frac{1}{m}\left(\Gamma^{\sigma\nu}{}_\mu - g^{\sigma\lambda} g^{\nu\alpha} \partial_\lambda g_{\mu\alpha}\right) p^t_\sigma p^t_\nu - \frac{\gamma}{m} p^t_\mu + f^\mu_t + e^\mu_i \eta^i_t \\ &\equiv -F^t_\mu + e^i_\mu \eta^i_t \ . \end{aligned} \tag{7}$$

The last equation defines the total apparent force F^t_μ, where $f^t_\mu = g_{\mu\nu}(q_t) f^\nu_t$. We mark the time dependence also by a superscript to avoid a pile-up of subscripts. Note that although the time derivative of the momentum $\dot{p}^t_\mu$ and the force F^t_μ do not transform as vectors under general coordinate transformations, Eq. (7) is nevertheless covariant under these transformations, just as the Langevin equation (5).

At each time t, the system following Eqs. (6) and (7) is in a microscopic state with the distribution function $\delta(p - p^t)\delta(q - q_t)$ (omitting the spatial indices μ). This can be thought of as a conditional distribution function determining the distribution of p and q for a given initial distribution function $\delta(p - p^0)\delta(q - q_0)$. If the initial values of p^t and q_t are distributed with the probability density $\rho(p^0, q_0)$, the distribution function at any later time t can be found by an average over the initial distribution

$$\rho^\eta_t(p, q) = \int dp^0 dq_0 \rho(p^0, q_0) \delta(p - p^t)\delta(q - q_t) \ , \tag{8}$$

where $q_t = q_t(p^0, q_0)$ and $p^t = p^t(p^0, q_0)$ are solutions of the system (6), (7) with the initial conditions $p^{t=0} = p^0$ and $q_{t=0} = q_0$. Upon taking a time derivative of (8) and making use of both (6) and (7) and of the identity

$$\frac{d}{dt}\delta(z - z_t) = \dot{z}_t \frac{\partial}{\partial z_t}\delta(z - z_t) = -\frac{\partial}{\partial z}\left[\dot{z}_t \delta(z - z_t)\right] = -\frac{\partial}{\partial z}\left[V(z)\delta(z - z_t)\right] \ , \tag{9}$$

where a dynamical variable z_t (any of p^t_μ or q^μ_t) obeys the equation of motion $\dot{z}_t = V(z_t)$, we find Kubo's stochastic Liouville equation

$$\partial_t \rho^\eta_t = \hat{L}(\eta_t)\rho^\eta_t \ , \tag{10}$$

with $\hat{L}(\eta_t)$ being the noise-dependent Liouville operator

$$\hat{L}(\eta_t) = -\frac{\partial}{\partial q^\mu} \circ g^{\mu\nu}(q)\frac{p_\nu}{m} + \frac{\partial}{\partial p_\mu} \circ \left[F_\mu(p, q) - e^i_\mu \eta^i_t\right] \ . \tag{11}$$

The symbol $\circ$ emphasizes the fact that the differential operators in front of it act on the product of the function behind it with ρ^η_t in Eq. (10).

It is straightforward to verify the invariance of the Liouville operator and, hence, of Kubo's equation with respect to general coordinate transformations.

A solution of Kubo's stochastic equation (10) is a noise-dependent distribution function which determines the probability to find a particle in an infinitesimal volume dq by

$$dP_t(q) = dq \int dp \langle \rho^\eta_t(p, q)\rangle \equiv dq \int dp \rho_t(p, q) \ . \tag{12}$$

It follows then from (8) that $\int dP_t(q) = \int dP_0(q) = 1$, i.e. the temporal evolution of the probability distribution described by Kubo's equation (10) is unitary.

The locality of the noise correlator (4) enables us to derive a Fokker-Planck equation governing the temporal evolution of the noise-averaged distribution $\rho_t(p,q)$. For this let us first calculate the average $\varphi_t^i = \langle \eta_t^i \delta(p-p^t)\delta(q-q_t)\rangle$. Generalizing the partial integration formula $\int e^{-a\eta^2/2}\eta f(\eta)d\eta = a^{-1}\int e^{-a\eta^2/2}f'(\eta) = -a^{-1}\int \partial_\eta e^{-a\eta^2/2}f(\eta)$ to any Gaussian noise, we find

$$\varphi_t^i = \int\limits_{-\infty}^{\infty} dt' \langle \eta_t^i \eta_{t'}^j \rangle \left\langle \frac{\delta}{\delta \eta_{t'}^j} \delta(p-p^t)\delta(q-q_t) \right\rangle$$
$$= -6\gamma kT \left\langle \left(\frac{\partial}{\partial p_\mu} \circ \frac{\delta p_\mu^t}{\delta \eta_t^i} + \frac{\partial}{\partial q^\mu} \circ \frac{\delta q_t^\mu}{\delta \eta_t^i} \right) \delta(p-p^t)\delta(q-q_t) \right\rangle \ , \tag{13}$$

where we have used the explicit form of the two-point noise correlation function (4) and the identity (9). To calculate the variational derivatives of dynamical variables with respect to noise in (13), we formally integrate (6) and (7):

$$p_\mu^t = p_\mu^0 + \int\limits_0^t dt' \left(-F_\mu^{t'} + e_\mu^i \eta_{t'}^i \right) \ ; \tag{14}$$

$$q_t^\mu = q_0^\mu + \frac{1}{m}\int\limits_0^t dt' g^{\mu\nu}(q_{t'}) p_\nu^{t'} \ . \tag{15}$$

Taking the variational derivative of (14), we obtain

$$\frac{\delta p_\mu^t}{\delta \eta_{t'}^i} = -\int\limits_{t'}^t dt'' \left[\frac{\delta F_\mu^{t''}}{\delta \eta_{t'}^i} - \eta_{t''}^j \frac{\delta e_\mu^j(q_{t''})}{\delta \eta_{t'}^i} \right] + \int\limits_{t'}^t dt'' e_i^\mu(q_{t''})\delta(t''-t') \ . \tag{16}$$

The Langevin equation (7) is causal which implies that $p_{t''}^\mu$ or $q_{t''}^\mu$ depend on $\eta_{t'}^i$ only for $t'' > t'$. This has been used to restrict the integration range in (16). The first integral in the left-hand side of (16) tends to zero as t' approaches t, whereas the second integral is equal to $e_i^\mu(q_{t'})\Theta(t-t')$ and, hence, is not uniquely determined, since it involves the value of Heaviside function at zero argument, $\Theta(0)$, when $t = t'$. Therefore a regularization is needed. We replace the delta-function in the correlator (4) by a smooth would-be delta-function of width ϵ, $\delta_\epsilon(t-t')$, which satisfies $\int_{-\infty}^{\infty} dt \delta_\epsilon(t) = 1$ and $\delta_\epsilon(t) = \delta_\epsilon(-t)$. It is easily seen from (3) that such a regularization can be achieved by retaining a weak dependence on ω in the coupling constant λ_ω [1]. With this regularization, we have to replace $\delta p_\mu^t/\delta\eta_t^i$ in (13) by

$$\int\limits_{-\infty}^{\infty} dt' \delta_\epsilon(t-t') \frac{\delta p_\mu^t}{\delta \eta_{t'}^i} = \int\limits_{-\infty}^{\infty} dt' \delta_\epsilon(t-t') e_\mu^i(q_{t'})\Theta(t-t') = \frac{1}{2} e_\mu^i(q_t) \ , \tag{17}$$

where we have dropped the contribution of the first integral of (16) since it vanishes for $\epsilon \to 0$.

Considering analogously $\delta q_t^\mu/\delta\eta_{t'}^i$, we conclude that the latter variational derivative vanishes as t' approaches t. Averaging φ_t^i with respect to the initial distribution $\rho(p_0,q_0)$ we find the following relation

$$\int dp_0 dq_0 \rho(p_0,q_0)\varphi_t^i = \langle \eta_t^i \rho_t^\eta \rangle = -3\gamma kT e_\mu^i \frac{\partial}{\partial p_\mu}\rho_t \ . \tag{18}$$

Taking the noise average of Kubo's stochastic equation (10) and making use of (18), we end up with the Fokker-Planck equation associated with the Langevin equation (5):

$$\partial_t \rho_t = \hat{L}_T \rho_t \ , \tag{19}$$

$$\hat{L}_T = -\frac{\partial}{\partial q^\mu} \circ g^{\mu\nu}\frac{p_\nu}{m} + \frac{\partial}{\partial p_\mu} \circ \left[F_\mu(p,q) + 3\gamma kT g^{\mu\nu}\frac{\partial}{\partial p_\nu} \right] \ , \tag{20}$$

---

[1] This physical regularization corresponds to the Stratonovich treatment of stochastic integrals [6], p.55. A symmetrically smoothed correlator appears also due to quantum effects (the fluctuation-dissipation theorem).

Thus, we have eliminated the nonholonomic mapping functions in favor of the well-defined affine connection and metric. Integrating (19) over all phase space we arrive at the probability conservation law: $d/dt \int dpdq\rho_t = 0$, where we have used the fact of vanishing surface integrals occurring upon the integration of the right-hand side of (19).

In the overdamped limit, we may drop the mass term in (5) and derive analogously an equation for the q-space probability distribution $D_t(q)$, defined by $dP_t(q) = dq\sqrt{g}D_t(q)$,

$$\begin{aligned}\partial_t D_t &= \left[\frac{3kT}{\sqrt{g}}\partial_\mu \circ e_i^\mu \partial_\nu \circ e_i^\nu \sqrt{g} - \frac{1}{\gamma\sqrt{g}}\partial_\mu \circ f^\mu \sqrt{g}\right] D_t \\ &= \left[\frac{3kT}{\sqrt{g}}\partial_\mu \circ g^{\mu\nu}\sqrt{g}(\partial_\nu + 2S_\nu - \kappa f_\nu)\right] D_t \ , \end{aligned} \tag{21}$$

where $\kappa = (3kT\gamma)^{-1}$, $S_\nu = S_{\nu\mu}{}^\mu$ and $g = \det g_{\mu\nu}$. Since S_ν and f_ν are vectors, the operator in the brackets is invariant under general coordinate transformations. In the absence of torsion, the limiting equation (21) agrees with a result in Ref. [7].

3. Since defects in crystals can be described geometrically by a nonvanishing torsion and curvature in a metric-affine space in which distances are measured by counting atomic steps [1], the Fokker-Planck equation (19) can be applied to the classical diffusion of interstitial atoms in crystals [8].

It must be realized, however, that the approximation is necessarily very crude. The reason for this is the twofold role played by the phonon bath. First, it provides the system with a random driving force which in our equations has been simulated by the model noise η_t^i. Second, it influences the hopping probabilities of the interstitials. The latter effect has not been accounted for in our initial Eq. (5). In crystals, the hopping probability depends on elastic distortions which correspond to time-independent general coordinate transformations in the above theory. The Langevin equation (5) is, however, invariant with respect to these transformations. A proper treatment should take into account the fact that elastic distortions are also produced by the off-shell phonons comprising the noise.

Another field of applications should be positrons or muons in a crystal at low temperatures below a few or about 80 Kelvin, respectively. These particles are bound in the lowest atomic orbits, so that their wave functions follow the tight-binding approximation. Spending most of their time near the atomic sites, they live effectively in the above crystal geometry.

It is worth noting that in the absence of external forces, the overdamped equation (21) agrees with an equation recently studied in Ref. [9].

Acknowledgment:
The authors thank R. Schmitz for sending them reprints of his work with R. Bausch and L.A. Turski, and for subsequent discussions. They also acknowledge a useful discussion with H. Grabert.

---

[1] Such mappings are standard in the theory of defects:
K. Kondo, in: Proc. 2nd Japan Nat. Congr. Applied Mechanics, Tokio, 1952
B.A. Bilby, R. Bullough and E. Smith, Proc. R. Soc. London **A231**, 263 (1955);
E. Kröner, in: *Physics of defects*, Les Houches summer school XXXV, North-Holland, Amsterdam 1981;
H. Kleinert, *Gauge fields in condensed matter*, Vol. II, World Scientific, Singapore, 1989;

[2] J.A. Schouten, *Ricci– Calculus*, Springer-Verlag, Berlin, 1954.

[3] E. Cartan, *On Manifolds with an affine connection and the theory of general relativity*, Bibliopolis, Italy, 1986;

[4] H. Kleinert, *Path Integrals in Quantum Mechanics, Statistics and Polymer Physics,*
World Scientific Publishing Co., Singapore 1990 and 1995 (second extended edition).

[5] Note that the usual action principle of classical mechanics requires a modification: The variation at the upper end of the time interval is no longer zero but equal to the Burgers vector of the "dislocations" enclosed by the path variation. See P. Fiziev and H. Kleinert, Berlin preprints, hep-th/9503074 and hep-th/9503074. The trajectories in a space with torsion are autoparallels rather than geodesics.

[6] H.Risken, *The Fokker-Planck equation*, Springer-Verlag, Heidelberg, 1984.

[7] J.Zinn-Justin, *Quantum Field Theory and Critical Phenomena*, Clarendon Press, Oxford, 1990.

[8] For a discussion of the physics of such processes see K. Schröder, *Theory of Diffusion Controlled Reactions of Point Defects in Metals*, Springer Tracts in Mod. Phys., **87**, 171 (1980); J.W. Haus and K.W. Kehr, Phys. Rep. **150**, 264 (1987).

[9] R. Bausch, R. Schmitz, and L.A. Turski, Phys.Rev.Lett. **73** (1994) 2382.