

Commuting Difference Operators with Polynomial Eigenfunctions

June 1993
funct-an/9306002

J.F. van Diejen*

Department of Mathematics and Computer Science, University of Amsterdam,
Plantage Muidergracht 24, 1018 TV Amsterdam, The Netherlands.

Abstract. We present explicit generators $\hat{D}_1, \dots, \hat{D}_n$ of an algebra of commuting difference operators in n variables with trigonometric coefficients. The algebra depends, apart from two scale factors, on five parameters. The operators are simultaneously diagonalized by Koornwinder's multivariable generalization of the Askey-Wilson polynomials. For special values of the parameters and via limit transitions, one obtains difference operators for the Macdonald polynomials that are associated with (admissible pairs of) the classical root systems: A_{n-1} , B_n , C_n , D_n and BC_n . By sending the step size of the differences to zero, the difference operators reduce to known hypergeometric differential operators. This limit corresponds to sending $q \rightarrow 1$; the eigenfunctions reduce to the multivariable Jacobi polynomials of Heckman and Opdam. Physically the algebra can be interpreted as an integrable quantum system that generalizes the (trigonometric) Calogero-Moser systems related to classical root systems.

1991 Mathematics Subject Classification: 39A70, 47A75, 33D45, 33D80, 81Q99.

Key words & Phrases: commuting difference operators, (joint) eigenfunctions, multivariable q-polynomials, classical root systems, integrable quantum systems.

Contents

1	Introduction	1
2	Introducing the Difference Operators	3
2.1	The Operator $\hat{D}_r$	3
2.2	Combinatorial Structure and Parameters	4
2.3	$g = 0$: Reduction to Rank $n = 1$	4
3	Simultaneous Diagonalization	5
3.1	Trigonometric Polynomials	5
3.2	Triangularity	6
3.3	The Spectrum	7
3.4	Symmetry	8
3.5	Diagonalization and Commutativity	10
4	$\beta \rightarrow 0$: The Transition to BC_n-type Hypergeometric PDO's	11
4.1	Eigenfunctions	11
4.2	Eigenvalues	12
4.3	Operators	13

*E-mail address: jand@fwi.uva.nl

5	Special Cases Related to Classical
5.1	Preliminaries
5.2	The Root System A_{n-1}
5.3	The Root Systems B_n , C_n and BC_n
5.4	The Root System D_n

Appendix A: Cancellation of Poles

Appendix B: Two Combinatorial Lemmas

Appendix C: $\hat{D}(\mathcal{A}^W) = 0 \implies \hat{D} = 0$

References

1 Introduction

Over the past few years, progress has been made in more than one variable. It has turned out that orthogonal polynomials depending on only one variable. Such generalizations can be viewed as root system R , there exist associated families of polynomials. The number of variables coincides with the rank n of the root system. For Jacobi polynomials related to root systems of type A , see [HO, H1]. Recently, a more elementary approach was given in [H2]. For $R = BC_1$ the Heckman-Opdam polynomials reduce to Jacobi polynomials in one variable. In a recent paper [H3] we introduced q-versions of the Heckman-Opdam polynomials. In a sequel of these results and [M3] for lectures devoted to the subject, we then Macdonald's polynomials coincide with the q-Jacobi polynomials. Information on continuous q-Jacobi polynomials and their properties. Results pertaining to the root system BC_n have been obtained. We find BC_n -type multivariable versions of the Heckman-Opdam polynomials. The one-variable case is recovered by specializing $q \rightarrow 1$.

A crucial ingredient in the construction of the difference operator of which the polynomials are eigenfunctions is the Jacobi polynomials. This operator is a second-order partial differential operator which is named hypergeometric differential operator. For the polynomials the relevant operator is an adjoint transformation turns the hypergeometric differential operator into its adjoint with respect to Lebesgue measure. Upon the latter PDO as being the Hamiltonian of a quantum system of dimension one. Such quantum systems have been studied in the literature; they are known as (generalized) Calogero-Moser systems. The systems studied in [C] and [Su1, Su2] are of the type studied in the survey paper [OP2] for the generalization of the corresponding classical systems has already been given (arbitrary R).

It has been shown by Heckman and Opdam that the difference operator is but one member of an algebra of commuting difference operators.

polynomials as their joint eigenfunctions [HO, H1, O1, O2]. This algebra is generated by n independent PDO's. This state of affairs can be expressed by saying that the corresponding (generalized) Calogero-Moser system is quantum integrable. For $R = A_{n-1}$, Liouville integrability of the classical system, i.e. the existence of n independent integrals in involution, was already proved by Moser [Mo] using a Lax pair formulation. For a partial generalization of this result to the other classical (i.e. non-exceptional) root systems, see [OP1] and [I].

Just as the hypergeometric PDO, Macdonald's difference operator for $R = A_{n-1}$ is related to certain known quantum systems of n particles. The systems of interest were originally introduced as a relativistic generalization of the Calogero-Moser systems [RS] (classical) and [R1] (quantum). (See [R2] for a survey and connections with certain soliton PDE's and exactly solvable quantum field theories). The relativistic generalization of the Calogero-Moser system is (quantum) integrable too: explicit formulas representing n independent commuting integrals are presented in [RS, R1]. At the quantum level these integrals are $\Lambda\Delta\mathcal{O}$'s, which are related to Macdonald's A_{n-1} -type difference operators E_{ω_r} via a similarity transformation [K1]. The operators E_{ω_r} (associated with the fundamental weights ω_r) generate an algebra of commuting $\Lambda\Delta\mathcal{O}$'s which have Macdonald's A_{n-1} -type polynomials as joint eigenfunctions.

Also for root systems other than A_{n-1} one expects that there exist algebras of commuting difference operators that are simultaneously diagonalized by the Macdonald polynomials. The purpose of the present paper is to introduce such difference operators for the Koornwinder polynomials. We will show that via limit transitions and/or specialization of parameters, this also leads to the corresponding AΔO's for those families of Macdonald polynomials that are connected with (admissible pairs of) the classical series: A_{n-1} , B_n , C_n , D_n and BC_n ; the Koornwinder polynomials then reduce to the latter Macdonald polynomials. By sending the step size of the differences to zero (this corresponds to the limit $q \rightarrow 1$) our AΔO's go over in PDO's. Thus, we recover the commuting hypergeometric PDO's associated with the classical root systems as a limit case.

Our difference operators constitute a new integrable quantum system of n particles in dimension one. In this paper, however, we will not pay much attention to this interpretation of the ADO's; instead we will emphasize the connection with orthogonal polynomials. In a forthcoming paper the author intends to return to the question of integrability of these and related n -particle systems, at the level of both quantum and classical mechanics [D]. In particular, possible generalization to integrable systems consisting of commuting difference operators with elliptic functions as coefficients will be discussed in [D].

Before outlining the contents of this paper in more detail, let us mention two more connections of interest. For special values of the parameters (namely those corresponding to root multiplicities), the system of hypergeometric PDO's coincides with the radial reduction of the algebra of invariant differential operators on certain symmetric spaces G/K [OP2, HO]. It seems natural to ask oneself the question whether, for special values of the parameters, our system of difference operators can be seen in some way as radial reduction of certain AΔO's connected with quantum homogeneous spaces. Recent results on the quantum group interpretation of Macdonald's A_{n-1} -type polynomials [K3, N] indeed seem to point in this direction. However, apart from this special case no relations of this kind are known to the author.

In 5.3 and 5.4 we show how the Macdon root systems can be obtained from the Koo of the parameters. In contrast to our ac contained, this involves various concepts f account more accessible by collecting some skipped at first reading and referred back t

Recently, Cherednik introduced commuting difference operators connected with Knizh-

2 Introducing the Difference Operators

In this section the operators $\hat{D}_r$, $r = 1, \dots, n$, are introduced and their combinatorial structure is discussed.

2.1 The Operator $\hat{D}_r$

In order to write down our difference operators we first introduce some notation. Let $v_a(z)$ and $v_b(z)$ be the following trigonometric functions:

$$v_a(z) = \frac{\sin \alpha(\mu + z)}{\sin \alpha z}, \quad (2.1)$$

$$v_b(z) = \frac{\sin \alpha(\mu_0 + z)}{\sin \alpha z} \frac{\cos \alpha(\mu_1 + z)}{\cos \alpha z} \frac{\sin \alpha(\mu'_0 + \gamma + z)}{\sin \alpha(\gamma + z)} \frac{\cos \alpha(\mu'_1 + \gamma + z)}{\cos \alpha(\gamma + z)}, \quad (2.2)$$

with α , γ and μ , μ_δ , μ'_δ , ($\delta = 0, 1$) complex parameters. For later purposes, it is convenient to parametrize γ according to:

$$\gamma \equiv i\beta/2. \quad (2.3)$$

We form the following multivariable functions using v_a and v_b as elementary constituents:

$$\begin{aligned} V_{\varepsilon J; K} &\equiv \prod_{j \in J} v_b(\varepsilon_j x_j) \prod_{\substack{j, j' \in J \\ j < j'}} v_a(\varepsilon_j x_j + \varepsilon_{j'} x_{j'}) v_a(\varepsilon_j x_j + \varepsilon_{j'} x_{j'} + 2\gamma) \\ &\times \prod_{\substack{j \in J \\ k \in K}} v_a(\varepsilon_j x_j + x_k) v_a(\varepsilon_j x_j - x_k), \end{aligned} \quad (2.4)$$

with

$$J, K \subset \{1, \dots, n\}, \quad J \cap K = \emptyset; \quad \varepsilon_j \in \{+1, -1\}. \quad (2.5)$$

The variables $x_1, \dots, x_n$ are assumed to be real. The function $V_{\varepsilon J; K}$ depends on the index sets J , K and on a collection of prescribed signs ε_j , $j \in J$; it serves as a building block from which the coefficients of $\hat{D}_r$ are constructed. The AΔO's read explicitly

$$\hat{D}_r \equiv \sum_{\substack{J \subset \{1, \dots, n\}, |J|=r \\ \varepsilon_j = \pm 1, j \in J}} \sum_{\substack{\emptyset \subsetneq J_1 \subsetneq \dots \subsetneq J_s = J \\ 1 \leq s \leq r}} (-1)^{s-1} \prod_{1 \leq s' \leq s} V_{\varepsilon(J_{s'} \setminus J_{s'-1}); J_{s'}^c} \left(e^{-\beta \hat{\theta}_{\varepsilon J_1}} - 1 \right), \quad (2.6)$$

$$r = 1, \dots, n,$$

with $J_0 \equiv \emptyset$ and

$$\hat{\theta}_{\varepsilon J} \equiv \sum_{j \in J} \varepsilon_j \hat{\theta}_j, \quad \hat{\theta}_j \equiv \frac{1}{i} \frac{\partial}{\partial x_j}. \quad (2.7)$$

Remarks i. $|J|$ denotes the cardinality of J and J^c is the complement set $\{1, \dots, n\} \setminus J$.
ii. The first summation in Eq. (2.6) is over all index sets $J \subset \{1, \dots, n\}$ with cardinality r and over all flippings of the signs $\varepsilon_j \in \{+1, -1\}$, $j \in J$; the second summation is over all strictly increasing sequences of subsets in J :

$$\emptyset \subsetneq J_1 \subsetneq J_2 \subsetneq \dots \subsetneq J_{s-1} \subsetneq J_s = J, \quad 1 \leq s \leq |J|. \quad (2.8)$$

iii. The exponential $\exp(-\beta \hat{\theta}_j)$ acts on a function of the variables $x_1, \dots, x_n$, as a (complex) shift:

$$(e^{-\beta \hat{\theta}_j} f)(x_1, \dots, x_n) = f(x_1, \dots, x_n + \beta x_j).$$

Hence, $\hat{D}_r$ is indeed an analytic difference operator.
iv. The ‘hats’ in Eqs. (2.6), (2.7) are used to denote operators rather than ordinary (complex) functions of the variables.
v. In the simplest case, i.e. for $r = 1$, Eq. (2.6) reads

$$\hat{D}_1 = \sum_{\substack{1 \leq j \leq n \\ \varepsilon = \pm 1}} v_b(\varepsilon x_j) \prod_{k \neq j} v_a(\varepsilon x_j + x_k) v_a(\varepsilon x_j - x_k).$$

vi. It is clear that the operator $\hat{D}_r$ is invariant under the shift $x_j \rightarrow x_j + \pi/(2\alpha)$, $j = 1, \dots, n$. Furthermore, the AΔO's are periodic with period $\pi/(2\alpha)$: a simultaneous shift of the variables x_j by $\pi/(2\alpha)$ leaves the parameters:

$$x_j \rightarrow x_j + \pi/(2\alpha), \quad j = 1, \dots, n.$$

(see Eq. (2.4) and Eqs. (2.1), (2.2)).

For some purposes it is more convenient to write such expressions read

$$\hat{D}_r = \sum_{\substack{J \subset \{1, \dots, n\}, |J|=r \\ \varepsilon_j = \pm 1, j \in J}} \sum_{\substack{\emptyset \subsetneq J_0 \subsetneq \dots \subsetneq J_s = J \\ 0 \leq s \leq r}} (-1)^{s-1} \prod_{1 \leq s' \leq s} V_{\varepsilon(J_{s'} \setminus J_{s'-1}); J_{s'}^c} \left(e^{-\beta \hat{\theta}_{\varepsilon J_1}} - 1 \right),$$

(with $J_{-1} \equiv \emptyset$) and

$$\hat{D}_r = \sum_{0 \leq s \leq r} \sum_{\substack{J \subset \{1, \dots, n\}, |J|=r \\ \varepsilon_j = \pm 1, j \in J}} (-1)^{s-1} \prod_{1 \leq s' \leq s} V_{\varepsilon(J_{s'} \setminus J_{s'-1}); J_{s'}^c} \left(e^{-\beta \hat{\theta}_{\varepsilon J_1}} - 1 \right),$$

with

$$\begin{aligned} W_{I,p} &\equiv \sum_{\substack{1 \leq q \leq p \\ \varepsilon_i = \pm 1, i \in I}} (-1)^q \sum_{\substack{\emptyset \subsetneq I_1 \subsetneq \dots \subsetneq I_q \subsetneq I \\ |I_q|=p}} (-1)^{q-1} \prod_{1 \leq s' \leq q} V_{\varepsilon(I_{s'} \setminus I_{s'-1}); I_{s'}^c} \left(e^{-\beta \hat{\theta}_{\varepsilon I_1}} - 1 \right), \\ W_{I,0} &\equiv 1. \end{aligned}$$

In Eq. (2.12), $\hat{D}_r$ is written in terms of the coefficients $W_{I,p}$. Accordingly, J_0 in Eq. (2.12) is the empty set. The fact that the coefficients of the translation operators $\hat{D}_r$ which does not commute with the translation operators

Note. From a physical point of view, one can interpret the operators $\hat{D}_r$ as describing a particle quantum system in dimension one. The parameters μ , μ_δ , μ'_δ , ($\delta = 0, 1$) and $v_a(\pm x_j \pm x_k)$ in the coefficients are interpretable for the interaction between the particles in the field.

2.2 Combinatorial Structure and Parameters

The increment sets $J_1, J_2 \setminus J_1, \dots, J_s \setminus J_{s-1}$ of the increasing sequence (2.8) form the *blocks* of a partition of J ; the second summation in (2.6) amounts to a sum over all ordered blocks. By breaking up $V_{\varepsilon J; K}$ (Eq. (2.4)) into three parts

$$V_{\varepsilon J; K} = V_{\varepsilon J}^1 V_{\varepsilon J}^2 V_{\varepsilon J; K}^3 \quad (2.15)$$

with

$$V_{\varepsilon J}^1 \equiv \prod_{j \in J} v_b(\varepsilon_j x_j), \quad (2.16)$$

$$V_{\varepsilon J}^2 \equiv \prod_{\substack{j, j' \in J \\ j < j'}} v_a(\varepsilon_j x_j + \varepsilon_{j'} x_{j'}) v_a(\varepsilon_j x_j + \varepsilon_{j'} x_{j'} + 2\gamma), \quad (2.17)$$

$$V_{\varepsilon J; K}^3 \equiv \prod_{\substack{j \in J \\ k \in K}} v_a(\varepsilon_j x_j + x_k) v_a(\varepsilon_j x_j - x_k), \quad (2.18)$$

one can rewrite $\hat{D}_r$ (Eq. (2.6)) as

$$\begin{aligned} \hat{D}_r = & \sum_{\substack{J \subset \{1, \dots, n\}, |J|=r \\ \varepsilon_j = \pm 1, j \in J}} \left\{ V_{\varepsilon J}^1 V_{\varepsilon J; J^c}^3 \right. \\ & \times \sum_{\substack{\emptyset \subsetneq J_1 \subsetneq \dots \subsetneq J_s = J \\ 1 \leq s \leq r}} (-1)^{s-1} \prod_{1 \leq s' \leq s} V_{\varepsilon(J_{s'} \setminus J_{s'-1})}^2 V_{\varepsilon(J_{s'} \setminus J_{s'-1}); J \setminus J_{s'}}^3 \left(e^{-\beta \hat{\theta}_{\varepsilon J_1}} - 1 \right) \Big\}. \end{aligned} \quad (2.19)$$

Eqs. (2.6), (2.12) and (2.13) are more compact than (2.19), but the latter has the virtue that different parts of the coefficient can be controlled independently. The index set J in Eq. (2.19) will be referred to as the *cell*. The first block J_1 determines the translator; this part of the cell will be called the *nucleus*. Notice that: *i.* $V_{\varepsilon J}^1 V_{\varepsilon J; J^c}^3$ depends on the cell J but not on its subdivision in blocks; *ii.* the product over $V_{\varepsilon(J_{s'} \setminus J_{s'-1})}^2$ depends on the partition of J , but not on the order of the blocks; *iii.* the product over $V_{\varepsilon(J_{s'} \setminus J_{s'-1}); J \setminus J_{s'}}^3$ depends both on the blocks and on their order.

The parameters α and β are scale factors; α determines the period of the trigonometric functions and β the complex shift of the translation operators $\exp(\pm \beta \hat{\theta}_j)$. Both parameters will be taken positive. The parameters μ, μ_δ and $\mu_{\delta'}$ determine the relative ‘weight’ of v_a and v_b in the coefficients of the $\Lambda\Delta\mathcal{O}$. For instance, $v_a \equiv 1$ for $\mu = 0$; therefore, v_a may be omitted for $\mu = 0$. The parameters μ, μ_δ and $\mu_{\delta'}$ will be assumed to be non-negative imaginary:

$$\begin{aligned} \mu &\equiv i\beta g, & \mu_\delta &\equiv i\beta g_\delta, & \mu_{\delta'} &\equiv i\beta g'_\delta, & (\delta = 0, 1) \\ \alpha, \beta &> 0; & g, g_\delta, g'_\delta &\geq 0. \end{aligned} \quad (2.20)$$

Notice that the above restrictions on the parameters guarantee that: *i.* $\exp(\pm \beta \hat{\theta}_j)$ yields a purely imaginary shift; *ii.* the commuting part of the coefficient, viz. $W_{J^c, r-s}$ (cf. Eqs. (2.13), (2.14)), is real because $v_c(z) = v_c(-z)$ ($c = a, b$) for z real.

If one picks $\alpha = 1/2$, then $\hat{D}_1$ (Eq. (2.10)) is a finite constant, with Koornwinder’s difference operators. The parameters used in [K4] are related to ours

$$q = e^{-\beta},$$

$$a = e^{-\beta g_0}, \quad b = -e^{-\beta g_1}, \quad c = e^{-\beta g_2}.$$

Note. The parameters g, g_δ and g'_δ can be taken as constants that determine the strengths of the interactions. Setting $g = 0$ (i.e. $\mu = 0$) yields a system without an external field.

2.3 $g = 0$: Reduction to Random Walks

By setting $g = 0$, the combinatorial structure of the coefficients in Eq. (2.19) no longer depends on g ($0 \Rightarrow V_{\varepsilon J}^2 = V_{\varepsilon J; K}^3 = 1$). It will be shown that $\hat{D}_r$ is an elementary symmetric function of the following

$$\hat{D}_1(x_j) \equiv \sum_{\varepsilon = \pm 1} v_b(\varepsilon x_j) \left(e^{-\beta \hat{\theta}_{\varepsilon J_1}} - 1 \right)$$

(For $n = 1$, $\hat{D}_1$ (2.10) coincides with $\hat{D}_1(x_1)$.)

By summation of all terms in $\hat{D}_r$ which contain $J_1 = I$, Eq. (2.19) reduces to

$$\hat{D}_r = \sum_{\substack{J \subset \{1, \dots, n\}, |J|=r \\ \varepsilon_j = \pm 1, j \in J}} v_b(\varepsilon J_1) \left(e^{-\beta \hat{\theta}_{\varepsilon J_1}} - 1 \right)$$

where

$$c_0 \equiv 1, \quad c_p \equiv \sum_{1 \leq j_1 < \dots < j_p \leq n} v_b(\varepsilon j_1) \left(e^{-\beta \hat{\theta}_{\varepsilon j_1}} - 1 \right)$$

with

$$N_{p,s} = \sum_{\substack{p_1 + \dots + p_s = p \\ p_j \geq 1}} \binom{p}{p_1, \dots, p_s}$$

To verify this, think of $N_{p,s}$ as the number of ways p objects can be distributed over s distinct slots.

Lemma 2.1

One has

$$c_p = \sum_{s=1}^p N_{p,s} \prod_{j=1}^s v_b(\varepsilon j_1) \left(e^{-\beta \hat{\theta}_{\varepsilon j_1}} - 1 \right)$$

Proof

The above interpretation of $N_{p,s}$ leads to the recursion relation

$$N_{p,s} = \sum_{s-1 \leq q \leq p-1} \binom{p}{p-q} N_{q,s-1}, \quad (p \geq s > 1), \quad N_{p,1} = 1. \quad (2.28)$$

Substituting (2.28) in (2.25) yields a recursion relation for c_p

$$c_p = - \sum_{0 \leq q \leq p-1} \binom{p}{q} c_q, \quad (2.29)$$

whose unique solution is (2.27). $\square$

Definition 2.2 Let $t_1, \dots, t_n$ belong to a commutative algebra. The r th elementary symmetric function S_r of $t_1, \dots, t_n$ is defined as

$$S_r(t_1, \dots, t_n) \equiv \sum_{\substack{J \subset \{1, \dots, n\} \\ |J|=r}} \prod_{j \in J} t_j, \quad r = 1, \dots, n. \quad (2.30)$$

Proposition 2.3 If $g = 0$, then

$$\hat{D}_r = S_r(\hat{D}_1(x_1), \dots, \hat{D}_1(x_n)), \quad r = 1, \dots, n \quad (2.31)$$

(with $\hat{D}_1(x_j)$ defined by Eq. (2.23)).

Proof

Substituting (2.27) in (2.24) yields

$$\hat{D}_r = \sum_{\substack{J \subset \{1, \dots, n\}, |J|=r \\ \varepsilon_j = \pm 1, j \in J}} V_{\varepsilon J}^1 \sum_{\substack{I \subset J \\ 0 \leq |I| \leq r}} (-1)^{r-|I|} e^{-\beta \hat{\theta}_{\varepsilon I}}. \quad (2.32)$$

Using Eq. (2.16), one completes the proof of the proposition:

$$\hat{D}_r = \sum_{\substack{J \subset \{1, \dots, n\}, |J|=r \\ \varepsilon_j = \pm 1, j \in J}} \prod_{j \in J} v_b(\varepsilon_j x_j) \left(e^{-\beta \varepsilon_j \hat{\theta}_j} - 1 \right) \quad (2.33)$$

$$= \sum_{\substack{J \subset \{1, \dots, n\} \\ |J|=r}} \prod_{j \in J} \hat{D}_1(x_j) = S_r(\hat{D}_1(x_1), \dots, \hat{D}_1(x_n)). \quad (2.34)$$

$\square$

Note. Proposition 2.3 is in accordance with the previously noted fact that for $g = 0$ the particles of the quantum system become independent.

3 Simultaneous Diagonalization

In this section it is shown that Koornwinder's eigenfunctions of $\hat{D}_1, \dots, \hat{D}_n$. We prove that the different eigenvalues. As a result, we obtain an explicit basis for the abelian algebra generated by $\hat{D}_1, \dots, \hat{D}_n$. For convenience, we will put $\alpha = 1/2$ from now on.

3.1 Trigonometric Polynomials

Let $\mathcal{A} = \mathbb{C}[\exp(ix_1), \dots, \exp(ix_n)]$ be the algebra of functions on the n -dimensional torus $\mathbb{T} = \mathbb{R}^n / 2\pi\mathbb{Z}^n$.

$\mathcal{A}$ is spanned by the Fourier basis $\{e^\lambda\}$ with

$$e^\lambda(x) \equiv e^{i \sum_{j=1}^n \lambda_j x_j}$$

Let W be the (Weyl) group of permutations of n objects (so $W \cong S_n \ltimes (\mathbb{Z}_2)^n$). The subalgebra of $\mathcal{A}$ consisting of symmetric polynomials on $\mathbb{T}$ is spanned by the basis $\{m_\lambda\}$

$$m_\lambda(x) = \sum_{\lambda' \in W\lambda} e^{\lambda'} \sim \sum_{\lambda' \in S_n\lambda} e^{\lambda'}$$

with $\sim$ denoting proportionality and $\mathcal{P}^+$ denoting the set of dominant weights

$$\mathcal{P}^+ = \{\lambda \in \mathcal{P} \mid \lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_n\}$$

The lattice $\mathcal{P}$ can be partially ordered in the following way:

Definition 3.1 [partial ordering of $\mathcal{P}$]

$$(\forall \lambda, \lambda' \in \mathcal{P}) : \quad \lambda' \leq \lambda \quad \text{iff} \quad \sum_{i=1}^n \lambda'_i \leq \sum_{i=1}^n \lambda_i$$

The above ordering induces a partial ordering on $\mathcal{A}$. For each dominant weight $\lambda \in \mathcal{P}^+$ we associate a highest weight λ :

$$\mathcal{A}_\lambda^W \equiv \text{span}\{m_{\lambda'}\}_{\lambda' \leq \lambda}$$

Occasionally we will also use the notation

$$|\lambda| \equiv \sum_{j=1}^n \lambda_j$$

3.2 Triangularity

In this subsection it is shown that $\hat{D}_r$ maps the highest weight spaces $\mathcal{A}_\lambda^W$ into itself.

Definition 3.2 A linear operator $\hat{D} : \mathcal{A}^W \rightarrow \mathcal{A}^W$ is called *triangular* iff

$$\hat{D}(\mathcal{A}_\lambda^W) \subset \mathcal{A}_\lambda^W, \quad \forall \lambda \in \mathcal{P}^+. \quad (3.8)$$

One can rewrite Eq. (3.8) in a more illuminating way:

$$(\forall \lambda \in \mathcal{P}^+) : \quad \hat{D} m_\lambda = \sum_{\lambda' \in \mathcal{P}^+, \lambda' \leq \lambda} [\hat{D}]_{\lambda, \lambda'} m_{\lambda'}, \quad \text{with } [\hat{D}]_{\lambda, \lambda'} \in \mathbb{C} \quad (3.9)$$

(i.e. $[\hat{D}]_{\lambda, \lambda'} = 0$ if $\lambda' \not\leq \lambda$). In order to prove that $\hat{D}_r$ is triangular, we first need to verify that the operator maps $\mathcal{A}^W$ into itself.

Proposition 3.3 (invariance of $\mathcal{A}^W$)

$$\hat{D}_r(\mathcal{A}^W) \subset \mathcal{A}^W, \quad r = 1, \dots, n. \quad (3.10)$$

Proof

Acting with $\hat{D}_r$ Eq. (2.19) on a monomial m_λ (3.3) yields the following W-invariant trigonometric function on the torus $\mathbb{T}$:

$$\begin{aligned} (\hat{D}_r m_\lambda)(x) = & \sum_{\substack{J \subset \{1, \dots, n\}, |J|=r \\ \varepsilon_j = \pm 1, j \in J}} V_{\varepsilon J}^1 V_{\varepsilon J^c}^3 \times \sum_{\substack{\emptyset \subsetneq J_1 \subsetneq \dots \subsetneq J_s = J \\ 1 \leq s \leq r}} \\ & (-1)^{s-1} \left\{ \prod_{1 \leq s' \leq s} V_{\varepsilon(J_{s'} \setminus J_{s'-1})}^2 V_{\varepsilon(J_{s'} \setminus J_{s'-1})}^3; J \setminus J_{s'} [m_\lambda(x + 2\gamma e_{\varepsilon J_1}) - m_\lambda(x)] \right\}, \end{aligned} \quad (3.11)$$

with

$$e_{\varepsilon J} \equiv \sum_{j \in J} \varepsilon_j e_j \quad (3.12)$$

($\{e_1, \dots, e_n\}$ denotes the standard basis of $\mathbb{R}^n$). The r.h.s. of (3.11) is rational in the exponentials $\exp(ix_j)$, $j = 1, \dots, n$. In order to prove the proposition, we need to show that $\hat{D}_r m_\lambda$ (3.11) is actually a polynomial in $\exp(ix_1), \dots, \exp(ix_n)$. Since the r.h.s. of (3.11) is symmetric in $x_1, \dots, x_n$ it suffices to verify that $\hat{D}_r m_\lambda$, viewed as a function of x_1 , is free of poles.

As a function of x_1 , the terms in (3.11) may have poles caused by zeros in the denominators of the coefficients of the $\Lambda\Delta\mathcal{O}$. These poles are located at (cf. Eqs. (2.16)-(2.18) and (2.1), (2.2)):

$$\begin{aligned} x_1 &= 0 \quad \text{mod } \pi, \\ &= \pm \gamma \quad \text{mod } \pi, \\ &= \pm x_j \quad \text{mod } 2\pi, \quad j = 2, \dots, n, \\ &= \pm x_j \pm 2\gamma \quad \text{mod } 2\pi, \quad j = 2, \dots, n. \end{aligned} \quad (3.13)$$

From now on the parameters $\gamma, \mu, \mu_\delta, \mu'_\delta$ ($\delta = 0, 1$) and the remaining variables $x_2, \dots, x_n$ are fixed in general position. Specifically, we choose these parameters and variables such that the poles in the terms of (3.11) are simple.

The residue at $x_1 = 0$ vanishes because x_j vanishes because (3.11) is invariant under $x_j \rightarrow x_j + \pi$. Furthermore, because $\hat{D} m_\lambda(x)$ is even in x_j over half the period (cf. Remark vi, Section

$$x_j \rightarrow x_j + \pi, \quad j = 1, \dots, n \Leftarrow$$

(with $|\lambda| = \sum_{j=1}^n \lambda_j$), we need only show that

$$\begin{aligned} x_1 &= -\gamma \\ &= -x_j - 2\gamma \end{aligned}$$

type I: $x_1 = -\gamma$.

The only terms in the r.h.s. of (3.11) that contribute to the residue at $x_1 = -\gamma$ are those corresponding to cells J with $1 \in J$ and $\varepsilon_1 = -1$. For a cell J and choose a configuration of signs ε_j , the total residue at $x_1 = -\gamma$ in the sum of all terms corresponding to cell J , with the signs prescribed, is zero. One might think this is because the general situation can be changed by flipping of signs of the variables x_j , $j \in J$. However, the sum of terms in (3.11) corresponding to a fixed cell J is regular at $x_1 = -\gamma$.

type II: $x_1 = -x_j - 2\gamma$.

The proof of this case is very similar to the proof of type I. The only terms that contribute to the residue at $x_1 = -x_j - 2\gamma$ are those corresponding to cells J with $j \in J$ and $\varepsilon_j = -1$ (cf. (2.1) and (2.17)). Lemma A.2 of Appendix A.1 shows that these terms correspond to a fixed cell J with the signs ε_j prescribed. Again the general case (corresponding to all cells J) can be obtained by an appropriate flipping of the signs.

We conclude that $\hat{D}_r m_\lambda$ is a W -invariant polynomial. Consequently, $\hat{D}_r m_\lambda$ must be a polynomial. This proves the proposition.

Proposition 3.3 says that $\hat{D}_r m_\lambda$ is a W -invariant polynomial. It must be a finite linear combination of monomials.

$$(\forall \lambda \in \mathcal{P}^+) : \quad \hat{D}_r m_\lambda = \sum_{\lambda' \in \mathcal{P}^+} c_{\lambda, \lambda'} m_{\lambda'}$$

with

$$\mathcal{P}_{\lambda, r}^+ \equiv \{\lambda' \in \mathcal{P}^+ \mid \lambda' \leq \lambda + r\gamma\}$$

In order for $\hat{D}_r$ to be triangular one must have

$$\lambda' \in \mathcal{P}_{\lambda, r}^+ \implies \lambda' \leq \lambda$$

(cf. Eq. (3.9)). We shall prove this property by studying the asymptotics of $(\hat{D}_r m_\lambda)(x)$ for $\text{Im } x_j \rightarrow -\infty$. The following limits will be useful (cf. Eqs. (2.1), (2.2) and (2.20)):

$$\lim_{R \rightarrow \infty} v_a(z + i\varepsilon R) = e^{\varepsilon \beta g/2}, \quad (3.19)$$

$$\lim_{R \rightarrow \infty} v_b(z + i\varepsilon R) = e^{\varepsilon \beta (g_0 + g_1 + g'_0 + g'_1)/2} \quad (3.20)$$

(with $\varepsilon = \pm 1$).

Proposition 3.4 (triangularity)

$$(\forall \lambda \in \mathcal{P}^+) : \quad \hat{D}_r(\mathcal{A}_\lambda^W) \subset \mathcal{A}_\lambda^W, \quad r = 1, \dots, n. \quad (3.21)$$

Proof

Fix an $r \in \{1, \dots, n\}$ and $\lambda \in \mathcal{P}^+$. Let $\omega_k \equiv \sum_{1 \leq j \leq k} e_j$ (the k th fundamental weight) and introduce (cf. (3.17))

$$M_{\lambda, r; k} \equiv \max\{(\lambda', \omega_k) \mid \lambda' \in \mathcal{P}_{\lambda, r}^+\}. \quad (3.22)$$

To derive a contradiction, assume (3.18) does not hold; i.e. assume that there exists a $k \in \{1, \dots, n\}$ such that

$$M_{\lambda, r; k} > (\lambda, \omega_k) \left(= \sum_{1 \leq j \leq k} \lambda_j \right). \quad (3.23)$$

Now it is easy to verify the asymptotics

$$m_{\lambda'}(x - iR\omega_k) \sim e^{R(\lambda', \omega_k)} \sum_{\lambda'' \in W_{\lambda'; k}(\lambda')} e^{\lambda''}(x), \quad R \rightarrow +\infty, \quad (3.24)$$

with

$$W_{\lambda'; k} \equiv \{w \in W \mid (w\lambda', \omega_k) = (\lambda', \omega_k)\}, \quad (3.25)$$

so using (3.16) we obtain

$$\lim_{R \rightarrow \infty} e^{-RM_{\lambda, r; k}} (\hat{D}_r m_\lambda)(x - iR\omega_k) = \sum_{\substack{\lambda' \in \mathcal{P}_{\lambda, r}^+ \\ (\lambda', \omega_k) = M_{\lambda, r; k}}} [\hat{D}_r]_{\lambda, \lambda'} \left(\sum_{\lambda'' \in W_{\lambda'; k}(\lambda')} e^{\lambda''}(x) \right). \quad (3.26)$$

On the other hand, Eq. (3.24) combined with the limits (3.19) and (3.20) entails the following asymptotics for (3.11):

$$(\hat{D}_r m_\lambda)(x - iR\omega_k) = O(e^{R(\lambda, \omega_k)}), \quad R \rightarrow \infty. \quad (3.27)$$

Consequently,

$$\lim_{R \rightarrow \infty} e^{-RM_{\lambda, r; k}} (\hat{D}_r m_\lambda)(x - iR\omega_k) = 0 \quad (3.28)$$

(because of inequality (3.23)).

The matrix elements $[\hat{D}_r]_{\lambda, \lambda'}$ in (3.26) are non-zero (by definition), and the exponentials $e^{\lambda''}(x)$ (3.2) corresponding to different weights $\lambda'' \in \mathcal{P}$ are linearly independent. Hence, by comparing the r.h.s. of Eqs. (3.26) and (3.28) one arrives at the desired contradiction. $\square$

3.3 The Spectrum

By extending $\leq$ (Definition 3.1) to a linear ordering, it is easy to see that the triangularity

$$\hat{D}_r m_\lambda = \sum_{\lambda' \in \mathcal{P}^+, \lambda' \leq \lambda} [\hat{D}_r]_{\lambda, \lambda'} m_{\lambda'}$$

has as consequence that the elements on the right-hand side are eigenvectors of the operator $\hat{D}_r : \mathcal{A}^W \rightarrow \mathcal{A}^W$. (It follows that these eigenvalues are semisimple). The proof is complete. $\square$

Let $y \in \mathbb{R}^n$ be a fixed vector subject to the condition

$$y_1 > y_2 > \dots > y_n$$

By combining the asymptotics (cf. (3.24))

$$m_{\lambda'}(iRy) \sim e^{R \sum_{j=1}^n y_j \lambda'_j}$$

with Eq. (3.29), one finds (use $\lambda' < \lambda \Rightarrow \sum_{j=1}^n y_j \lambda'_j < \sum_{j=1}^n y_j \lambda_j$)

$$[\hat{D}_r]_{\lambda, \lambda} = \lim_{R \rightarrow \infty} e^{-R \sum_{j=1}^n y_j \lambda_j}$$

In the next proposition, we will evaluate this limit.

Proposition 3.5 (eigenvalues)

One has

$$[\hat{D}_r]_{\lambda, \lambda} = 2^r E_{r, n}(\text{ch} \beta(\lambda_1 + \rho_1), \dots, \text{ch} \beta(\lambda_n + \rho_n))$$

with

$$E_{r, n}(t_1, \dots, t_n; p_1, \dots, p_n) \equiv \sum_{0 \leq s \leq r} (-1)^{r+s} \left(\sum_{\substack{J \subset \{1, \dots, n\} \\ |J|=s}} \prod_{j \in J} t_j \prod_{j \notin J} p_j \right)$$

and

$$\rho_j \equiv (n-j)g + (g_0 + g_1 + \dots + g_n)$$

Proof

Using Eq. (2.13) we obtain

$$(\hat{D}_r m_\lambda)(x) = \sum_{0 \leq s \leq r} \sum_{\substack{J \subset \{1, \dots, n\}, |J|=s \\ \varepsilon_j = \pm 1, j \in J}} m_{\lambda'}(x)$$

with $V_{\varepsilon J;K}$ and $W_{I,p}$ defined by Eqs. (2.4) and (2.14), respectively. In order to compute the limit (3.32), we first derive some preliminary asymptotics:

i. m_λ (cf. Eq. (3.24)):

$$m_\lambda(x + i\beta e_{\varepsilon J})|_{x=iRy} \sim e^{R \sum_{j=1}^n \lambda_j y_j} e^{\beta \sum_{j \in J} \varepsilon_j \lambda_j}, \quad R \rightarrow \infty. \quad (3.37)$$

ii. $V_{\varepsilon J;J^c}$:

From (3.19), (3.20) one deduces

$$\lim_{R \rightarrow \infty} V_{\varepsilon J;J^c}|_{x=iRy} = e^{\beta[gN_{\varepsilon J} + (g_0 + g_1 + g'_0 + g'_1)M_{\varepsilon J}]/2}, \quad (3.38)$$

with

$$\begin{aligned} M_{\varepsilon J} &= |\{j \in J \mid \varepsilon_j = +1\}| - |\{j \in J \mid \varepsilon_j = -1\}| \\ &= \sum_{j \in J} \varepsilon_j \end{aligned} \quad (3.39)$$

and

$$\begin{aligned} N_{\varepsilon J} &= 2|\{j, k \in J \mid j < k, \varepsilon_j = +1\}| - 2|\{j, k \in J \mid j < k, \varepsilon_j = -1\}| + \\ &\quad 2|\{j \in J, k \in J^c \mid j < k, \varepsilon_j = +1\}| - 2|\{j \in J, k \in J^c \mid j < k, \varepsilon_j = -1\}| \\ &= 2|\{j \in J, k = 1, \dots, n \mid j < k, \varepsilon_j = +1\}| - \\ &\quad 2|\{j \in J, k = 1, \dots, n \mid j < k, \varepsilon_j = -1\}| \\ &= 2 \sum_{j \in J} (n - j) \varepsilon_j. \end{aligned} \quad (3.40)$$

Consequently,

$$\lim_{R \rightarrow \infty} V_{\varepsilon J;J^c}|_{x=iRy} = e^{\beta \sum_{j \in J} \varepsilon_j \rho_j} \quad (3.41)$$

with ρ_j defined by (3.35).

iii. $W_{I,p}$:

Using (2.14) and (3.41) it is not hard to see that $\lim_{R \rightarrow \infty} W_{I,p}|_{x=iRy}$ exists and depends only on the cardinality of I and on p (but not on the number of variables n). We define

$$F_{|I|,p} \equiv 2^{-p} \lim_{R \rightarrow \infty} W_{I,p}|_{x=iRy}. \quad (3.42)$$

Notice that (cf. Eq. (2.14))

$$F_{m,0} = 1, \quad m = 0, \dots, n. \quad (3.43)$$

After these preliminaries, we are now ready to compute limit (3.32). Substituting (3.36) in (3.32), and making use of (3.37), (3.41) and (3.42), we obtain

$$[\hat{D}_r]_{\lambda, \lambda} = 2^r \sum_{\substack{J \subset \{1, \dots, n\}, |J|=s \\ 0 \leq s \leq r}} \left(\prod_{j \in J} \text{ch} \beta (\lambda_j + \rho_j) \right) F_{n-s, r-s}. \quad (3.44)$$

It remains to calculate $F_{m,p}$, $1 \leq p \leq m \leq n$. For $p = 0$, $F_{m,0} = 1$ (see Eq. (2.6)); one obtains, therefore $\lambda = 0$:

$$\sum_{\substack{J \subset \{1, \dots, n\}, |J|=s \\ 0 \leq s \leq r}} \left(\prod_{j \in J} \text{ch}(\beta \rho_j) \right)$$

For a fixed number of variables n , this yields the same coefficients $F_{m,p}$ occur, now with $0 \leq p \leq m \leq n$ for $n' = 1, \dots, n$, and making use of Eq. (3.44) for n' equations in the $n(n+1)/2$ variables $F_{m,p}$.

In order to solve this system it is convenient to notice that $\tilde{\rho}_j$ (unlike ρ_j) does not depend on n . Substitution the $n(n+1)/2$ equations becomes

$$\sum_{\substack{J \subset \{1, \dots, n'\}, |J|=s \\ 0 \leq s \leq r}} \left(\prod_{j \in J} \text{ch}(\beta \tilde{\rho}_j) \right)$$

with condition (3.43). In Lemma B.1 of [A] it is shown that (3.44) has a unique solution:

$$F_{m,p} = (-1)^p \sum_{1 \leq i_1 \leq i_2 \leq \dots \leq i_p \leq m}$$

Substituting (3.47) in (3.44) and using $\tilde{\rho}_j$ (3.35).

For $g = 0$ all components of the vector ρ

$$\rho_1, \rho_2, \dots, \rho_n = (g_0, g_1, \dots, g_n)$$

Then, one can rewrite the above expression in terms of symmetric functions (see the remark following (3.35)).

$$\begin{aligned} [\hat{D}_r]_{\lambda, \lambda} &= 2^r \sum_{\substack{J \subset \{1, \dots, n\}, |J|=r}} \prod_{j \in J} (\text{ch} \beta (\lambda_j + \rho_j)) \\ &= 2^r S_r (\text{ch} \beta (\lambda_1 + \bar{\rho}), \dots, \text{ch} \beta (\lambda_n + \bar{\rho})) \end{aligned}$$

This equation is in agreement with Proposition 3.3.

3.4 Symmetry

In Ref. [K4] the following weight function is introduced:

$$\Delta(x) \equiv \prod_{1 \leq j < k \leq n} d_a(x_j + x_k)$$

with

$$d_c(z) = d_c^+(z)d_c^+(-z), \quad c = a, b \quad (3.51)$$

$$d_a^+(z) \equiv \frac{(e^{iz}; e^{-\beta})_\infty}{(e^{-\beta g} e^{iz}; e^{-\beta})_\infty}, \quad (3.52)$$

$$d_b^+(z) \equiv \frac{(e^{2iz}; e^{-\beta})_\infty}{(e^{-\beta g_0} e^{iz}, -e^{-\beta g_1} e^{iz}, e^{-\beta(g'_0+1/2)} e^{iz}, -e^{-\beta(g'_1+1/2)} e^{iz}; e^{-\beta})_\infty}. \quad (3.53)$$

The so-called q-shifted factorials are defined in the usual way:

$$(a; q)_\infty \equiv \prod_{m=1}^{\infty} (1 - aq^m), \quad (a_1, \dots, a_r; q)_\infty \equiv \prod_{s=1}^r (a_s; q)_\infty. \quad (3.54)$$

Notice that the conditions on our parameters, viz. (2.20), guarantee that the infinite products in Eqs. (3.52) and (3.53) converge. Recall that in order to compare our formulas with those of [K4], one has to reparametrize according to Eqs. (2.21) and (2.22).

Let $L_W^2(\mathbb{T}, \Delta dx)$ be the space of W-invariant functions on $\mathbb{T}$ that are square integrable with respect to the measure Δdx . We define

$$\langle f, g \rangle_\Delta \equiv \int_{\mathbb{T}} f \bar{g} \Delta dx, \quad f, g \in L_W^2(\mathbb{T}, \Delta dx). \quad (3.55)$$

The space of W-invariant polynomials $\mathcal{A}^W$ is a dense subspace of $L_W^2(\mathbb{T}, \Delta dx)$. The purpose of the present section is to show that the AΔO's $\hat{D}_1, \dots, \hat{D}_n$ are symmetric with respect to $\langle \cdot, \cdot \rangle_\Delta$. We need the following lemma:

Lemma 3.6 *Let $\alpha = 1/2$ and $z \in \mathbb{R}$; furthermore, let the parameters be subject to condition (2.20). Then the functions $d_c^{(+)}$ ($c = a, b$) satisfy the following first order difference equations:*

i. d_c^+ :

$$d_a^+(z + i\beta) = e^{-\beta g/2} v_a(z) d_a^+(z), \quad (3.56)$$

$$d_b^+(z + i\beta) = e^{-\beta(g_0+g_1+g'_0+g'_1)/2} v_b(z) d_b^+(z); \quad (3.57)$$

ii. d_c^- :

$$v_c(z - i\beta) d_c(z - i\beta) = \overline{v_c(z)} d_c(z), \quad c = a, b. \quad (3.58)$$

Proof

i. Eq. (3.56) is an immediate consequence of definition (3.52):

$$d_a^+(z + i\beta)/d_a^+(z) = \frac{(1 - e^{-\beta g} e^{iz})}{(1 - e^{iz})} = e^{-\beta g/2} v_a(z). \quad (3.59)$$

Eq. (3.57) can be reduced to the former case by observing that $d_b^+(z)$ factorizes:

$$d_b^+(z) = \frac{(e^{iz}; e^{-\beta})_\infty}{(e^{-\beta g_0} e^{iz}; e^{-\beta})_\infty} \frac{(-e^{iz}; e^{-\beta})_\infty}{(-e^{-\beta g_1} e^{iz}; e^{-\beta})_\infty} \times \frac{(e^{-\beta/2} e^{iz}; e^{-\beta})_\infty}{(e^{-\beta(g'_0+1/2)} e^{iz}; e^{-\beta})_\infty} \frac{(-e^{-\beta/2} e^{iz}; e^{-\beta})_\infty}{(-e^{-\beta(g'_1+1/2)} e^{iz}; e^{-\beta})_\infty}. \quad (3.60)$$

ii. Using Eqs. (3.56) or (3.57), respectively

$$\begin{aligned} v_c(z - i\beta) d_c(z - i\beta) &= v_c \frac{d_c^+}{d_c^+} \\ &= \frac{d_c^+}{d_c^+} \\ &= \frac{d_c^+}{d_c^+} \end{aligned}$$

Part ii. of the above lemma leads to the fo

Corollary 3.7 *One has*

$$e^{\beta \hat{\theta}_{\varepsilon J}} (V_{\varepsilon J}; J)$$

We now arrive at the main result of this proof hinges on relation (3.62).

Proposition 3.8 (symmetry)

$$\langle \hat{D}_r m_\lambda, m_{\lambda'} \rangle_\Delta = \langle m_\lambda, \hat{D}_r m_{\lambda'} \rangle_\Delta$$

Proof

First consider the following contour integrals

$$\oint_{C_j} W^{J^c, r-s} V_{\varepsilon J; J^c}$$

with $V_{\varepsilon J; K}$ and $W_{I, p}$ as in (2.4) and (2.14), place over the closed contour

$$C_j = [-\pi, \pi] \cup [\pi, \pi - i\varepsilon_j \beta] \cup [\pi - i\varepsilon_j \beta, -\pi]$$

Let all parameters and the variables x_k , $k=1, \dots, n$, the integrand has simple poles inside C_j and $\Delta(x)$. However, one easily verifies that $\Delta(x)$ has a zero in $\Delta(x)$; similarly, poles inside C_j and $V_{\varepsilon J; J^c}$. Consequently, integral (3.64) vanishes. The contributions to (3.64) which are due to the poles on the real axis, respectively, cancel each other because the integrand is odd. The upshot is that, when integrating the integrand over the integration path to $[-\pi - i\varepsilon_j \beta, \pi - i\varepsilon_j \beta]$ with

Armed with this conclusion and Eq. (3.62), the result can be written

$$\langle \hat{D}_r m_\lambda, m_{\lambda'} \rangle_\Delta \stackrel{\text{Eq. (2.13)}}{=} \sum_{\substack{|J|=s, 0 \leq s \leq r \\ \varepsilon_j = \pm 1, j \in J}} \int_{\mathbb{T}} \prod_{j \in J} V_{\varepsilon_j J; J^c}$$

Deformation of the integration paths of x_j , $j \in J$, from $[-\pi, \pi]$ to $[-\pi - i\varepsilon_j\beta, \pi - i\varepsilon_j\beta]$, followed by a change of variables $x_j \rightarrow x_j - i\beta\varepsilon_j$, $j \in J$, yields

$$\langle \hat{D}_r m_\lambda, m_{\lambda'} \rangle_\Delta = \sum_{\substack{|J|=s, 0 \leq s \leq r \\ \varepsilon_j = \pm 1, j \in J}} \int_{\mathbb{T}} W_{J^c, r-s} m_\lambda (e^{\beta \hat{\theta}_{\varepsilon J}} \overline{m_{\lambda'}} V_{\varepsilon J; J^c} \Delta) dx. \quad (3.67)$$

Using Corollary 3.7 and the fact that $W_{J^c, r-s}$ is real (for parameters subject to (2.20)) entails

$$\begin{aligned} \langle \hat{D}_r m_\lambda, m_{\lambda'} \rangle_\Delta &= \sum_{\substack{|J|=s, 0 \leq s \leq r \\ \varepsilon_j = \pm 1, j \in J}} \int_{\mathbb{T}} m_\lambda \overline{W_{J^c, r-s} V_{\varepsilon J; J^c} (e^{-\beta \hat{\theta}_{\varepsilon J}} m_{\lambda'})} \Delta dx \\ &= \langle m_\lambda, \hat{D}_r m_{\lambda'} \rangle_\Delta. \end{aligned} \quad (3.68)$$

□

3.5 Diagonalization and Commutativity

If $g = g_0 = g_1 = g'_0 = g'_1 = 0$, then $d_a = d_b = 1$ (see Eqs. (3.51), (3.52) and (3.60)), and thus $\langle \cdot, \cdot \rangle_\Delta$ reduces to the inner product on $\mathbb{T}$ with respect to Lebesgue measure ($\Delta = 1$). The basis of monomials $\{m_\lambda\}_{\lambda \in \mathcal{P}^+}$ is an orthogonal basis of $L_W^2(\mathbb{T}, dx)$. For arbitrary parameters however, the orthogonality of the monomials with respect to $\langle \cdot, \cdot \rangle_\Delta$ no longer holds. By subtracting from m_λ the orthogonal projection of m_λ onto $\text{span}\{m_{\lambda'}\}_{(\lambda' \in \mathcal{P}^+, \lambda' < \lambda)} \subset \mathcal{A}_\lambda^W$, one obtains an alternative basis $\{p_\lambda\}_{\lambda \in \mathcal{P}^+}$ of $\mathcal{A}^W$. This is the basis of Koornwinder polynomials.

Definition 3.9 [Koornwinder polynomials]

Koornwinder's polynomial $p_\lambda \in \mathcal{A}_\lambda^W$ is defined by the conditions

$$p_\lambda = m_\lambda + \sum_{\lambda' \in \mathcal{P}^+, \lambda' < \lambda} c_{\lambda, \lambda'} m_{\lambda'} \quad (3.69)$$

and

$$\langle p_\lambda, m_{\lambda'} \rangle_\Delta = 0, \quad \forall \lambda' \in \mathcal{P}^+, \lambda' < \lambda. \quad (3.70)$$

We now prove that $\{p_\lambda\}_{\lambda \in \mathcal{P}^+}$ is a basis of joint eigenfunctions of $\hat{D}_1, \dots, \hat{D}_n$. For convenience, the notation for the eigenvalues Eq. (3.33) is sometimes abbreviated by putting

$$E_{r,n}(\text{ch}\beta\theta_1, \dots, \text{ch}\beta\theta_n; \text{ch}\beta\rho_r, \dots, \text{ch}\beta\rho_n) \longrightarrow E_{r,n}(\theta), \quad (\theta \in \mathbb{R}^n). \quad (3.71)$$

Theorem 3.10 (eigenfunctions)

$$\hat{D}_r p_\lambda = E_{r,n}(\lambda + \rho) p_\lambda, \quad \forall \lambda \in \mathcal{P}^+ \quad (3.72)$$

(with $r = 1, \dots, n$ and $\rho = (\rho_1, \dots, \rho_n)$, see Eq. (3.35)).

Proof

It follows from (3.69) and Proposition 3.4 that

$$(\hat{D}_r - [\hat{D}_r])$$

On the other hand, one has (use the proposition 3.4)

$$\langle (\hat{D}_r - [\hat{D}_r])_{\lambda, \lambda} p_\lambda, m_{\lambda'} \rangle_\Delta \stackrel{\text{symmetry}}{=} \langle p_\lambda, (\hat{D}_r - [\hat{D}_r])_{\lambda, \lambda} m_{\lambda'} \rangle_\Delta$$

Combining (3.73) and (3.74) entails $\hat{D}_r p_\lambda = E_{r,n}(\lambda + \rho) p_\lambda$ to complete the proof.

In Appendix C it is shown that if a difference operator is diagonalizable, then all its coefficients must be zero. Combining this with the commutativity of the $\Lambda\Delta$ O's:

Theorem 3.11 (commutativity)

The operators $\hat{D}_1, \dots, \hat{D}_n$ mutually commute.

Proof

The polynomials $\{p_\lambda\}_{\lambda \in \mathcal{P}^+}$ form a basis of $\mathcal{A}^W$. The operators $\hat{D}_1, \dots, \hat{D}_n$ (Theorem 3.10). Hence, it is sufficient to show that $[\hat{D}_r, \hat{D}_{r'}] p_\lambda = 0$ on $\mathcal{A}^W$. In other words, the commutator vanishes on $\mathcal{A}^W$:

$$[\hat{D}_r, \hat{D}_{r'}](\mathcal{A}^W) = 0,$$

It now follows from Proposition C.1 in Appendix C that the operators are identically zero.

Consider the real algebra of difference operators

$$\mathbb{D} \equiv \mathbb{R}[\hat{D}_1, \dots, \hat{D}_n]$$

It is clear that $\mathbb{D}$ is an abelian algebra (Theorem 3.11) and is simultaneously diagonalized by the Koornwinder polynomials.

Theorem 3.12 ($\mathbb{D} \cong \mathbb{R}[\text{ch}\beta\theta_1, \dots, \text{ch}\beta\theta_n]$)

For each symmetric function $S(\theta) \in \mathbb{R}[\text{ch}\beta\theta_1, \dots, \text{ch}\beta\theta_n]$, there is a unique operator $\hat{D} \in \mathbb{D}$ such that

$$\hat{D} p_\lambda = S(\lambda + \rho) p_\lambda$$

Proof

$E_{r,n}(\theta)$ (3.71) is a linear combination of elementary symmetric functions

$$E_{r,n}(\theta) = S_r(\text{ch}\beta\theta_1, \dots, \text{ch}\beta\theta_n)$$

l.d. stands for terms of lower degree in $\text{ch}\beta\theta_j$, $j = 1, \dots, n$. The elementary symmetric functions form a set of algebraically independent generators of the symmetric algebra (this fact is the ‘fundamental theorem on symmetric functions’, see e.g. [M1]). Hence, Eq. (3.78) implies that the same is true for the functions $E_{r,n}(\theta)$: every element in $\mathbb{R}[\text{ch}\beta\theta_1, \dots, \text{ch}\beta\theta_n]^{S_n}$ can be written uniquely as a polynomial in $E_{r,n}(\theta)$, $r = 1, \dots, n$.

Now we use Theorem 3.10 to conclude that for every symmetric function $S(\theta)$ there exists a difference operator $\hat{D} \in \mathbb{D}$ such that Eq. (3.77) holds. That such a difference operator $\hat{D}$ is unique follows from Proposition C.1 (Appendix C).

□

The symmetric functions $S(\theta) \in \mathbb{R}[\text{ch}\beta\theta_1, \dots, \text{ch}\beta\theta_n]^{S_n}$ separate the points of the wedge

$$\{\theta \in \mathbb{R}^n \mid \theta_1 \geq \theta_2 \geq \dots \geq \theta_n \geq 0\}. \quad (3.79)$$

To see this, first notice that $(-1)^{n-r} S_r(\text{ch}\beta\theta_1, \dots, \text{ch}\beta\theta_n)$ is the coefficient of ν^{n-r} in the characteristic polynomial $\det(T - \nu I)$ of the diagonal matrix $T = \text{diag}(\text{ch}\beta\theta_1, \dots, \text{ch}\beta\theta_n)$. Consequently, the values of $S_r(\text{ch}\beta\theta_1, \dots, \text{ch}\beta\theta_n)$, $r = 1, \dots, n$ determine θ in the wedge (3.79) uniquely. This fact combined with Theorem 3.12 can be used to prove the orthogonality of the basis $\{p_\lambda\}$:

Corollary 3.13 (orthogonality)

$$\langle p_\lambda, p_{\lambda'} \rangle_\Delta = 0, \quad \forall \lambda, \lambda' \in \mathcal{P}^+, \lambda \neq \lambda'. \quad (3.80)$$

Proof

Let $\lambda, \lambda' \in \mathcal{P}^+$, and $\lambda \neq \lambda'$. Since the symmetric functions in $\mathbb{R}[\text{ch}\beta\theta_1, \dots, \text{ch}\beta\theta_n]^{S_n}$ separate the points of the wedge (3.79), there exists an $S(\theta) \in \mathbb{R}[\text{ch}\beta\theta_1, \dots, \text{ch}\beta\theta_n]^{S_n}$ such that

$$S(\lambda + \rho) \neq S(\lambda' + \rho). \quad (3.81)$$

Hence, by Theorem 3.12 there exists a $\hat{D} \in \mathbb{D}$ for which p_λ and $p_{\lambda'}$ are eigenfunctions corresponding to different eigenvalues. But then the polynomials p_λ and $p_{\lambda'}$ must be orthogonal with respect to $\langle \cdot, \cdot \rangle_\Delta$ because $\hat{D}$ is symmetric, cf. Proposition 3.8.

□

Note. The orthogonality of the basis $\{p_\lambda\}_{\lambda \in \mathcal{P}^+}$ was already shown by Koornwinder [K4]. His proof exploits the continuity of $\langle p_\lambda, p_{\lambda'} \rangle_\Delta$ in the parameters.

Corollary 3.14 (self-adjointness)

Every difference operator $\hat{D} \in \mathbb{D}$ is essentially self-adjoint on $\mathcal{A}^W \subset L_W^2(\mathbb{T}, \Delta dx)$.

Proof

This is an immediate consequence of the fact that every $\mathbf{A}\Delta\mathbf{O}$ in $\mathbb{D}$ acts as a real multiplication operator on the orthogonal basis $\{p_\lambda\}_{\lambda \in \mathcal{P}^+}$ of $L_W^2(\mathbb{T}, \Delta dx)$.

□

Remarks i. Theorem 3.12 states that the as

$$\hat{D}_r \xrightarrow{HC} E_{r,n}(\theta)$$

induce a Harish-Chandra-type algebra isom

ii. Recall that p_λ is defined as m_λ restricted to $\text{span}\{m_{\lambda'}\}_{\lambda' < \lambda}$. Using the orthogonality of the basis $\{p_\lambda\}$ and the following recursion relation for p_λ :

$$p_\lambda = m_\lambda - \sum_{\lambda' \in \mathcal{P}^+} \dots$$

iii. If the partial ordering 3.1 is extended to $\mathcal{P}^+$, it is possible to orthogonalize the basis $\{m_\lambda\}$ using Corollary 3.13, the result does not depend on the ordering: the resulting orthogonal basis is unique up to the restrictive property of the measure Δdx .

iv. According to Theorem 3.11, the Harish-Chandra-type integrable n -particle system (cf. the note e

4 $\beta \rightarrow 0$: The Transition to the pergeometric PDO's

By sending the step size β of the differences to zero, the pergeometric PDO's associated with the root system Φ converge to the BC_n -type Jacobi polynomials $\{p_\lambda\}$. In this section we will make the dependence of all objects of interest, e.g.: $\hat{D}_{r,\beta}$, Δ_β and p_λ on β explicit.

4.1 Eigenfunctions

Consider the following weight function on $\mathbb{T}$:

$$\Delta_0(x) \equiv \prod_{1 \leq j < k \leq n} |\sin \alpha(x_j + x_k)| \prod_{1 \leq j \leq n} |\sin(\alpha x_j)|^{2\tilde{g}_0} |\cos \alpha x_j|^{2\tilde{g}_1}$$

and let $\langle \cdot, \cdot \rangle_{\Delta_0}$ be the inner product on $L^2(\mathbb{T}, \Delta_0 dx)$. We introduce W -invariant polynomials on $\mathbb{T}$ (Definition 3.9); these are the BC_n -type Jacobi polynomials.

Definition 4.1 [BC_n -type Jacobi polynomial] The *Jacobi polynomial* $p_{\lambda,0} \in \mathcal{A}_\lambda^W$ is defined by

$$p_{\lambda,0} = m_\lambda + \dots$$

and

$$\langle p_{\lambda,0}, m_{\lambda'} \rangle_{\Delta_0} = 0, \quad \forall \lambda' \in \mathcal{P}^+, \lambda' < \lambda. \quad (4.3)$$

Remark. Usually Δ_0 is written in a slightly different form, which emphasizes the relation with the root system BC_n . Use

$$|\sin(\alpha x_j)|^{2\tilde{g}_0} |\cos(\alpha x_j)|^{2\tilde{g}_1} \sim |\sin(\alpha x_j)|^{2k_0} |\sin(2\alpha x_j)|^{2k_1} \quad (4.4)$$

with $\tilde{g}_0 = k_0 + k_1$ and $\tilde{g}_1 = k_1$, to compare (4.1) with the usual expression for the BC_n -type weight function.

We need the following convergence result from [K2] to connect $p_{\lambda,0}$ with Koornwinder's polynomial $p_{\lambda,\beta}$, $\beta > 0$:

Lemma 4.2 *Let $k_1, k_2 \in \mathbb{R}$ and $q \in]0, 1[$. Then*

$$\lim_{q \uparrow 1} \frac{(q^{k_1} z; q)_\infty}{(q^{k_2} z; q)_\infty} = (1 - z)^{k_2 - k_1}, \quad (4.5)$$

uniformly for z in compacts of the punctured disc $\{z \in \mathbb{C} \mid |z| \leq 1, z \neq 1\}$.

Proof

See Proposition A.2 of Ref. [K2, Appendix A].

□

It is immediate from Eqs. (3.52), (3.60) and the above lemma that

$$\lim_{\beta \rightarrow 0} d_{a,\beta}(z) = |2 \sin(z/2)|^{2g}, \quad (4.6)$$

$$\lim_{\beta \rightarrow 0} d_{b,\beta}(z) = |2 \sin(z/2)|^{2(g_0 + g'_0)} |2 \cos(z/2)|^{2(g_1 + g'_1)}, \quad (4.7)$$

uniformly for z in compacts of $] \pi, \pi[\setminus \{0\}$.

Consequently, for $\beta \rightarrow 0$, Δ_β (3.50) converges to a weight function that is proportional to Δ_0 :

$$\lim_{\beta \rightarrow 0} \Delta_\beta = c \Delta_0, \quad (4.8)$$

with

$$\tilde{g}_0 = g_0 + g'_0 (\geq 0), \quad \tilde{g}_1 = g_1 + g'_1 (\geq 0) \quad (4.9)$$

and $c = 4^{n(n-1)g} 4^{n(\tilde{g}_0 + \tilde{g}_1)} (> 0)$.

Proposition 4.3 *One has*

$$\lim_{\beta \rightarrow 0} p_{\lambda,\beta} = p_{\lambda,0}, \quad \forall \lambda \in \mathcal{P}^+, \quad (4.10)$$

and

$$\langle p_{\lambda,0}, p_{\lambda',0} \rangle_{\Delta_0} = 0, \quad \forall \lambda, \lambda' \in \mathcal{P}^+, \lambda \neq \lambda'. \quad (4.11)$$

Proof

Using (3.83), (4.8) and an induction argument, one satisfies a recursion relation of the type (3.8) and Eq. (4.8) to conclude that the resulting $\langle \cdot, \cdot \rangle_{\Delta_0}$. But then the recursion relation for the orthogonal projection (with respect to Definition 4.1 one obtains (4.10).

Note. The limit $\beta \rightarrow 0$ corresponds to the limit of the polynomials the $q \rightarrow 1$ limit to the Jacobi polynomials studied in [M2], for arbitrary root systems. This result was proved in [H1] (again for arbitrary root systems).

4.2 Eigenvalues

The purpose of this subsection is to investigate the eigenvalues (Proposition 3.5) for $\beta \rightarrow 0$.

Proposition 4.4 *One has (for $r = 1, \dots, n$),*

$$\begin{aligned} \lim_{\beta \rightarrow 0} \beta^{-2r} E_{r,n}(\text{ch}\beta\theta_1, \dots, \text{ch}\beta\theta_n; \text{ch}\beta\rho_r, \dots, \text{ch}\beta\rho_n) \\ = E_{r,n}(\theta_1^2/2, \dots, \theta_n^2/2; \rho_r^2/2, \dots, \rho_n^2/2), \end{aligned}$$

Proof

Our proof of Eq. (4.12) hinges on a recursion relation to Lemma B.2 of Appendix B:

$$\begin{aligned} E_{r,n}(\text{ch}\beta\theta_1, \dots, \text{ch}\beta\theta_n; \text{ch}\beta\rho_r, \dots, \text{ch}\beta\rho_n) \\ = (\text{ch}\beta\theta_n - \text{ch}\beta\rho_n) E_{r-1,n-1}(\text{ch}\beta\theta_1, \dots, \text{ch}\beta\theta_{n-1}; \text{ch}\beta\rho_r, \dots, \text{ch}\beta\rho_{n-1}) \\ + E_{r,n-1}(\text{ch}\beta\theta_1, \dots, \text{ch}\beta\theta_{n-1}; \text{ch}\beta\rho_r, \dots, \text{ch}\beta\rho_{n-1}) \end{aligned}$$

with the convention

$$E_{0,n} \equiv 1,$$

We divide Eq. (4.13) by β^{2r} and then use

$$\begin{aligned} \lim_{\beta \rightarrow 0} \beta^{-2r} E_{r,n}(\text{ch}\beta\theta_1, \dots, \text{ch}\beta\theta_n; \text{ch}\beta\rho_r, \dots, \text{ch}\beta\rho_n) \\ = (\theta_n^2/2 - \rho_n^2/2) E_{r-1,n-1}(\theta_1^2/2, \dots, \theta_{n-1}^2/2; \rho_r^2/2, \dots, \rho_{n-1}^2/2) \\ + E_{r,n-1}(\theta_1^2/2, \dots, \theta_{n-1}^2/2; \rho_r^2/2, \dots, \rho_{n-1}^2/2) \end{aligned}$$

Now we use again Lemma B.2 to conclude (4.14).

4.3 Operators

Expansion of $\hat{D}_{r,\beta}$ in β yields a formal power series of the form

$$\hat{D}_{r,\beta} = \sum_{m=0}^{\infty} \hat{D}_r^{(m)} \beta^m. \quad (4.16)$$

The coefficients $\hat{D}_r^{(m)}$ are polynomials in the partials $\hat{\theta}_j$, $j = 1, \dots, n$; this means that these coefficients are PDO's. We define the leading differential operator $\hat{D}_{r,0}$ of $\hat{D}_{r,\beta}$ as the first nonzero coefficient in expansion (4.16):

Definition 4.5 [leading PDO]

Let

$$m_r \equiv \min\{m \in \mathbb{N} \mid \hat{D}_r^{(m)} \neq 0\} \quad (4.17)$$

(with $\hat{D}_r^{(m)}$ defined by expansion (4.16)). Then,

$$\hat{D}_{r,0} \equiv \hat{D}_r^{(m_r)} \quad (4.18)$$

is called the *leading PDO* of $\hat{D}_{r,\beta}$.

The BC_n -type Jacobi polynomials are joint eigenfunctions of $\hat{D}_{1,0}, \dots, \hat{D}_{n,0}$:

Theorem 4.6 *One has $m_r = 2r$ and*

$$\hat{D}_{r,0} p_{\lambda,0} = E_{r,n}((\lambda_1 + \rho_1)^2, \dots, (\lambda_n + \rho_n)^2; \rho_r^2, \dots, \rho_n^2) p_{\lambda,0}, \quad \forall \lambda \in \mathcal{P}^+, \quad (4.19)$$

with $r = 1, \dots, n$ and ρ as in (3.35).

Proof

Consider the eigenvalue equation (3.72):

$$\hat{D}_{r,\beta} (p_{\lambda,\beta}) = 2^r E_{r,n}(\text{ch}\beta(\lambda_1 + \rho_1), \dots, \text{ch}\beta(\lambda_n + \rho_n); \text{ch}\beta\rho_r, \dots, \text{ch}\beta\rho_n) p_{\lambda,\beta}. \quad (4.20)$$

First, apply Taylor's theorem to the l.h.s. of Eq. (4.20) and make use of Definition 4.5 and Limit (4.10) to conclude that

$$\hat{D}_{r,\beta} p_{\lambda,\beta} = \hat{D}_{r,0} p_{\lambda,0} \beta^{m_r} + o(\beta^{m_r}). \quad (4.21)$$

Next, use Proposition 4.4 and Limit (4.10) to derive the asymptotic behavior for $\beta \rightarrow 0$ of the r.h.s. of (4.20):

$$2^r E_{r,n}(\text{ch}\beta(\lambda_1 + \rho_1), \dots, \text{ch}\beta(\lambda_n + \rho_n); \text{ch}\beta\rho_r, \dots, \text{ch}\beta\rho_n) p_{\lambda,\beta} = E_{r,n}((\lambda_1 + \rho_1)^2, \dots, (\lambda_n + \rho_n)^2; \rho_r^2, \dots, \rho_n^2) p_{\lambda,0} \beta^{2r} + o(\beta^{2r}). \quad (4.22)$$

By Definition 4.5 and Proposition C.1 it is possible to pick a $\lambda \in \mathcal{P}^+$ such that $\hat{D}_{r,0} p_{\lambda,0} \neq 0$, so $m_r \geq 2r$. It is not difficult to see that there also exist $\lambda \in \mathcal{P}^+$ such that

$$E_{r,n}((\lambda_1 + \rho_1)^2, \dots, (\lambda_n + \rho_n)^2; \rho_r^2, \dots, \rho_n^2) \neq 0,$$

so $m_r \leq 2r$. This entails $m_r = 2r$ and (4.19).

Corollary 4.7

$$\hat{D}_{r,0} = \lim_{\beta \rightarrow 0} \hat{D}_{r,\beta}$$

Explicit computation of (4.23) for $r = 1$ yields

$$\begin{aligned} \hat{D}_{1,0} &= \sum_{1 \leq j \leq n} \hat{\theta}_j^2 - 2i\alpha g \sum_{1 \leq j < k \leq n} \{\cot \alpha x_j \cot \alpha x_k - \cot \alpha x_j \cot \alpha x_k\} \\ &\quad - 2i\alpha \sum_{1 \leq j \leq n} \{\tilde{g}_0 \cot(\alpha x_j)\} \end{aligned}$$

(with $\alpha = 1/2$ and $\tilde{g}_0, \tilde{g}_1$ as in Eq. (4.9)).

An immediate consequence of Theorem 4.6 is that $\hat{D}_{r,0}$, $r = 1, \dots, n$.

Corollary 4.8 *The differential operators $\hat{D}_{1,0}, \dots, \hat{D}_{n,0}$ are simultaneously diagonalized by the BC_n -type Jacobi polynomials.*

Let $\mathbb{D}_0 \equiv \mathbb{R}[\hat{D}_{1,0}, \dots, \hat{D}_{n,0}]$. The algebra $\mathbb{D}_0$ is simultaneously diagonalized by the BC_n -type Jacobi polynomials.

Theorem 4.9 ($\mathbb{D}_0 \cong \mathbb{R}[\theta_1^2, \dots, \theta_n^2]^{S_n}$)
For every symmetric polynomial $S(\theta) \in \mathbb{R}[\theta_1^2, \dots, \theta_n^2]^{S_n}$, there exists a unique operator $\hat{D}_0 \in \mathbb{D}_0$ such that

$$\hat{D}_0 p_{\lambda,0} = S(\lambda + \rho) p_{\lambda,0}$$

Corollary 4.10 (self-adjointness)

The PDO's $\hat{D}_0 \in \mathbb{D}_0$ are essentially self-adjoint on $L^2(\mathbb{R}^n)$.

The proofs of Theorem 4.9 and Corollary 4.10 are given in [13] and Corollary 3.14, respectively.

Remarks i. The operator $\hat{D}_{1,0}$ (4.24) can be identified with the hypergeometric PDO associated with the root system BC_1 with the usual notation which emphasizes the dependence on $\tan(\alpha x_j)$ from (4.24) by means of the relation $\tan(\alpha x_j) = \frac{1}{2}(\theta_j^2 - \theta_j^{-2})$ (the remark under Definition 4.1).

ii. The existence of an abelian algebra of differential operators on $\mathbb{R}[\theta_1^2, \dots, \theta_n^2]^{S_n}$ via a Harish-Chandra-type map was shown by Heckman and Opdam [H1, O2]. This result can be extended to consider arbitrary root systems.

iii. For an arbitrary root system R , the operator $\hat{D}_0$ can be identified with the generalized Calogero-Sutherland system via a similarity transformation [OP2]. This amounts to the conjugation with $\Delta^{1/2}$, where Δ is the Plancherel measure with weight function Δ .

$$\begin{aligned}
\hat{H} &= \Delta_0^{1/2} \hat{D}_{1,0} \Delta_0^{-1/2} + E_0 \\
&= \sum_{1 \leq j \leq n} \hat{\theta}_j^2 + 2g(g-1) \alpha^2 \sum_{1 \leq j < k \leq n} \{ \sin^{-2} \alpha(x_j + x_k) + \sin^{-2} \alpha(x_j - x_k) \} \\
&\quad + \alpha^2 \sum_{1 \leq j \leq n} \{ \tilde{g}_0(\tilde{g}_0 - 1) \sin^{-2}(\alpha x_j) + \tilde{g}_1(\tilde{g}_1 - 1) \cos^{-2}(\alpha x_j) \}
\end{aligned} \tag{4.26}$$

with $E_0 = 4\alpha^2(\rho, \rho)$ and $\rho_j = (n-j)g + (\tilde{g}_0 + \tilde{g}_1)/2$, cf. (3.35).

Corollary 4.8 can be interpreted as the quantum integrability of the BC_n -type Calogero-Sutherland system. For arbitrary root systems integrability follows from [H1, O2] (cf. Remark ii).

5 Special Cases Related to Classical Root Systems

By limit transitions and/or specialization of the parameters, the operators $\hat{D}_1, \dots, \hat{D}_n$ reduce to commuting $A\Delta O$'s, which are simultaneously diagonalized by Macdonald's polynomials. Such difference operators are obtained for all Macdonald families associated with (admissible pairs of) the classical root systems: A_{n-1} , B_n , C_n , D_n and BC_n .

Note. Most results in this section have an obvious counterpart for $\beta = 0$ (which amounts to $q = 1$).

5.1 Preliminaries

First, we outline very briefly some of the main points of the construction presented by Macdonald [M2]. A more detailed summary of his results can be found in [M4] and [K4]. (For our purposes, especially the second summary is useful). Here, we only want to introduce some terminology which facilitates clarifying the connection between the preceding sections and Ref. [M2]. For general information on root systems the reader is referred to e.g. [B, S]. Although most of the remaining part of the paper should be accessible without a detailed knowledge of root systems, a glance at the 'planches' in Bourbaki [B] might be of some help.

Ref. [M2] uses the concept of admissible pairs of root systems. The pair (R, S) is admissible if R and S are root systems (assumed irreducible) such that $S \subset R$ is reduced and generates the same Weyl group as R . Let V be the real vector space spanned by R and consider the torus $\mathbb{T}_R \equiv V/(2\pi\mathbb{Z}R^\vee)$. Let $\mathcal{P}_R^+$ be the dominant cone of the weight lattice of R (which equals the character lattice of $\mathbb{T}_R$) and let $\mathcal{A}_R^W$ denote the algebra of W -invariant (trigonometric) polynomials on $\mathbb{T}_R$ (this algebra is isomorphic to the W -invariant part of the group algebra over the weight lattice). To every admissible pair (R, S) Macdonald associates a weight function $\Delta_{(R,S)}$ on $\mathbb{T}_R$ and finds a corresponding orthogonal basis $\{p_{\lambda, (R,S)}\}_{\lambda \in \mathcal{P}_R^+}$ of $\mathcal{A}_R^W$. Furthermore, he introduces difference operators $\hat{D}_\sigma$ associated with the so-called (quasi-)minuscule weights σ of $S^\vee$. These operators are diagonalized by the basis of Macdonald polynomials $\{p_{\lambda, (R,S)}\}_{\lambda \in \mathcal{P}_R^+}$.

We will show that additional $A\Delta O$'s for Macdonald's polynomials arise as special cases of $\hat{D}_1, \dots, \hat{D}_n$ (2.6). This leads to difference operators associated with the fundamental weights of $S^\vee$, for every admissible pair consisting of classical root systems. These $A\Delta O$'s

generate an abelian algebra $\mathbb{D}_{(R,S)}$ of difference operators diagonalized by the basis $\{p_{\lambda, (R,S)}\}_{\lambda \in \mathcal{P}_R^+}$. The W -invariant part of the real group algebra

Remark. If all roots in R have the same length, then $S = R$ and there exists only one admissible pair. If R is not simply laced, then there are several possibilities for the pair (R, S) ; there are six such possibilities; these correspond to the six classical root systems.

5.2 The Root System A_{n-1}

Let $\hat{D}_{r,lead}$ consist of those terms in $\hat{D}_r$ which are diagonalized by $\exp(-\beta\hat{\theta}_j)$, $j = 1, \dots, n$ (cf. Eq. (2.13)):

$$\hat{D}_{r,lead} = \sum_{\substack{J \subset \{1, \dots, n\} \\ |J|=r}} V_{J,J}$$

One picks up these leading terms via the formula

$$\hat{D}_{r,lead} = \lim_{R \rightarrow \infty} e^{-\beta\hat{\theta}_j}$$

with

$$\Lambda_R \equiv e^{-\beta\hat{\theta}_j}$$

Let

$$\tilde{\Delta}_+ \equiv \prod_{1 \leq j < k \leq n} (e^{g(x_j + x_k)/2i} d_a^+(x_j + x_k))$$

Conjugation of $\hat{D}_{r,lead}$ with $\tilde{\Delta}_+$ results in (3.57):

$$\begin{aligned}
\hat{D}'_r &= \tilde{\Delta}_+ \hat{D}_{r,lead} \tilde{\Delta}_+^{-1} \\
&= \sum_{\substack{J \subset \{1, \dots, n\} \\ |J|=r}} \left(\prod_{\substack{j \in J \\ k \in J^c}} v_a(x_j) \right)
\end{aligned}$$

It is clear from Theorem 3.11 and Eqs. (3.57) that $\hat{D}'_r$ and $\hat{D}'_s$ commute. For $r = n$, (5.6) reduces to an operator in the coordinates: $\hat{D}'_n = \exp(-\beta(\hat{\theta}_1 + \dots + \hat{\theta}_n))$ (up to a multiplicative constant) with Macdonald polynomials as eigenfunctions.

$$\hat{D}'_r = e^{-\beta r(\hat{\theta}_1 + \dots + \hat{\theta}_n)/n} \hat{D}_r$$

The first part causes a translation $x \rightarrow x + i\beta r x/n$ (up to a multiplicative constant) with Macdonald polynomials as eigenfunctions.

$$E_{\omega_r} = c_r \hat{D}_{r, A_{n-1}}, \quad c_r \equiv e^{-\beta r(\hat{\theta}_1 + \dots + \hat{\theta}_n)/n}$$

The operator E_{ω_r} is associated with the r th fundamental weight ω_r of the root system A_{n-1} . The parameters in [M2] are related to ours via Eq. (2.21).

Notes i. For $r = 1$, Eq. (5.6) reduces to

$$\hat{D}'_1 = \sum_{1 \leq j \leq n} \left(\prod_{k \neq j} v_a(x_j - x_k) \right) e^{-\beta \hat{\theta}_j}. \quad (5.9)$$

ii. After transformation to Lebesgue measure, the operator $\hat{D}'_r$ goes over in the r -th quantum integral $\hat{S}_r$ [R1, (Eq. (2.3))] of the relativistic Calogero-Moser system with trigonometric coefficients. More precisely, let

$$\Delta_{A_{n-1}}(x) \equiv \prod_{1 \leq j < k \leq n} d_a(x_j - x_k), \quad (5.10)$$

then

$$\begin{aligned} \hat{S}_r &= \Delta_{A_{n-1}}^{1/2} \hat{D}'_r \Delta_{A_{n-1}}^{-1/2} \\ &= \sum_{\substack{J \subset \{1, \dots, n\} \\ |J|=r}} \left(\prod_{\substack{j \in J \\ k \in J^c}} v_a(x_j - x_k) \right)^{1/2} e^{-\beta \hat{\theta}_J} \left(\prod_{\substack{j \in J \\ k \in J^c}} v_a(x_k - x_j) \right)^{1/2}. \end{aligned} \quad (5.11)$$

This relation between the n -particle relativistic CM system introduced by Ruijsenaars and Macdonald's difference operators for the root system A_{n-1} was first observed by Koornwinder [K1]. It generalizes the relation between the n -particle Calogero-Sutherland system and the hypergeometric PDO's associated with $R = A_{n-1}$ [OP2] (cf. Remark *iii* at the end of Section 4.3).

iii. The operators $\hat{D}_{r, A_{n-1}}$ act as a real multiplication on functions that depend only on $x_1 + \dots + x_n$. The transition $\hat{D}'_r \longrightarrow \hat{D}_{r, A_{n-1}}$ can be interpreted physically as restricting attention to the motion in the center of mass hyperplane $x_1 + \dots + x_n = 0$.

We show next that the joint eigenfunctions of $\hat{D}_{r, A_{n-1}}$, i.e. Macdonald's A_{n-1} -type polynomials, can be obtained from Koornwinder's polynomials by a certain limit transition. Let $m_{\lambda, lead}$ be the sum of terms in m_{λ} (3.3) which are of the highest degree in $\exp(ix_j)$, $j = 1, \dots, n$:

$$m_{\lambda, lead} = \sum_{\lambda' \in S_n \lambda} e^{\lambda'}, \quad \lambda \in \mathcal{P}^+. \quad (5.12)$$

Recall that according to Definition 3.9, p_{λ} is a linear combination of monomials of the form

$$p_{\lambda} = \sum_{\lambda' \in \mathcal{P}^+, \lambda' \leq \lambda} c_{\lambda, \lambda'} m_{\lambda'}, \quad \lambda \in \mathcal{P}^+, \quad (5.13)$$

with $c_{\lambda, \lambda'}$ certain complex coefficients (which depend only on the parameters) such that (3.70) holds and $c_{\lambda, \lambda} = 1$. We set

$$p_{\lambda, lead} \equiv \sum_{\substack{\lambda' \in \mathcal{P}^+, \lambda' \leq \lambda \\ |\lambda'| = |\lambda|}} c_{\lambda, \lambda'} m_{\lambda', lead}, \quad \lambda \in \mathcal{P}^+. \quad (5.14)$$

Let

$$\omega \equiv e_1$$

From the asymptotics

$$m_{\lambda}(x - iR\omega) \sim m_{\lambda, lead}$$

and Eqs. (5.13) and (5.14), one derives

$$p_{\lambda, lead}(x) = \lim_{R \rightarrow \infty} m_{\lambda}(x - iR\omega)$$

The polynomial $p_{\lambda, lead}$ is homogeneous of degree $|\lambda|$. Consequently, a translation causes an automorphism of the space of polynomials. By multiplying $p_{\lambda, lead}$ by $\exp(iR|\lambda|)$, one obtains a basis of translation-invariant polynomials.

$$p_{\lambda, A_{n-1}}(x) \equiv e^{-i|\lambda|(x_1 + \dots + x_n)/n} p_{\lambda, lead}(x)$$

with

$$m_{\lambda', A_{n-1}}(x) \equiv e^{-i|\lambda'|x_1}$$

As the notation suggests, it will turn out that $p_{\lambda, A_{n-1}}$ are the Macdonald polynomials associated with the root system A_{n-1} .

First we need a lemma. It says that the p_{λ} are linearly independent. (Recall that the eigenvalues of $\hat{D}_1$ are given by p_{λ} .)

Lemma 5.1 *Let*

$$F_{\beta}(\theta) \equiv$$

Then, for all $\lambda, \lambda' \in \mathcal{P}^+$:

$$\lambda > \lambda' \implies F_{\beta}(\lambda) > F_{\beta}(\lambda')$$

(with ρ given by (3.35)).

Proof

It is clear from Definition 3.1 that $\lambda > \lambda'$ if and only if

$$\lambda = \lambda' + \sum_{1 \leq j \leq n-1} a_j (e_j - e_n)$$

with at least one of the a_j 's positive.

Obviously, it suffices to verify (5.21) for $\lambda > \lambda'$ positive. Now for $j = n$ this is immediate, since F_{β} is a convex function:

$$\text{ch}\beta(x + a) + \text{ch}\beta(x) > \text{ch}\beta(x)$$

if $x \geq y$ and $a > 0$.

□

Proposition 5.2 Let $\hat{D}_{r,A_{n-1}}$ be determined by (5.6)-(5.7) and let $p_{\lambda,A_{n-1}}$ be defined by (5.18). Then

$$\hat{D}_{r,A_{n-1}} p_{\lambda,A_{n-1}} = E_{r,A_{n-1}}(\lambda + \rho') p_{\lambda,A_{n-1}}, \quad (5.24)$$

with

$$E_{r,A_{n-1}}(\theta) \equiv e^{\beta r(\theta_1 + \dots + \theta_n)/n} S_r(e^{-\beta\theta_1}, \dots, e^{-\beta\theta_n}), \quad (5.25)$$

$$\rho'_j \equiv g(n+1-2j)/2, \quad 1 \leq j \leq n. \quad (5.26)$$

(S_r denotes the r th elementary symmetric function (Definition 2.2)).

Proof

The operator $\hat{D}'_r$ (5.6) is invariant both under permutations of x_j and under translations of the form $x \rightarrow x + R\omega$ (with ω as in (5.15)). We use this and the asymptotics of $(\hat{D}'_r m_{\lambda,lead})(-iRy)$ for $R \rightarrow \infty$ (with y such that (3.30) holds) to derive (cf. Proposition 3.3, Proposition 3.4, and their proofs)

$$\hat{D}'_r m_{\lambda,lead} = \sum_{\substack{\lambda' \in \mathcal{P}^+, \lambda' \leq \lambda \\ |\lambda'| = |\lambda|}} [\hat{D}'_r]_{\lambda,\lambda'} m_{\lambda',lead}, \quad (5.27)$$

with

$$[\hat{D}'_r]_{\lambda,\lambda} = S_r(e^{-\beta(\lambda_1 + \rho'_1)}, \dots, e^{-\beta(\lambda_n + \rho'_n)}). \quad (5.28)$$

(The poles at $x_j = x_k$, $j \neq k$, cancel because of the permutation symmetry; the condition $|\lambda'| = |\lambda|$ in sum the (5.27) stems from the translational invariance of $\hat{D}'_r$).

We will now show that $p_{\lambda,lead}$ is an eigenfunction of $\hat{D}'_r$ (with eigenvalue (5.28)). Consider the eigenvalue equation (3.72) for $r = 1$:

$$(\hat{D}_1 p_\lambda)(x) = 2 \left(\sum_{1 \leq j \leq n} [\text{ch}\beta(\lambda_j + \rho_j) - \text{ch}\beta\rho_j] \right) p_\lambda(x), \quad \lambda \in \mathcal{P}^+, \quad (5.29)$$

with $\hat{D}_1$ given by (2.10). Substitute

$$x \rightarrow x - iR\omega \quad (5.30)$$

and divide both sides of the equation by $\exp(R|\lambda|)$; Sending $R \rightarrow \infty$ entails (use (3.19),(3.20) and (5.17))

$$\begin{aligned} (c_1 \hat{D}'_1 + c_1^{-1} (\hat{D}'_n)^{-1} \hat{D}'_{n-1} - c_2) p_{\lambda,lead} = \\ 2 \left(\sum_{1 \leq j \leq n} [\text{ch}\beta(\lambda_j + \rho_j) - \text{ch}\beta\rho_j] \right) p_{\lambda,lead}, \end{aligned} \quad (5.31)$$

with

$$\begin{aligned} c_1 &= e^{-\beta g(n-1)/2} e^{-\beta(g_0 + g_1)} \\ c_2 &= c_1 \sum_{1 \leq j \leq n} \prod_{k \neq j} v_a(x_j - x_k) \\ &\stackrel{*}{=} 2 \sum_{1 \leq j \leq n} \text{ch}\beta\rho_j. \end{aligned}$$

(To verify equality $*$, first check that both sides are independent of x_j ; consequently, these parts are constants. Then check the value of these constants by putting $x = 0$ and sending $R \rightarrow \infty$).

It is clear from the commutativity of $\hat{D}'_1$ and $\hat{D}'_n$ on the l.h.s. of (5.31). Therefore, $\hat{D}'_r p_{\lambda,lead}$ corresponding to the same eigenvalue as $p_{\lambda,lead}$. $\hat{D}'_r p_{\lambda,lead}$ lies in $\text{span}\{p_{\lambda',lead} \mid \lambda' \leq \lambda, |\lambda'| = |\lambda|\}$ that $p_{\lambda,lead}$ must be an eigenfunction of $\hat{D}'_r$ (5.27), (5.28).

One obtains the expressions (5.24)-(5.26) from (5.27) and (5.28).

Notice that $m_{\lambda',A_{n-1}} = m_{\lambda,A_{n-1}}$ iff $\lambda' - \lambda \in \mathcal{P}^+$ can be relabeled by the projection of $\mathcal{P}^+$ onto $\mathcal{P}^+$. This projection of $\mathcal{P}^+$ coincides with the set of $\lambda' - \lambda \in \mathbb{Z}(e_1 + \dots + e_n)$. The polynomials $p_{\lambda,A_{n-1}}$ can also be relabeled by $\lambda' - \lambda \in \mathbb{Z}(e_1 + \dots + e_n)$. (This follows from (5.24) for $r = 1, \dots, n-1$). Since the operators $\hat{D}'_r$ are difference operators up to a constant, we can write

Corollary 5.3 The function $p_{\lambda,A_{n-1}}$ coincides with the function corresponding to the weight vector $\lambda - |\lambda|(e_1 + \dots + e_n)$ as in (2.21)).

Notes i. The transition $p_{\lambda,lead} \rightarrow p_{\lambda,A_{n-1}}$ is a shift of the center of mass.

ii. For $\beta \rightarrow 0$ (i.e. $q = \exp(-\beta) \rightarrow 1$), the polynomials associated with A_{n-1} [M2]. It follows from (5.14), (5.18) relates the Jacobi polynomial

$$p_{\lambda,A_{n-1}} = e^{-i|\lambda|(x_1 + \dots + x_n)/n} \lim_{R \rightarrow \infty} e^{iR|\lambda|} p_{\lambda,lead}$$

iii. Recently, a completely different limit of BC_n to those associated with A_{n-1} has been found with for $q \neq 1$.

B_n -type polynomials in $m_{\omega_n} \mathcal{A}^W$ can also be expressed in terms of Koornwinder's polynomials. To see this, notice that $m_{\omega_n}^2 \Delta_{(B_n, S)}$ coincides with a weight function of the type (3.50)-(3.53) with parameters

$$\mu_0 = \nu_1, \quad \mu_1 = i\beta, \quad \mu'_0 = \begin{cases} 0, & S = B_n \\ \nu_1, & S = C_n \end{cases}, \quad \mu'_1 = 0. \quad (5.49)$$

By multiplying m_{ω_n} and the Koornwinder polynomials with parameters as in (5.49), we obtain an orthogonal basis of $m_{\omega_n} \mathcal{A}^W$; the latter polynomials coincide with the anti-periodic Macdonald polynomials. To be more explicit, we have:

$$p_{\lambda, (B_n, S)} = p_{\lambda} \quad \begin{cases} \mu_0 = \nu_1, & \mu'_0 = \mu_1 = \mu'_1 = 0, & S = B_n \\ \mu_0 = \mu'_0 = \nu_1, & \mu_1 = \mu'_1 = 0, & S = C_n \end{cases} \quad (5.50)$$

$$p_{\lambda + \omega_n, (B_n, S)} = m_{\omega_n} p_{\lambda} \quad \begin{cases} \mu_0 = \nu_1, & \mu_1 = i\beta, & \mu'_0 = \mu'_1 = 0, & S = B_n \\ \mu_0 = \mu'_0 = \nu_1, & \mu_1 = i\beta, & \mu'_1 = 0, & S = C_n \end{cases} \quad (5.51)$$

(with $\lambda \in \mathcal{P}^+$).

Let us now turn to the corresponding difference operators. Let $\hat{D}_{1, (R, S)}, \dots, \hat{D}_{n, (R, S)}$ denote the operators $\hat{D}_1, \dots, \hat{D}_n$ (2.6) with parameters given by the table. We claim that the polynomials $p_{\lambda', (R, S)}$, $\lambda' \in \mathcal{P}_R^+$, are joint eigenfunctions of the operators $\hat{D}_{1, (R, S)}, \dots, \hat{D}_{n, (R, S)}$. For $R = (B)C_n$, and for the polynomials (5.50), this is an immediate consequence of Theorem 3.10. For the polynomials (5.51) this is seen as follows. By conjugating $\hat{D}_{1, (B_n, S)}$ with m_{ω_n} (5.47), one obtains (up to an additive constant) the operator $\hat{D}_1$ (2.10) with parameters (5.49):

$$\begin{aligned} m_{\omega_n}^{-1} \hat{D}_{1, (B_n, S)} m_{\omega_n} & \quad (5.52) \\ &= \sum_{\substack{1 \leq j \leq n \\ \varepsilon = \pm 1}} v_b(\varepsilon x_j) \prod_{k \neq j} v_a(\varepsilon x_j + x_k) v_a(\varepsilon x_j - x_k) \left(\frac{\cos \alpha(i\beta + \varepsilon x_j)}{\cos \alpha \varepsilon x_j} e^{-\varepsilon \beta \hat{\theta}_j} - 1 \right) \quad (\alpha = 1/2) \\ &= \sum_{\substack{1 \leq j \leq n \\ \varepsilon = \pm 1}} \frac{\cos \alpha(i\beta + \varepsilon x_j)}{\cos \alpha \varepsilon x_j} v_b(\varepsilon x_j) \prod_{k \neq j} v_a(\varepsilon x_j + x_k) v_a(\varepsilon x_j - x_k) \left(e^{-\varepsilon \beta \hat{\theta}_j} - 1 \right) + \text{const} \end{aligned}$$

(calculate the residues to verify the second equality). Therefore, the polynomials (5.51) are eigenfunctions of $\hat{D}_{1, (B_n, S)}$. One generalizes this to $\hat{D}_{r, (B_n, S)}$, $r > 1$, via similar reasoning as in the proof of Proposition 5.2: first one shows that $\hat{D}_{r, (B_n, S)}$ leaves invariant the space of anti-periodic polynomials $m_{\omega_n} \mathcal{A}^W$ (by calculating the residues); from the asymptotics for $\text{Im } x \rightarrow \infty$ in the positive Weyl chamber, it then follows that the operator is triangular. One uses Eq. (5.52), the monotony of the spectrum of $\hat{D}_{1, (B_n, S)}$ (Lemma 5.1), and the commutativity of the operators to conclude that the polynomials (5.51) are joint eigenfunctions of $\hat{D}_{1, (B_n, S)}, \dots, \hat{D}_{n, (B_n, S)}$.

The value of the additive constant in the r.h.s. of (5.52) can be easily obtained by comparing the spectrum of the operators on both sides of the equation (cf. eqs. (3.33)-(3.35), for $r = 1$, and (5.37)):

$$\text{const} = 2 \sum_{1 \leq j \leq n} (\text{ch} \beta(\rho_j + 1/2) - \text{ch} \beta \rho_j) \quad (5.53)$$

with $\rho_j = (n - j)g + k_1/2$ ($S = B_n$) or $\rho_j = (n - j)g + k_1/2 + 1/2$ ($S = C_n$). One could generalize (5.52) to an expression in terms of $\hat{D}_1, \dots, \hat{D}_r$ with parameters (5.49). The proof follows by comparing the spectrum of the operators $\hat{D}_{1, (R, S)}, \dots, \hat{D}_{r, (R, S)}$ (3.71) in terms of $E_{1, n}(\theta), \dots, E_{r, n}(\theta)$ with the spectrum of $\hat{D}_1, \dots, \hat{D}_r$ (2.10).

Remarks i. The operator $\hat{D}_{1, (R, S)}$ coincides with the Macdonald difference operator D_{π} that is associated with $\omega_1 = e_1$ of $S^{\vee}$. For technical reasons, Macdonald's operator is scaled by factor 2 if $S = C_n$ and $R = B(C)_n$. Specifically, $\hat{D}_{1, (R, S)} \sim D_{2e_1}$ if $S = C_n$ and $R = B(C)_n$. *ii.* The operators $\hat{D}_{1, (R, S)}, \dots, \hat{D}_{n, (R, S)}$ are simultaneous eigenoperators that are simultaneous eigenfunctions of $S = C_n$ and consider the operator

$$\hat{D}'_{n, (R, C_n)} \equiv \sum_{\varepsilon_1, \dots, \varepsilon_n = \pm 1} \prod_{1 \leq j \leq n} v'_b(\varepsilon_j x_j)$$

with

$$v'_b(z) \equiv \frac{\sin \alpha(\nu_1 + z)}{\sin \alpha(\nu_2 + z)}$$

This AΔO coincides up to a multiplicative constant with the operator associated with the n th fundamental weight ω_n . One has (cf. Remark i) $\hat{D}'_{n, (R, C_n)} \sim E_{\omega_n}$ if $R = B(C)_n$.

It follows from [M2] that $\hat{D}'_{n, (R, C_n)}$ is diagonalizable. However, that $\hat{D}'_{n, (R, C_n)}$ is not in $\mathbb{D}$ but its square is. This follows from eigenvalues and using Theorem 3.12).

iii. From a group-theoretic perspective, $\hat{D}'_{n, (R, C_n)}$ is isomorphic to $\mathbb{R}[\mathcal{P}]^W$ (the W-invariant polynomials on $\mathcal{P} = \mathbb{Z}^n$). If $S = B_n$, then one has $\mathcal{P}_{S^{\vee}} = \mathcal{P}$. If $S = C_n$, $\mathcal{P}_{S^{\vee}}$ is a subgroup of $\mathcal{P}_{S^{\vee}}$ with index two; so $\mathcal{P}_{S^{\vee}} \neq \mathcal{P}$. In the latter case, one can extend $\mathbb{D}$ to an algebra by replacing the generator $\hat{D}_{n, (R, C_n)}$ by $\hat{D}'_{n, (R, C_n)}$.

5.4 The Root System D_n

We conclude by briefly sketching the state of the art (this should not have much difficulty to supply more details in a subsection). Put

$$\mu_0 = \mu_1 = \dots = \mu_n = 0$$

Then $d_b(z) = 1$ and $\Delta(x)$ (3.50) reduces to $\Delta(x) = 1$. The correspondence of parameters is via Elmacdonald's operator D_{ω_1} associated with

For $R = D_n$, the torus $\mathbb{T}_R = \mathbb{R}^n / (2\pi\mathbb{Z}R^\vee)$ is the same as for $R = B_n$. The Weyl group, however, is smaller: only an even number of sign flips of the variables x_j , $j = 1, \dots, n$, is allowed. For $k = 1, \dots, n-2$, the fundamental weights ω_k of D_n are the same as in (5.42), but ω_{n-1} and ω_n are now given by the half-spin weights

$$\omega_{n-1} = (e_1 + \dots + e_{n-1} - e_n)/2, \quad \omega_n = (e_1 + \dots + e_{n-1} + e_n)/2. \quad (5.57)$$

It is not hard to see that the cone of dominant weights $\mathcal{P}_{D_n}^+$ generated by ω_k , $k = 1, \dots, n$, consists of the vectors

$$(\lambda + \delta\omega_n)_\varepsilon \equiv (\lambda_1 + \delta/2, \dots, \lambda_{n-1} + \delta/2, \varepsilon(\lambda_n + \delta/2)) \quad (5.58)$$

with

$$\lambda \in \mathcal{P}^+, \quad \delta = 0, 1, \quad \varepsilon = \pm 1. \quad (5.59)$$

The Macdonald polynomials p_{λ', D_n} , $\lambda' \in \mathcal{P}_{D_n}^+$ form an orthogonal basis of $L_{W_{D_n}}^2(\mathbb{T}_{D_n}, \Delta_{D_n} dx)$. By combining the polynomials associated with $(\lambda + \delta\omega_n)_+$ and $(\lambda + \delta\omega_n)_-$ one obtains polynomials that are even in x_j , $j = 1, \dots, n$. These are related to Koornwinder's polynomials in the following way (cf. Eqs. (5.50), (5.51)):

$$\begin{aligned} p_{\lambda_+, D_n} &= p_{\lambda_-, D_n} = p_\lambda, & \mu_0 &= \mu_1 = \mu'_0 = \mu'_1 = 0, & \text{if } \lambda_n &= 0, \\ p_{\lambda_+, D_n} + p_{\lambda_-, D_n} &= p_\lambda, & \mu_0 &= \mu_1 = \mu'_0 = \mu'_1 = 0, & \text{if } \lambda_n &> 0, \\ p_{(\lambda+\omega_n)_+, D_n} + p_{(\lambda+\omega_n)_-, D_n} &= m_{\omega_n} p_\lambda, & \mu_0 &= \mu'_0 = \mu'_1 = 0, \mu_1 = i\beta. \end{aligned} \quad (5.60)$$

In the second and the third line of the above formula one obtains a sum of D_n polynomials rather than the polynomials themselves. Nevertheless, Eq. (5.60) determines the D_n polynomials uniquely. This is because flipping the sign of one of the x_j 's in $p_{(\lambda+\delta\omega_n)_+, D_n}$ results in $p_{(\lambda+\delta\omega_n)_-, D_n}$. Consequently, the coefficients of the expansion of $p_{(\lambda+\delta\omega_n)_\varepsilon, D_n}$ in D_n -type monomial symmetric functions are determined in terms of the coefficients occurring in (3.69).

As regards the difference operators with parameters (5.56), the algebra $\mathbb{D}$ consists of commuting $\Lambda\Delta\mathcal{O}$'s with the D_n -type polynomials as joint eigenfunctions. One can extend $\mathbb{D}$ to an algebra that is isomorphic to $\mathbb{R}[\mathcal{P}_{D_n}^\vee]^{W_{D_n}}$ by replacing the generators $\hat{D}_{n-1}$ and $\hat{D}_n$ by

$$\hat{D}'_{n-1} = \sum_{\substack{\varepsilon_1, \dots, \varepsilon_n = \pm 1 \\ \varepsilon_1 \dots \varepsilon_n = -1}} \prod_{1 \leq j < k \leq n} v_a(\varepsilon_j x_j + \varepsilon_k x_k) e^{-\beta(\varepsilon_1 \hat{\theta}_1 + \dots + \varepsilon_n \hat{\theta}_n)/2}, \quad (5.61)$$

$$\hat{D}'_n = \sum_{\substack{\varepsilon_1, \dots, \varepsilon_n = \pm 1 \\ \varepsilon_1 \dots \varepsilon_n = +1}} \prod_{1 \leq j < k \leq n} v_a(\varepsilon_j x_j + \varepsilon_k x_k) e^{-\beta(\varepsilon_1 \hat{\theta}_1 + \dots + \varepsilon_n \hat{\theta}_n)/2}. \quad (5.62)$$

These operators are proportional to Macdonald's operators $E_{\omega_{n-1}}$ and E_{ω_n} , which are associated with the half-spin weights (5.57).

Appendix A: Cancellation of Poles

In this appendix we prove two results, which were needed to demonstrate that $\hat{D}_r$ maps $\mathcal{A}^W$ into itself. It was claimed in the proof of Proposition 3.3 (Section 3.2) that the

following expression:

$$V_J^1 V_{J^c}^3 \sum_{\substack{\emptyset \subsetneq J_1 \subsetneq \dots \subsetneq J_s = J \\ 1 \leq s \leq |J|}} (-1)^s \prod_{1 \leq s' \leq s} V_{J_{s'} \setminus J_s}^2$$

(with $\lambda \in \mathcal{P}^+$ and $J \subset \{1, \dots, n\}$, $J_0 \equiv \emptyset$, and at $x_1 = -x_j - 2\gamma$, $j = 2, \dots, n$ (poles of the coefficients) to zeros in the denominators of the coefficients) that we assume that the parameters γ , μ , μ_δ , μ'_δ are chosen in such a way that these poles are cancelled by the above regularity claims, thereby completing the proof.

Before turning to the details, let us outline the structure of a sum of terms of the type

$$(-1)^s V_J^1 V_{J^c}^3 \prod_{1 \leq s' \leq s} V_{B_{s'}}^2 V_{B_{s'}^c}^3$$

where the index sets $B_{s'} \subset J$, $s' = 1, \dots, s$

$$B_{s'} \equiv J_{s'} \setminus J_{s' \setminus 1}$$

The terms (A.2) are associated with the sequence

$$\emptyset \subsetneq J_1 \subsetneq J_2 \subsetneq \dots \subsetneq J_s$$

(with the cell J fixed). Each term in (A.2) can be obtained by constructing an involutive operation σ ($\sigma^2 = \text{id}$) on the sequence $J_1, \dots, J_s$ in a way that the terms associated with a sequence and its image cancel each other in the residue. Therefore, the poles in (A.1) cancel.

Lemma A.1 (pole of type I)

Let γ , μ , μ_δ , μ'_δ ($\delta = 0, 1$) and $x_2, \dots, x_n$ be generic parameters and poles. Then (A.1) is regular as a function of x_1 .

Proof

First note that the lemma is trivial if $1 \notin J$. If $1 \in J$, then V_J^1 gives rise to a pole at x_1 .

Assume $1 \in J$ and let B_{s_1} denote the block of J containing 1. Define the following map σ on the collection of sequences (A.2):

1.A If $|B_{s_1}| > 1$, then σ maps (A.4) to the sequence

$$\emptyset \subsetneq J_1 \subsetneq J_2 \subsetneq \dots \subsetneq J_{s_1-1}$$

1.B If $|B_{s_1}| = 1$ and $s_1 > 1$, then σ maps (A.4) to the sequence

$$\emptyset \subsetneq J_1 \subsetneq J_2 \subsetneq \dots \subsetneq J_{s_1}$$

2. If $|B_{s_1}| = 1$ and $s_1 = 1$ (i.e. $J_1 = \{1\}$) then σ maps (A.4) to the sequence

Phrased in words: unless $B_{s_1} = \{1\}$, the map σ pulls the index 1 out of B_{s_1} and places it in a newly created block, which is sandwiched between $B_{s_1} \setminus \{1\}$ and B_{s_1+1} (case 1.A); when B_{s_1} contains only the index 1, then σ merges the blocks $B_{s_1} = \{1\}$ and B_{s_1-1} if $s_1 > 1$ (case 1.B) or, if $s_1 = 1$, then it leaves the sequence (A.4) unchanged (case 2.).

Thus defined, σ is indeed an involution on the collection of sequences (A.4): the cases 1.A and 1.B are inverse to each other (see Fig. 1. below).




Fig. 1. A graphical representation of the map σ .

We claim that in the first situation (i.e. 1.A or 1.B) the pole at $x_1 = -\gamma$ in the term (A.2) (which is associated with (A.4)) cancels against the pole in the term corresponding with the σ -image of the sequence (A.4). To see this, we may assume that we are in situation 1.A. One obtains the term corresponding to the sequence (A.5) from (A.2) by making the substitutions

$$s \rightarrow s + 1, \quad (\text{A.7})$$

$$V_{B_{s_1}}^2 \rightarrow V_{B_{s_1} \setminus \{1\}}^2 V_{\{1\}}^2 = V_{B_{s_1} \setminus \{1\}}^2, \quad (\text{A.8})$$

$$\begin{aligned} V_{B_{s_1}; J \setminus J_{s_1}}^3 &\rightarrow V_{B_{s_1} \setminus \{1\}; (J \setminus J_{s_1}) \cup \{1\}}^3 V_{\{1\}; J \setminus J_{s_1}}^3 \\ &= V_{B_{s_1}; J \setminus J_{s_1}}^3 V_{B_{s_1} \setminus \{1\}; \{1\}}^3, \end{aligned} \quad (\text{A.9})$$

$$m_\lambda(x + 2\gamma e_{B_1}) \rightarrow m_\lambda(x + 2\gamma e_{B_1} - 2\gamma \delta_{1, s_1} e_1) \quad (\text{A.10})$$

($\delta_{j,k}$ denotes the Kronecker symbol). This substitution in (A.2) amounts to replacing the part

$$\begin{aligned} &\left(V_{B_{s_1}}^2 / V_{B_{s_1} \setminus \{1\}}^2 \right) [m_\lambda(x + 2\gamma e_{B_1}) - m_\lambda(x)] = \\ &\prod_{j \in B_{s_1} \setminus \{1\}} v_a(x_j + x_1) v_a(x_j + x_1 + 2\gamma) [m_\lambda(x + 2\gamma e_{B_1}) - m_\lambda(x)] \end{aligned} \quad (\text{A.11})$$

by

$$\begin{aligned} &-V_{B_{s_1} \setminus \{1\}; \{1\}}^3 [m_\lambda(x + 2\gamma e_{B_1} - 2\gamma \delta_{1, s_1} e_1) - m_\lambda(x)] = \\ &= - \prod_{j \in B_{s_1} \setminus \{1\}} v_a(x_j + x_1) v_a(x_j - x_1) [m_\lambda(x + 2\gamma e_{B_1} - 2\gamma \delta_{1, s_1} e_1) - m_\lambda(x)]. \end{aligned} \quad (\text{A.12})$$

At $x_1 = -\gamma$ the r.h.s. of (A.11) and (A.12) cancel.

If we are in situation 2., i.e. $B_1 = \{1\}$, then in V_J^1 is compensated by a zero in the difference.

$$[m_\lambda(x + 2\gamma e_1) - m_\lambda(x)]$$

This shows that the total residue at $x_1 = -\gamma$ is zero, completing the proof of the lemma.

Lemma A.2 (poles of type II)

Let $\gamma, \mu, \mu_\delta, \mu'_\delta$ ($\delta = 0, 1$) and $x_2, \dots, x_n$ be poles. Then (A.1) is regular as a function of x_1 .

Proof

The proof is very similar to that of Lemma A.1. If J does not contain the pair $\{1, j\}$, because $x_1 = -x_j - 2\gamma$.

Assume for the remaining part of the proof that J contains the pair $\{1, j\}$. We consider two cases: 1.A and 1.B.

1.A If the pair $\{1, j\}$ is contained in one of the blocks J_{s_j} with $|B_{s_j}| > 2$, then σ_j maps sequence (A.4) to

$$\emptyset \subsetneq J_1 \subsetneq \dots \subsetneq J_{s_j-1} \subsetneq J_{s_j} \subsetneq \dots \subsetneq J_n$$

1.B If one of the blocks of sequence (A.4) contains the pair $\{1, j\}$, then σ_j maps sequence (A.4) to

$$\emptyset \subsetneq J_1 \subsetneq \dots \subsetneq J_{s_j} \subsetneq \dots \subsetneq J_n$$

2. If $B_1 = \{1, j\}$, then σ_j maps sequence (A.4) to

3. If the pair $\{1, j\}$ is not contained in any of the blocks of sequence (A.4) to itself.

It is clear that σ_j is an involution, the cases 1.A and 1.B are inverse to each other (see Fig. 2. below).




Fig. 2. A graphical representation of the map σ_j .

Consider situation 1., assuming case 1.A. The application of σ_j boils down to making the following substitutions in the associated term (A.2):

$$s \rightarrow s + 1 \quad (\text{A.16})$$

$$V_{B_{s_j}}^2 \rightarrow V_{B_{s_j} \setminus \{1, j\}}^2 V_{\{1, j\}}^2 \quad (\text{A.17})$$

$$V_{B_{s_j}; J \setminus J_{s_j}}^3 \rightarrow V_{B_{s_j}; J \setminus J_{s_j}}^3 V_{B_{s_j}; \{1, j\}}^3. \quad (\text{A.18})$$

These substitutions amount to the following change in the term (A.2): replace the part

$$\prod_{\substack{k \in B_{s_j} \\ k \neq 1, j}} v_a(x_k + x_1) v_a(x_k + x_1 + 2\gamma) v_a(x_k + x_j) v_a(x_k + x_j + 2\gamma) \quad (\text{A.19})$$

by

$$- \prod_{\substack{k \in B_{s_j} \\ k \neq 1, j}} v_a(x_k + x_1) v_a(x_k - x_1) v_a(x_k + x_j) v_a(x_k - x_j). \quad (\text{A.20})$$

At $x_1 = -x_j - 2\gamma$, (A.19) and (A.20) differ only by sign. Consequently, the residues at $x_1 = -x_j - 2\gamma$ of the corresponding terms in (A.1) add up to zero.

In situation 2. the pole in the coefficient of (A.2), which is caused by $V_{B_1}^2$, cancels against the zero in the difference of the monomial symmetric functions:

$$[m_\lambda(x + 2\gamma e_{\{1, j\}}) - m_\lambda(x)]_{(x_1 = -x_j - 2\gamma)} = 0. \quad (\text{A.21})$$

In situation 3. the denominator of the coefficient of (A.2) has no zero at $x_1 = -x_j - 2\gamma$, so the term is regular at $x_1 = -x_j - 2\gamma$.

We conclude from the above analysis that the total residue at $x_1 = -x_j - 2\gamma$ in the sum (A.1) is zero, thus completing the proof of the lemma. $\square$

Appendix B: Two Comb

In this appendix we prove two technical results on the eigenvalues of our difference operator; its solution resulted in explicit formulae. Lemma B.2 deals with a recursion relation, and Lemma B.3 deals with a relation concerning the behavior of the eigenvalues.

Lemma B.1 *The functions*

$$F_{m,p} = (-1)^p \sum_{1 \leq i_1 \leq \dots \leq i_p \leq m-p}$$

form the unique solution of the linear system

$$\sum_{\substack{J \subset \{1, \dots, n\}, |J|=s \\ 0 \leq s \leq r}} \left(\prod_{j \in J} t_j \right) F_{m,p}$$

with the convention

$$F_{m,0} = 1,$$

Proof

After splitting off the term in (B.2) corresponding to the r.h.s. of the equation, one arrives at

$$F_{n,r} = - \sum_{\substack{J \subset \{1, \dots, n\}, |J|=s \\ 1 \leq s \leq r}} \left(\prod_{j \in J} t_j \right) F_{n,r}$$

It is clear that (B.4) with condition (B.3) holds. Hence, the system (B.2), (B.3) has a unique solution.

In order to prove that this solution is indeed the expression

$$\sum_{\substack{J \subset \{1, \dots, n\}, |J|=s \\ 0 \leq s \leq r}} (-1)^s \left(\prod_{j \in J} t_j \right) \left(\prod_{j \in J} t_j \right)$$

vanishes identically (for $1 \leq r \leq n$). To this end, we consider the sum of monomials in the variables $t_1, \dots, t_n$ of the form

$$(-1)^s (t_{j_1} t_{j_2} \cdots t_{j_s})$$

which correspond to pairs of the form

$$\{(j_1, j_2, \dots, j_s), (i_1, i_2, \dots, i_s)\}$$

subject to the condition

$$1 \leq j_1 < j_2 < \cdots < j_s \leq n, \quad 1 \leq i_1 \leq i_2 \leq \cdots \leq i_{r-s} \leq n - r + 1. \quad (\text{B.8})$$

We shall now show that these monomials cancel in pairs.

Let σ be the following operation defined on the above collection of pairs (B.7) with condition (B.8):

A. If $i_1 < j_1$ or $s = 0$, then

$$\{(j_1, j_2, \dots, j_s), (i_1, i_2, \dots, i_{r-s})\} \xrightarrow{\sigma} \{(i_1, j_1, \dots, j_s), (i_2, \dots, i_{r-s})\}, \quad (\text{B.9})$$

B. If $i_1 \geq j_1$ or $s = r$, then

$$\{(j_1, j_2, \dots, j_s), (i_1, i_2, \dots, i_{r-s})\} \xrightarrow{\sigma} \{(j_2, \dots, j_s), (j_1, i_1, \dots, i_{r-s})\}. \quad (\text{B.10})$$

Roughly speaking, σ compares the first entries of the two elements constituting the pair (B.7) and moves the smallest of these two to the first entry of the other element. One easily verifies that: first, σ is well defined in the sense that the image of (B.7) is again a pair satisfying (B.8); second, σ is an involution ($\sigma^2 = id$), the cases A. and B. being inverse to each other.

For the associated monomial (B.6), acting with σ amounts to an increase (case A.) or a decrease (case B.) of the number s by one, i.e. it flips the sign of the corresponding monomial. Therefore, combining the term (B.6) associated with a pair (B.7) with the one associated with its image under σ entails the vanishing of the sum (B.5), which completes the proof. $\square$

Remark. If one replaces the upper bound $n - r + 1$ of the second summation in (B.5) by n , then, for $r = n$, expression (B.5) also vanishes. (Indeed, the above proof again applies). In this case the vanishing of (B.5) amounts to a well-known relation between the elementary symmetric functions and the complete symmetric functions (see e.g. [M1]).

Lemma B.2 *The function*

$$E_{r,n}(t_1, \dots, t_n; p_r, \dots, p_n) = \sum_{0 \leq s \leq r} (-1)^{r+s} \left(\sum_{1 \leq j_1 < \cdots < j_s \leq n} t_{j_1} \cdots t_{j_s} \right) \left(\sum_{r \leq i_1 \leq \cdots \leq i_{r-s} \leq n} p_{i_1} \cdots p_{i_{r-s}} \right), \quad 1 \leq r \leq n, \quad (\text{B.11})$$

is the unique solution of the recursion relation

$$E_{r,n}(t_1, \dots, t_n; p_r, \dots, p_n) = (t_n - p_n) E_{r-1,n-1}(t_1, \dots, t_{n-1}; p_r, \dots, p_n) + E_{r,n-1}(t_1, \dots, t_{n-1}; p_r, \dots, p_{n-1}), \quad 1 \leq r \leq n \quad (\text{B.12})$$

with the convention

$$E_{0,n} \equiv 1, \quad E_{r,n} \equiv 0 \text{ if } n < r. \quad (\text{B.13})$$

Proof

It is clear that (B.12) with condition (B.13) is satisfied. After splitting up the sum in (B.11) into two parts, one indeed solves Eq. (B.12):

- i. terms with $j_s = n$;
- ii. terms with $j_s < n$ and $i_{r-s} = n$;
- iii. terms with $j_s < n$ and $i_{r-s} < n$;

Remark. In some cases Lemma B.2 can be used to prove that the sum (B.5) is zero. For instance, one easily verifies with the aid of (B.13) that

$$E_{r,n}(t_1, \dots, t_n; p_r, p_{r+1}, \dots, p_n) = 0$$

then

$$E_{r,n} = \sum_{\substack{J \subset \{1, \dots, n\} \\ |J|=r}} \left(\prod_{j \in J} (t_j - p_j) \right) E_{r,n}(t_1, \dots, t_n; p_r, \dots, p_n)$$

In particular, $E_{n,n}(t_1, \dots, t_n; p_n) = (t_1 - p_n) \cdots (t_n - p_n)$.

Appendix C: $\hat{D}(\mathcal{A}^W) = 0$

In this appendix we present a result due to Heckman [He84]: if an AΔO or PDO is zero on all symmetric functions, then it is zero. This fact was used in Section 3.5 to prove (Theorem 3.11), and again in Section 4.3 to prove that the BC_n -type hypergeometric PDO's of Heckman [He84] are zero. Let

$$\hat{T}_\kappa \equiv \hat{t}_1^{\kappa_1} \cdots \hat{t}_n^{\kappa_n}, \quad \hat{t}_j \equiv t_j$$

with $\kappa = (\kappa_1, \dots, \kappa_n)$ in $\mathbb{R}^n$ or $\mathbb{N}^n$ in which case the AΔO's/PDO's of interest are of the form:

$$\hat{D} = \sum_{1 \leq r \leq n} \hat{t}_r^{\kappa_r} \hat{D}_r$$

with the n -dimensional vectors $\kappa^{(1)}, \dots, \kappa^{(n)}$ in $\mathbb{R}^n$.

$$V_r : \mathcal{U} \subset \mathbb{R}^n \longrightarrow \mathbb{R}$$

continuous on an open dense set $\mathcal{U} \subset \mathbb{R}^n$.

Proposition C.1 ($\hat{D}(\mathcal{A}^W) = 0 \Rightarrow \hat{D} = 0$)
Let $\hat{D}$ be an AΔO/PDO of the form (C.2).

$$\hat{D} m_\lambda = 0,$$

then

$$V_1(x) = V_2(x) = \cdots = 0$$

Proof ([R3])

Introduce the following vector-valued functions:

$$t_\lambda : \mathcal{U} \rightarrow \mathbb{C}^M, \quad x \xrightarrow{t_\lambda} ((\hat{T}_{\kappa(1)} m_\lambda)(x), \dots, (\hat{T}_{\kappa(M)} m_\lambda)(x)), \quad \lambda \in \mathcal{P}^+, \quad (\text{C.6})$$

$$v : \mathcal{U} \rightarrow \mathbb{C}^M, \quad x \xrightarrow{v} (\overline{V_1(x)}, \dots, \overline{V_M(x)}). \quad (\text{C.7})$$

The fact that m_λ is in the kernel of $\hat{D}$ translates itself geometrically in the orthogonality of t_λ and v :

$$\hat{D} m_\lambda = 0 \iff (t_\lambda, v) = 0. \quad (\text{C.8})$$

We will assume $v \not\equiv 0$ and derive a contradiction. Let $\lambda^{(1)}, \dots, \lambda^{(M)}$ be vectors in $\mathcal{P}^+$. One has $v \perp t_{\lambda^{(1)}}, \dots, t_{\lambda^{(M)}}$. Therefore, the vectors $t_{\lambda^{(1)}}(x), \dots, t_{\lambda^{(M)}}(x)$ must be linearly dependent for all $x \in \mathcal{U}$ for which $v(x) \neq 0$. Since $v(x)$ is continuous in x , there exists an open ball $\mathcal{B} \subset \mathcal{U}$ on which $v(x) \neq 0$. The fact that the vectors $t_{\lambda^{(s)}}(x)$, $1 \leq s \leq M$ are real-analytic in x then entails

$$\det(\text{Col}[t_{\lambda^{(1)}}(x), \dots, t_{\lambda^{(M)}}(x)]) = 0, \quad \forall x \in \mathcal{U}. \quad (\text{C.9})$$

We will now show that an appropriate choice of the vectors $\lambda^{(1)}, \dots, \lambda^{(M)}$ contradicts the vanishing of the above determinant.

Let $\lambda \in \mathcal{P}^+$ and $y \in \mathbb{R}^n$ be such that

$$\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_n > 0, \quad y_1 > y_2 > \dots > y_n > 0. \quad (\text{C.10})$$

From the asymptotics (cf. Eq. (3.37))

$$(\hat{T}_\kappa m_\lambda)(x)|_{x=iRy} \sim \tau_{\kappa, \lambda} e^{(\lambda, y)R}, \quad R \rightarrow \infty, \quad (\text{C.11})$$

with

$$\tau_{\kappa, \lambda} = \begin{cases} e^{\beta(\kappa, \lambda)} & (\text{A}\Delta\text{O}) \\ (-1)^{|\kappa|} (\lambda_1)^{\kappa_1} \dots (\lambda_n)^{\kappa_n} & (\text{PDO}) \end{cases}, \quad (\text{C.12})$$

one derives:

$$\lim_{R \rightarrow \infty} e^{i(\lambda, x)} t_\lambda(x)|_{x=iRy} = (\tau_{\kappa(1), \lambda}, \dots, \tau_{\kappa(M), \lambda}). \quad (\text{C.13})$$

Pick the vector λ (subject to condition (C.10)) in such a way that

$$\tau_{\kappa(r), \lambda} \neq \tau_{\kappa(p), \lambda}, \quad 1 \leq r < p \leq M. \quad (\text{C.14})$$

That such a λ exists follows in the $\text{A}\Delta\text{O}$ case from (C.12) and the fact that the vectors $\kappa^{(1)}, \dots, \kappa^{(M)}$ are distinct; in the PDO case one can pick distinct prime numbers for the components of λ .

We use $\lambda = (\lambda_1, \dots, \lambda_n)$ to form the vectors $\lambda^{(1)}, \dots, \lambda^{(M)}$ in the following way:

$$\lambda^{(s)} = \begin{cases} (s-1)(\lambda_1, \dots, \lambda_n) & (\text{A}\Delta\text{O}) \\ ((\lambda_1)^{s-1}, \dots, (\lambda_n)^{s-1}) & (\text{PDO}) \end{cases}, \quad 1 \leq s \leq M. \quad (\text{C.15})$$

On the one hand, Eqs. (C.9) and (C.13) imply

$$\tau \equiv \begin{vmatrix} \tau_{\kappa(1), \lambda^{(1)}} & \dots & \tau_{\kappa(1), \lambda^{(M)}} \\ \vdots & \ddots & \vdots \\ \tau_{\kappa(M), \lambda^{(1)}} & \dots & \tau_{\kappa(M), \lambda^{(M)}} \end{vmatrix} = 0. \quad (\text{C.16})$$

On the other hand, for the above choice of determinant:

$$\tau_{\kappa(r), \lambda^{(s)}} = (\tau_{\kappa(r), \lambda})^{s-1}$$

Therefore,

$$\tau = \prod_{1 \leq r < p \leq M} (\tau_{\kappa(r), \lambda} - \tau_{\kappa(p), \lambda})$$

Because λ is chosen such that $\tau_{\kappa(r), \lambda} \neq \tau_{\kappa(p), \lambda}$, determinant $\tau \neq 0$, contradicting (C.16).

Hence, v (C.7) must be zero.

Acknowledgments. The author would like to thank for helpful conversations and useful suggestions T. H. Koornwinder. Thanks are also due to T. H. Koornwinder.

References

- [AW] Askey, R., Wilson, J.: *Some basic hypergeometric series*. Amer. Math. Soc. **319** (1985)
- [BK] Beerends, R. J., Koornwinder, T. H.: *On the Jacobi function of BC_n* . In preparation
- [B] Bourbaki, N.: *Groupes et algèbres de Lie*. Hermann, Paris (1968)
- [C] Calogero, F.: *Solutions of the one-body problems and/or inversely quadratic pair potentials*. J. Math. Phys. **9**, 61-71 (1968)
- [Ch1] Cherednik, I.: *Quantum Knizhnik-Polyakov-Bershteyn equations*. Commun. Math. Phys. **150**, 117-148 (1993)
- [Ch2] Cherednik, I.: *Double affine Hecke algebras and Macdonald's operators*. Int. Math. Res. Not. **1995**, 1-40 (1995)
- [D] van Diejen, J. F.: *Difference Calogero-Moser systems*. In preparation
- [HO] Heckman, G. J., Opdam, E. M.: *Root systems and hypergeometric functions*. Compos. Math. **64**, 329-352 (1987)
- [H1] Heckman, G. J.: *Root systems and hypergeometric functions*. Adv. Math. **64**, 353-373 (1987)
- [H2] Heckman, G. J.: *An elementary approach to the theory of root systems*. Opdam. Invent. Math. **103**, 341-350 (1989)
- [I] Inozemtsev, V. I.: *Lax representations of integrable particle systems*. Lett. Math. Phys. **38**, 1-10 (1995)
- [K1] Koornwinder, T. H.: private notes (1995)
- [K2] Koornwinder, T. H.: *Jacobi function of BC_n* . J. Math. Anal. and Appl. **148**, 44-55 (1990)

- [K3] Koornwinder, T. H.: *Orthogonal polynomials in connection with quantum groups*. In: Nevai, P. (ed.), *Orthogonal polynomials: theory and practice*. NATO ASI Series C 294, pp. 257-292. Dordrecht: Kluwer Academic Publishers 1990
- [K4] Koornwinder, T. H.: *Askey-Wilson polynomials for root systems of type BC*. In: Richards, D. St. P. (ed.), *Hypergeometric functions on domains of positivity, Jack polynomials, and applications*. Contemp. Math. **138**, pp. 189-204 (1992)
- [M1] Macdonald, I. G.: *Symmetric functions and Hall polynomials*. Oxford: Clarendon Press 1979
- [M2] Macdonald, I. G.: *Orthogonal polynomials associated with root systems*. Oxford preprint (1988)
- [M3] Macdonald, I. G.: *A new class of symmetric functions*. In: Cerlienco, L., Foata, D. (eds.), *Actes 20^e Séminaire Lotharingien*, pp. 131-171. Strasbourg: Publ. I. R. M. A. 1988
- [M4] Macdonald, I. G.: *Orthogonal polynomials associated with root systems*. In: Nevai, P. (ed.), *Orthogonal polynomials: theory and practice*. NATO ASI Series C 294, pp. 311-318. Dordrecht: Kluwer Academic Publishers 1990
- [Mo] Moser, J.: *Three integrable Hamiltonian systems connected with isospectral deformations*. Adv. Math. **16**, 197-220 (1975)
- [N] Noumi, M.: *Macdonald's symmetric polynomials as zonal spherical functions on some quantum homogeneous spaces*. Informal summary, March 1992
- [O1] Opdam, E. M.: *Root systems and hypergeometric functions III*. Compos. Math. **67**, 21-49 (1988)
- [O2] Opdam, E. M.: *Root systems and hypergeometric functions IV*. Compos. Math. **67**, 191-209 (1988)
- [OP1] Olshanetsky, M. A., Perelomov, A. M.: *Classical integrable finite-dimensional systems related to Lie algebras*. Phys. Reps. **71**, 313-400 (1981)
- [OP2] Olshanetsky, M. A., Perelomov, A. M.: *Quantum integrable systems related to Lie algebras*. Phys. Reps. **94**, 313-404 (1983)
- [RS] Ruijsenaars, S. N. M., Schneider, H.: *A new class of integrable systems and its relation to solitons*. Ann. Phys. (N.Y.) **170**, 370-405 (1986)
- [R1] Ruijsenaars, S. N. M.: *Complete integrability of relativistic Calogero-Moser systems and elliptic function identities*. Commun. Math. Phys. **110**, 191-213 (1987)
- [R2] Ruijsenaars, S. N. M.: *Finite-dimensional soliton systems*. In: Kupershmidt, B. (ed.), *Integrable and superintegrable systems*. pp. 165-206. Singapore: World Scientific 1990
- [R3] Ruijsenaars, S. N. M.: private notes (1993)
- [S] Serre, J. P.: *Complex semisimple Lie algebras*. New York: Springer 1987
- [Su1] Sutherland, B.: *Exact results for a quantum many-body problem in one dimension*. Phys. Rev. **A4**, 2019-2021 (1971)
- [Su2] Sutherland, B.: *Exact results for a quantum many-body problem in one dimension II*. Phys. Rev. **A5**, 1372-1376 (1972)