arXiv:hep-th/0611348v1 30 Nov 2006

Topological Charge and the Laminar Structure of the QCD Vacuum

H. B. Thacker

University of Virginia, Charlottesville, VA 22904

Abstract. Monte Carlo studies of pure glue $SU(3)$ gauge theory using the overlap-based topological charge operator have revealed a laminar structure in the QCD vacuum consisting of extended, thin, coherent, locally 3-dimensional sheets of topological charge embedded in 4D space, with opposite sign sheets interleaved. In this talk I discuss the interpretation of these Monte Carlo results in terms of our current theoretical understanding of theta-dependence and topological structure in asymptotically free gauge theories.

Keywords: QCD, topological charge, D-branes
PACS: 11.15.Ha, 11.30.Rd, 12.38.Gc, 12.39.Fe

Recent studies of topological charge using the overlap-based topological charge density operator on the lattice have revealed a type of long-range structure that is profoundly different from what might have been expected in an instanton-based model of the QCD vacuum. These studies [1] (see also [2]) produced the surprising result that the $q(x)$ distribution in a typical Monte Carlo gauge configuration is dominated by extended, coherent, thin 3-dimensional sheets of topological charge. In each configuration, sheets of opposite sign are juxtaposed and are everywhere close together in what can roughly be described as a dipole layer which is spread throughout the 4-dimensional Euclidean space (with various folds and wrinkles). The vacuum is thus permeated with what is locally a laminar structure consisting of alternating sign sheets or membranes of topological charge. The thickness of these membranes is typically a few lattice spacings, independent of the physical mass scale, and thus the membranes apparently become infinitely thin in the continuum limit. This kind of "subdimensional" ordering, where coherence takes place on manifolds of lower dimensionality than spacetime itself, is closely related to the appearance of a contact term in the two-point topological charge correlator. The Euclidean correlator $G(x) = \langle q(x)q(0)\rangle$ *is required by spectral considerations to be negative for any nonzero separation* $|x| > 0$. In practice (i.e. in Monte Carlo calculations), this requirement places severe restrictions on what type of topological charge fluctuations can be dominant. For example, the negativity of the correlator rules out the dominance of bulk-coherent lumps of topological charge (e.g. finite size instantons), as this would lead to a positive correlator over distances smaller than the instanton size. Since the topological susceptibility can be obtained by integrating the 2-point correlator over all x, a positive susceptibility can only arise from a delta-function contact term at the origin. The observed arrangement of thin, nearby layers of $q(x)$ with opposite sign builds up a positive contact term at $x = 0$ while maintaining the required negativity of the correlator for finite separation. The range of this contact term on the lattice is associated with the thickness of the sheets, both of these being a few lattice

spacings and approaching zero in physical units. This thickness is consistent with the range of non-ultralocality of the overlap Dirac operator.

Since the initial discovery of coherent topological charge sheets in QCD, similar methods have been applied to the study of 2-dimensional CP^{N-1} sigma models [3]. For $N > 3$, the topological charge distribution was found to be dominated by thin 1-dimensionally coherent membrane-like structures with interleaved membranes of opposite sign. This interleaved arrangement is exactly what one would expect as the analog of the observed 3-dimensional structures in 4D gauge theory. In both cases, the coherent structure has codimension 1, i.e. the dimensionality of a domain wall.

To interpret these Monte Carlo results, we recall the arguments of Luscher [4] that nonzero topological susceptibility implies the presence of a massless pole in the two-point correlator of the Chern-Simons current. Let us define the *abelian* 3-index Chern-Simons tensor

$$A_{\mu\nu\rho} = -Tr\left(B_\mu B_\nu B_\rho + \frac{3}{2}B_{[\mu}\partial_\nu B_{\rho]}\right) \tag{1}$$

where B_μ is the Yang-Mills gauge potential. We consider the Chern-Simons current that is dual to this tensor,

$$j_\mu^{CS} = \varepsilon_{\mu\nu\rho\sigma}A_{\nu\rho\sigma} \; . \tag{2}$$

Although j_μ^{CS} is not gauge invariant, its divergence is the gauge invariant topological charge density

$$\partial_\mu j_\mu^{CS} = TrF\tilde{F} = 32\pi^2 q(x) \; . \tag{3}$$

Choosing a covariant gauge, $\partial_\mu A_{\mu\nu\rho} = 0$, the correlator of two Chern-Simons currents has the form

$$\langle j_\mu^{CS}(x) j_\nu^{CS}(0)\rangle = \int \frac{d^4p}{(2\pi)^4}\, e^{-ip\cdot x}\, \frac{p_\mu p_\nu}{p^2} G(p^2) \; . \tag{4}$$

From (3) we see that $G(p^2)$ must have a $p^2 = 0$ pole whose residue is the topological susceptibility,

$$G(p^2) \sim \frac{\chi_t}{p^2} \; . \tag{5}$$

This long-range correlation constitutes a "secret long-range order" of gauge fields associated with their topological charge fluctuations. Since the CS current is not gauge invariant, the presence of a $q^2 = 0$ pole does not imply the existence of a massless particle. On the other hand, the pole has a gauge invariant residue ($\propto \chi_t$) and cannot be transformed away. So it characterizes a physically significant long range coherence in the gauge field associated with topological charge fluctuations.

Luscher's analysis of QCD topological structure in terms of Wilson bags can be understood as a generalization of the analysis of similar properties in the 2-dimensional CP^{N-1} sigma models. These models provide a quite detailed 2D analog of the coherent structure observed in 4D QCD. The CP^{N-1} models have a $U(1)$ gauge invariance and have classical instanton solutions which come in all sizes. (Just like pure-glue QCD these models are classically scale invariant and acquire a mass scale via a conformal anomaly.) In 2D U(1) gauge theories like CP^{N-1}, the topological charge density in the

continuum is just $(1/2\pi)\varepsilon_{\mu\nu}\partial_\mu A_\nu$ and the Chern-Simons current is just the dual of the gauge potential,

$$j_\mu^{CS} = \frac{1}{2\pi}\varepsilon_{\mu\nu}A_\nu \tag{6}$$

Just as in 4D QCD, nonzero topological susceptibility implies the presence of a $q^2 = 0$ pole in the CS current correlator. But in the two-dimensional case, this same pole appears in the A_μ correlator and is responsible for confinement of $U(1)$ charge via a linear coulomb potential. Thus, in 2-dimensional $U(1)$ theories, topological susceptibility and confinement of $U(1)$ charge are equivalent phenomena. An instructive way to illustrate this is to introduce a nonzero θ term in the action over a two-volume V enclosed by a boundary $C = \partial V$ with $\theta = 0$ outside the boundary. After integration by parts, the theta term is equivalent to a Wilson loop around the boundary carrying a charge $\theta/2\pi$:

$$\exp\left[\frac{i}{2\pi}\int d^2x\theta(x)\varepsilon_{\mu\nu}F^{\mu\nu}\right] = \exp\left[\frac{i\theta}{2\pi}\oint_C A\cdot dx\right] \tag{7}$$

If the topological susceptibility is nonzero

$$\chi_t = \frac{\partial^2 E(\theta)}{\partial\theta^2}|_{\theta=0} > 0 \tag{8}$$

then for small nonzero θ, the vacuum energy density $E(\theta)$ inside the Wilson loop will be greater than that outside the loop, so it will obey an area law,

$$\langle W(C)\rangle \propto \exp\left[-\left(E(\theta) - E(0)\right)V\right] \tag{9}$$

In a Hamiltonian framework, this corresponds to applying a background electric field θ by putting opposite charges at either end of the 1-dimensional spatial box. The topological susceptibility is just the vacuum polarizability with respect to this field. Periodicity in $\theta \to \theta + 2\pi$ arises via a discontinuous process of charged pair production at $\theta = \pi$, which screens a unit of electric flux. At $\theta = 2\pi$ the confining force is completely screened. This is essentially Coleman's original interpretation of θ-dependence in the massive Schwinger model [5].

In specifying the analogy between 2D U(1) theories and 4D SU(N) gauge theories, we take the Chern-Simons currents (2) and (6) to be directly analogous. This means that the gauge field A_μ in the 2D theory should be identified *not* with the 4-dimensional gauge field, but with the abelian 3-index Chern-Simons tensor (1). Like the gauge field A_μ in 2 dimensions, this is dual to the Chern-Simons current (2). Similarly, the Wilson loop or line excitations in the 2-dimensional $U(1)$ models correspond not to Wilson loops in 4D, but to "Wilson bags," i.e. integrals of the Chern-Simons tensor over a 3-surface Σ.

$$B(\Sigma) = \exp\left[i(\theta/2\pi)\int_\Sigma A_{\mu\nu\lambda}dx_\mu dx_\nu dx_\lambda\right] \tag{10}$$

This is the analog of a Wilson loop in 2D $U(1)$ in the sense that, if Σ is a closed 3-surface that forms the boundary of a 4-dimensional volume V, inserting the Wilson bag factor (10) in the gauge field path integral is equivalent to including a θ-term in the gauge

action over the 4-volume V. The discussion of what happens as we vary θ from 0 to 2π is completely analogous to the screening of the 2D Wilson loop. For a fractional bag charge $\theta/2\pi$, with $0 < \theta < 2\pi$, the vacuum inside the bag will have a higher energy than the $\theta = 0$ vacuum outside. The expectation of the Wilson bag integral thus satisfies a volume law analogous to the area law for the Wilson loop in 2D, and there will be "bag confinement," a confining force between the walls of a fractionally charged bag. At $\theta = 2\pi$, the confining force between bag walls disappears. Integer charged bag surfaces are thus free to percolate throughout the vacuum. The topological charge is the curl of the Chern-Simons tensor, so for a uniform $A_{\mu\nu\lambda}$ which is nonzero on a flat bag surface, the topological charge distribution is a dipole layer consisting of thin, coherent positive and negative layers on either side of the bag surface. Like the Wilson line excitations in the CP^{N-1} models [3], a vacuum full of unit-charged Wilson bags provides a reasonable model for interpreting the topological charge structure observed in lattice Monte Carlo simulations.

The role of Wilson bag excitations in QCD is greatly illuminated by string/gauge duality, exploiting the dual relationship between topological charge in gauge theory and Ramond-Ramond charge in IIA string theory [7]. As Witten showed [6], the string/gauge correspondence nicely confirms the k-vacuum/Wilson bag/domain wall scenario originally suggested by large-N_c chiral Lagrangian arguments [8]. It also points to the correct candidate for the string theory analog of a Wilson bag in gauge theory. In IIA string theory, the Wilson bag can be interpreted as the holographic image of a D6-brane (which is wrapped around a compact S_4 and therefore appears as a 2-brane or membrane in 3+1 dimensions). In particular, the defining property of the Wilson bag, namely that the value of θ jumps by $\pm 2\pi$ when crossing the surface, is in fact nothing but the statement of quantization of Ramond-Ramond charge on a D6-brane.

This work was supported in part by the Department of Energy under grant DE-FG02-97ER41027.

REFERENCES

1. I. Horvath et al, Phys. Rev. D68, 114505 (2003).
2. E.-M. Ilgenfritz et al., Nucl. Phys. Proc. Suppl. 153: 328 (2006); and hep-lat/0610087.
3. S. Ahmad, J. T. Lenaghan, H. B. Thacker, Phys. Rev. D72: 114511 (2005); Y. Lian and H. B. Thacker, hep-lat/0607026.
4. M. Luscher, Phys. Lett. B78, 465 (1978).
5. S. Coleman, Ann. Phys. 101: 239 (1976).
6. E. Witten, Phys. Rev. Lett. 81: 2862 (1998).
7. For a discussion, see H. Thacker, hep-lat/0610049.
8. E. Witten, Ann. Phys. 128: 363 (1980).