arXiv:hep-th/9502159v1 28 Feb 1995

DPNU-95-06
hep-th/9502159

Topology and Inequivalent Quantizations of Abelian Sigma Model[1]

Shogo Tanimura[2]

Department of Physics, Nagoya University, Nagoya 464-01, Japan[3]

The abelian sigma model in (1+1) dimensions is a field theoretical model which has a field $\phi : S^1 \to S^1$. An algebra of the quantum field is defined respecting the topological aspect of the model. It is shown that the zero-mode has an infinite number of inequivalent quantizations. It is also shown that when a central extension is introduced into the algebra, the winding operator and the momenta operators satisfy anomalous commutators.

1. Introduction

In both field theory and string theory there are several models which have manifold-valued variables, for instance, the nonlinear sigma model and the toroidal compactification model of closed bosonic string. To study global aspects of these models in quantum theory, we should have a quantization scheme respecting topological nature. However in the usual scheme of canonical quantization and perturbation method, the global aspects are obscure. On the other hand it is known [1], [2] that there exist an infinite number of inequivalent quantizations on a topologically nontrivial manifold even when it is a finite-dimensional manifold. In this note we consider the abelian sigma model in (1+1) dimensions to explore a system having infinite degrees of freedom. The model has a manifold-valued field $\phi : S^1 \to S^1$. For detail see the reference [3].

2. Ohnuki-Kitakado representation

We briefly review quantum mechanics of a particle on S^1 [1]. Let us assume that a unitary operator $\hat{U}$ and a self-adjoint operator $\hat{P}$ satisfy the commutation relation

$$[\,\hat{P}\,,\,\hat{U}\,] = \hat{U}. \tag{1}$$

An irreducible representation of the above algebra is called quantum mechanics on S^1. The operators $\hat{U}$ and $\hat{P}$ are called a position operator and a momentum operator, respectively [4].

A representation space is constructed as follows. Let $|\alpha\rangle$ be an eigenvector of $\hat{P}$ with a real eigenvalue α; $\hat{P}\,|\alpha\rangle = \alpha\,|\alpha\rangle$. Assume that $\langle\alpha|\alpha\rangle = 1$. The commutator (1) implies that the operator $\hat{U}$ increases an eigenvalue of $\hat{P}$ by a unit. If we put

$$|n+\alpha\rangle := \hat{U}^n\,|\alpha\rangle \quad (n = 0, \pm1, \pm2, \cdots), \tag{2}$$

[1]Presented on 19 January 1995 at Second Pasific Winter School for Theoretical Physics held at Sorak Mountain in Korea. At that time the title of seminar was "Winding number, zero-mode and commutators of abelian sigma model in (1+1) dimensions", but it is changed here.

[2]e-mail address : tanimura@eken.phys.nagoya-u.ac.jp

[3]Leaving for Department of Physics, University of Tokyo, Bunkyo-ku,Tokyo 113, Japan

it is easily seen that

$$\hat{P}\,|n+\alpha\rangle = (n+\alpha)\,|n+\alpha\rangle, \quad \hat{U}\,|n+\alpha\rangle = |n+1+\alpha\rangle. \tag{3}$$

Unitarity of $\hat{U}$ and self-adjointness of $\hat{P}$ imply that

$$\langle m+\alpha|n+\alpha\rangle = \delta_{m\,n}. \tag{4}$$

Let H_α denote the Hilbert space defined by completing the space of finite linear combinations of $|n+\alpha\rangle\,(n = 0, \pm1, \pm2, \cdots)$. By (3), H_α becomes an irreducible representation space of the algebra (1). H_α and H_β are unitary equivalent if and only if the difference $(\alpha-\beta)$ is an integer. Consequently there exists an inequivalent representation for each value of the parameter α ranging over $0 \le \alpha < 1$. Physical significance of α is further discussed in the reference [4].

3. Algebra

Next we shall propose an algebra of the sigma model. In the context of classical theory, the model has a field variable $\phi \in Q = \mathrm{Map}(S^1; S^1)$. Let $\Gamma = \mathrm{Map}(S^1; U(1))$ a group by pointwise composition. The group Γ acts on the configuration space Q by pointwise action of $U(1)$ on S^1. To quantize this system let us assume that $\hat{\phi}(\theta)$ is a unitary operator for each point $\theta \in S^1$ and $\hat{V}(\gamma)$ is a unitary operator for each element $\gamma \in \Gamma$. Moreover we assume the following algebra

$$\hat{\phi}(\theta)\,\hat{\phi}(\theta') = \hat{\phi}(\theta')\,\hat{\phi}(\theta), \tag{5}$$

$$\hat{V}(\gamma)^\dagger\,\hat{\phi}(\theta)\,\hat{V}(\gamma) = \gamma(\theta)\,\hat{\phi}(\theta), \tag{6}$$

$$\hat{V}(\gamma_1)\,\hat{V}(\gamma_2) = e^{-i\,c(\gamma_1,\gamma_2)}\,\hat{V}(\gamma_1\cdot\gamma_2) \quad (\gamma_1,\,\gamma_2 \in \Gamma). \tag{7}$$

In the last line a function $c : \Gamma\times\Gamma \to \boldsymbol{R}$ is called a central extension, which satisfies the cocycle condition; $c(\gamma_1,\gamma_2)+c(\gamma_1\gamma_2,\gamma_3) = c(\gamma_1,\gamma_2\gamma_3)+c(\gamma_2,\gamma_3)\ (\mathrm{mod}\,2\pi)$. We call the algebra (5)-(7) the fundamental algebra of the abelian sigma model. To consider a nontrivial central extension we decompose γ as

$$\gamma(\theta) = e^{i\,(\mu+f(\theta)+m\,\theta)}, \tag{8}$$

where $\mu \in \boldsymbol{R}$, $f \in \mathrm{Map}_0(S^1;\boldsymbol{R}) := \{g : S^1 \to \boldsymbol{R}\,|\,C^\infty,\ \int_0^{2\pi} g(\theta)\,d\theta = 0\}$, and $m \in \boldsymbol{Z}$. For γ_1 shown at (8) and $\gamma_2(\theta) = \exp[i\,(\mu_2 + f_2(\theta) + m_2\,\theta)]$, we define

$$c(\gamma_1,\gamma_2) := k\left\{m_1\,\mu_2 - \mu_1\,m_2 + \frac{1}{4\pi}\int_0^{2\pi}\left(f_1'(\theta)f_2(\theta) - f_1(\theta)f_2'(\theta)\right)d\theta\right\}, \tag{9}$$

where k is a non-zero integer. This central extension is the simplest but nontrivial one which is invariant under the action of $\mathrm{Diff}^+(S^1)$; $c(\gamma_1\circ\omega, \gamma_2\circ\omega) = c(\gamma_1,\gamma_2)$ for any $\omega \in \mathrm{Diff}^+(S^1)$. The group Γ associated with such an invariant central extension is called a Kac-Moody group [5].

a. Algebra without central extension : To clarify geometrical implication of the algebra we shall decompose the degrees of freedom of $\hat{\phi}$ and $\hat{V}$. At first we put $c \equiv 0$. We introduce a unitary operator $\hat{U}$, self-adjoint operators[4] $\hat{\varphi}(\theta)$ for each

[4]Expressing rigorously $\hat{\varphi}(\theta)$ is an operator-valued distribution. $\exp(i\hat{\varphi}(\theta))$ must be regularized by the normal product procedure[3].

$\theta \in S^1$ constrained by $\int_0^{2\pi} \hat{\varphi}(\theta)\, d\theta = 0$, and a self-adjoint operator $\hat{N}$ satisfying $\exp(2\pi i \hat{N}) = 1$, which is called the integer condition for $\hat{N}$. We demand that the quantum field $\hat{\phi}(\theta)$ is expressed by them as

$$\hat{\phi}(\theta) = \hat{U}\, e^{i\,(\hat{\varphi}(\theta) + \hat{N}\theta)}. \tag{10}$$

Geometrical meaning of this decomposition is apparent; $\hat{U}$ describes the zero-mode or collective motion of the field $\hat{\phi}$; $\hat{\varphi}$ describes fluctuation or local degrees of freedom; $\hat{N}$ is nothing but the winding number. Topologically nontrivial parts are $\hat{U}$ and $\hat{N}$. Next, corresponding to (8) we introduce a self-adjoint operator $\hat{P}$, self-adjoint operators $\hat{\pi}(\theta)$ for each $\theta \in S^1$ constrained by $\int_0^{2\pi} \hat{\pi}(\theta)\, d\theta = 0$, and a unitary operator $\hat{W}$. When γ is given by (8), we define $\hat{V}(\gamma)$ by

$$\hat{V}(\gamma) = e^{-i\,\mu\,\hat{P}}\, \exp\left[-i \int_0^{2\pi} f(\theta)\, \hat{\pi}(\theta)\, d\theta\right] \hat{W}^m. \tag{11}$$

Using these operators the fundamental algebra is now rewritten as

$$[\,\hat{P}, \hat{U}\,] = \hat{U}, \tag{12}$$

$$[\,\hat{\varphi}(\theta), \hat{\pi}(\theta')\,] = i\Big(\delta(\theta - \theta') - \frac{1}{2\pi}\Big), \tag{13}$$

$$[\,\hat{N}, \hat{W}\,] = \hat{W}, \tag{14}$$

and all other commutators vanish. In (13) the δ-function is periodic with the periodicity 2π. Observing the relation (14) we call $\hat{W}$ the winding operator.

b. Algebra with central extension : If the central extension (9) is included, the decomposition (11) of $\hat{V}$ should be modified a little. It is replaced by

$$\hat{V}(\gamma) = e^{-ikm\mu}\, e^{-i\mu\hat{P}}\, \exp\left[-i \int_0^{2\pi} f(\theta)\, \hat{\pi}(\theta)\, d\theta\right] \hat{W}^m. \tag{15}$$

Furthermore the following commutation relations should be added to (12)-(14);

$$[\,\hat{P}, \hat{W}\,] = -2k\, \hat{W}, \tag{16}$$

$$[\,\hat{\pi}(\theta), \hat{\pi}(\theta')\,] = -\,\frac{ik}{\pi}\, \delta'(\theta - \theta'). \tag{17}$$

(16) says that the zero-mode momentum $\hat{P}$ is decreased by $2k$ units when the winding number $\hat{N}$ is increased by one unit under the operation of $\hat{W}$. This is an inevitable consequence of the central extension. We call this phenomenon "twist".

4. Representations

a. Without the central extension : Representations of the algebra defined by (12)-(14) and other vanishing commutators are constructed below. $\hat{P}$ and $\hat{N}$ are self-adjoint and that $\hat{U}$ and $\hat{W}$ are unitary. Both of the relations (12) and (14) are isomorphic to (1). Hence the Ohnuki-Kitakado representation can be used for them. $\hat{P}$ and $\hat{U}$ act on the Hilbert space H_α via (3). $\hat{N}$ and $\hat{W}$ act on another Hilbert space H_β via $\hat{N}\,|n+\beta\rangle = (n+\beta)\,|n+\beta\rangle$ and $\hat{W}\,|n+\beta\rangle = |n+1+\beta\rangle$. The value of α is

arbitrary. However β is restricted to be an integer if we impose the integer condition $\exp(2\pi i\,\hat{N}) = 1$. For $\hat{\varphi}$ and $\hat{\pi}$ we define operators $\hat{a}_n$ and $\hat{a}_n^\dagger$ by

$$\hat{\varphi}(\theta) = \frac{1}{2\pi}\sum_{n\neq 0}\sqrt{\frac{\pi}{|n|}}\,(\hat{a}_n\,e^{i\,n\,\theta} + \hat{a}_n^\dagger\,e^{-i\,n\,\theta}),\quad \hat{\pi}(\theta) = \frac{i}{2\pi}\sum_{n\neq 0}\sqrt{\pi|n|}\,(-\hat{a}_n\,e^{i\,n\,\theta} + \hat{a}_n^\dagger\,e^{-i\,n\,\theta}). \tag{18}$$

It is easily verified that the commutator (13) is equivalent to

$$[\,\hat{a}_m, \hat{a}_n^\dagger] = \delta_{m\,n}\quad (m, n = \pm 1, \pm 2, \cdots) \tag{19}$$

with the other vanishing commutators. Hence the ordinary Fock space F gives a representation of $\hat{a}$'s and $\hat{a}^\dagger$'s. Consequently the tensor product space $H_\alpha \otimes H_0 \otimes F$ $(0 \leq \alpha < 1)$ gives an irreducible representation of the algebra (12)-(14).

b. With the central extension : Representations of the algebra defined by (12), (13), (14), (16) and (17) and other vanishing commutators are also constructed in a similar way. The twist relation (16) is represented by

$$\begin{cases} \hat{P}\,|\,p+\alpha;\,n\rangle = (p+\alpha)\,|\,p+\alpha;\,n\rangle, & \hat{U}\,|\,p+\alpha;\,n\rangle = |\,p+1+\alpha;\,n\rangle, \\ \hat{N}\,|\,p+\alpha;\,n\rangle = n|\,p+\alpha;\,n\rangle, & \hat{W}\,|\,p+\alpha;\,n\rangle = |\,p-2k+\alpha;\,n+1\rangle. \end{cases} \tag{20}$$

The Hilbert space formed by completing the space of linear combinations of $|\,p+\alpha;\,n\rangle$ is denoted by $T_{k\alpha}$. (T indicates "twist".) Taking the anomalous commutator (17) into account the Fourier expansions (18) of $\hat{\varphi}$ and $\hat{\pi}$ are changed to

$$\begin{aligned} \hat{\varphi}(\theta) &= \sum_{n\neq 0}\frac{1}{\sqrt{2|kn|}}\,(\hat{a}_n\,e^{i\,n\,\theta} + \hat{a}_n^\dagger\,e^{-i\,n\,\theta}), \\ \hat{\pi}(\theta) &= \frac{i}{2\pi}\sum_{n\neq 0}\theta(kn)\,\sqrt{2|kn|}\,(-\hat{a}_n\,e^{i\,n\,\theta} + \hat{a}_n^\dagger\,e^{-i\,n\,\theta}). \end{aligned} \tag{21}$$

It should be noticed that only positive n's appear in the expansion of $\hat{\pi}$ even though both of positive and negative n's appear in $\hat{\varphi}$. Thus the tensor product space $T_{k\alpha} \otimes F$ $(0 \leq \alpha < 1)$ gives an irreducible representation of the fundamental algebra.

Acknowledgment

I thank deeply Yoonbai Kim, who encouraged me to attend the winter school.

References

[1] Y. Ohnuki and S. Kitakado, *J. Math. Phys.* **34** (1993) 2827.

[2] G. W. Mackey, *Induced Representations of Groups and Quantum Mechanics*, (W. A. Benjamin, INC., New York, 1968); N. P. Landsman and N. Linden, *Nucl. Phys.* **B365** (1991) 121; S. Tanimura, Nagoya University preprint DPNU-93-21 (1993), hep-th/9306144.

[3] S. Tanimura, *Phys. Lett.* **340B** (1994) 57; Nagoya University preprint DPNU-94-60 (1994), hep-th/9412174 (submitted to *Nucl. Phys.*).

[4] S. Tanimura, *Prog. Theor. Phys.* **90** (1993) 271.

[5] G. Segal, *Commun. Math. Phys.* **80** (1981) 301.