

PARTIAL CONNECTION FOR p -TORSION LINE BUNDLES IN CHARACTERISTIC $p > 0$

HÉLÈNE ESNAULT

To S. S. Chern, in memoriam

ABSTRACT. The aim of this brief note is to give a construction for p -torsion line bundles in characteristic $p > 0$ which plays a similar rôle as the standard connection on an n -torsion line bundle in characteristic 0.

1. INTRODUCTION

In [3] (see also [4]) we gave an algebraic construction of characteristic classes of vector bundles with a flat connection (E, ∇) on a smooth algebraic variety X defined over a field k of characteristic 0. Their value at the generic point $\text{Spec}(k(X))$ was studied and redefined in [1], and then applied in [2] to establish a Riemann-Roch formula. One way to understand Chern classes of vector bundles (without connection) is via the Grothendieck splitting principle: if the receiving groups $\oplus_n H^{2n}(X, n)$ of the classes form a cohomology theory which is a ring and is functorial in X , then via the Whitney product formula it is enough to define the first Chern class. Indeed, on the flag bundle $\pi : \text{Flag}(E) \rightarrow X$, $\pi^*(E)$ acquires a complete flag $E_i \subset E_{i+1} \subset \pi^*(E)$ with E_{i+1}/E_i a line bundle, and $\pi^* : H^{2n}(X, n) \rightarrow H^{2n}(\text{Flag}(E), n)$ is injective, so it is enough to construct the classes on $\text{Flag}(E)$. However, if ∇ is a connection on E , $\pi^*(\nabla)$ does not stabilize the flag E_i . So the point of [3] is to show that there is a differential graded algebra $A^\bullet$ on $\text{Flag}(E)$, together with a morphism of differential graded algebras $\Omega_{\text{Flag}(E)}^\bullet \xrightarrow{\tau} A^\bullet$, so that $R\pi_* A^\bullet \cong \Omega_X^\bullet$ and so that the operator defined by the composition $\pi^*(E) \xrightarrow{\pi^*(\nabla)} \Omega_{\text{Flag}(E)}^1 \otimes_{\mathcal{O}_{\text{Flag}(E)}} \pi^*(E) \xrightarrow{\tau \otimes 1} A^1 \otimes_{\mathcal{O}_{\text{Flag}(E)}} \pi^*(E)$ stabilizes E_i . We call the induced operator $\nabla_i : E_i \rightarrow A^1 \otimes_{\mathcal{O}_{\text{Flag}(E)}} E_i$ a (flat) τ -connection. So it is a k -linear map which fulfills the τ -Leibniz

Date: August 7, 2006.

Partially supported by the DFG Leibniz Preis.

formula

$$(1.1) \quad \nabla_i(\lambda \otimes e) = \tau d(\lambda) \otimes e + \lambda \nabla_i(e)$$

for λ a local section of $\mathcal{O}_{\text{Flag}(E)}$ and e a local section of E_i . It is flat when $0 = \nabla_i \circ \nabla_i \in H^0(X, A^2 \otimes_{\mathcal{O}_X} \mathcal{E}nd(E))$, with the appropriate standard sign for the derivation of forms with values in E_i . The last point is then to find the correct cohomology which does not get lost under π^* . It is a generalization of the classically defined group

$$(1.2) \quad \mathbb{H}^1(X, \mathcal{O}_X^\times \xrightarrow{d\log} \Omega_X^1 \xrightarrow{d} \Omega_X^2 \xrightarrow{d} \dots)$$

of isomorphism classes of rank one line bundles on X with a flat connection.

A typical example of such a connection is provided by a torsion line bundle: if L is a line bundle on X which is n -torsion, that is which is endowed with an isomorphism $L^n \cong \mathcal{O}_X$, then the isomorphism yields an $\mathcal{O}_X$ -étale algebra structure on $\mathcal{A} = \bigoplus_0^{n-1} L^i$, hence a finite étale covering $\sigma : Y = \text{Spec}_{\mathcal{O}_X} \mathcal{A} \rightarrow X$, which is a principal bundle under the group scheme μ_n of n -th roots of unity, thus is Galois cyclic as soon as $\mu_n \subset k^\times$. Since the μ_n -action commutes with the differential $d_Y : \mathcal{O}_Y \rightarrow \Omega_Y^1 = \sigma^* \Omega_X^1$, it defines a flat connection $\nabla_L : L \rightarrow \Omega_X^1 \otimes_{\mathcal{O}_X} L$. Concretely, if $g_{\alpha,\beta} \in \mathcal{O}_X^\times$ are local algebraic transition functions for L , with trivialization

$$(1.3) \quad g_{\alpha,\beta}^n = u_\beta u_\alpha^{-1}, u_\alpha \in \mathcal{O}_X^\times,$$

then

$$(1.4) \quad (g_{\alpha,\beta}, \frac{1}{n} \frac{du_\alpha}{u_\alpha}) \in (\mathcal{C}^1(\mathcal{O}_X^\times) \times \mathcal{C}^0((\Omega_X^1)_{\text{clsd}}))_{d\log-\delta}$$

$$\frac{dg}{g} = \delta\left(\frac{du}{u}\right)$$

is a Čech cocycle for the class

$$(1.5) \quad (L, \nabla_L) \in \mathbb{H}^1(X, \mathcal{O}_X^\times \xrightarrow{d\log} \Omega_X^1 \xrightarrow{d} \Omega_X^2 \xrightarrow{d} \dots).$$

Clearly (1.4) is meaningless if the characteristic p of k is positive and divides n . The purpose of this short note is to give an Ersatz of this canonical construction in the spirit of the τ -connections explained above when p divides n .

2. A PARTIAL CONNECTION FOR p -TORSION LINE BUNDLES

Let X be a scheme of finite type over a perfect field k of characteristic $p > 0$. Let L be a n -torsion line bundle on X , thus endowed with an

isomorphism

$$(2.1) \quad \theta : L^n \cong \mathcal{O}_X.$$

Then θ defines an $\mathcal{O}_X$ -algebra structure on $\mathcal{A} = \bigoplus_0^{n-1} L^i$ which is étale if and only if $(p, n) = 1$. It defines the principal μ_n -covering

$$(2.2) \quad \sigma : Y = \operatorname{Spec}_{\mathcal{O}_X} \mathcal{A} \rightarrow X$$

which is étale if and only if $(p, n) = 1$, else decomposes into

$$(2.3) \quad \sigma : Y \xrightarrow{\iota} Z \xrightarrow{\sigma'} X$$

with σ' étale and ι purely inseparable. More precisely, if $n = m \cdot p^r$, $(m, p) = 1$, and $M = L^{p^r}$, θ defines an $\mathcal{O}_X$ -étale algebra structure on $\mathcal{B} = \bigoplus_0^{m-1} M^i$, which defines $\sigma' : Z = \operatorname{Spec}_{\mathcal{O}_X} \mathcal{B} \rightarrow X$ as an (étale) μ_m -principal bundle. The isomorphism θ also defines an isomorphism $(L')^{p^r} \cong \mathcal{O}_Z$ as it defines the isomorphism $(\sigma')^*(M) \cong \mathcal{O}_Z$, where $L' = (\sigma')^*(L)$. So $\mathcal{C} = \bigoplus_0^{p^r-1} (L')^i$ becomes a finite purely inseparable $\mathcal{O}_Z$ -algebra defining the principal μ_{p^r} -bundle $\iota : Y = \operatorname{Spec}_{\mathcal{O}_Z} \mathcal{C} \rightarrow Z$.

If $(n, p) = 1$, that is if $r = 0$, the formulae (1.3), (1.4) define (L, ∇) as in (1.5). We assume from now on that $(n, p) = p$. Then, as is well known, as a consequence of (1.3) one sees that the form

$$(2.4) \quad \omega_L := \frac{du_\alpha}{u_\alpha} \in \Gamma(X, \Omega_X^1)_{\text{clsd}}^{\text{Cartier}=1}$$

is globally defined and Cartier invariant. Let e_α be local generators of L , with transition functions $g_{\alpha, \beta}$ with $e_\alpha = g_{\alpha, \beta} e_\beta$. The isomorphism θ yields a trivialization

$$(2.5) \quad \sigma^* L \cong \mathcal{O}_Y$$

thus local units v_α on Y with

$$(2.6) \quad v_\alpha \in \mathcal{O}_Y^\times, \quad g_{\alpha, \beta} = v_\beta v_\alpha^{-1}$$

so that $1 = v_\alpha \sigma^*(e_\alpha) = v_\beta \sigma^*(e_\beta)$.

Definition 2.1. One defines the $\mathcal{O}_X$ -coherent sheaf Ω_L^1 as the subsheaf of $\sigma_* \Omega_Y^1$ spanned by $\operatorname{Im}(\Omega_X^1)$ and $\frac{dv_\alpha}{v_\alpha}$.

Lemma 2.2. Ω_L^1 is well defined and one has the exact sequence

$$(2.7) \quad 0 \rightarrow \mathcal{O}_X \xrightarrow{\omega_L} \Omega_X^1 \xrightarrow{\sigma^*} \Omega_L^1 \xrightarrow{s} \mathcal{O}_X \rightarrow 0$$

$$s\left(\frac{dv_\alpha}{v_\alpha}\right) = 1.$$

Proof. The relation (2.6) implies

$$(2.8) \quad \frac{dg_{\alpha,\beta}}{g_{\alpha,\beta}} = \frac{dv_\beta}{v_\beta} - \frac{dv_\alpha}{v_\alpha}$$

so $\frac{dv_\beta}{v_\beta} \equiv \frac{dv_\alpha}{v_\alpha} \in \sigma_*\Omega_Y^1/\text{Im}(\Omega_X^1).$

Hence the sheaf Ω_L^1 is well defined. If e'_α is another basis, then one has $e_\alpha = w_\alpha e'_\alpha$ for local units $w_\alpha \in \mathcal{O}_X^\times$. The new v_α are then multiplied by local units in $\mathcal{O}_X^\times$, so the surjection s is well defined. It remains to see that $\text{Ker}(\sigma^*) = \text{Im}(\cdot\omega_L)$. By definition, on the open of X on which L has basis e_α , one has

$$(2.9) \quad Y = \text{Spec } \mathcal{O}_X[v_\alpha]/(v_\alpha^n - u_\alpha).$$

This implies $\Omega_Y^1 = \langle \text{Im}(\Omega_X^1), dv_\alpha \rangle_{\mathcal{O}_Y} / \langle du_\alpha \rangle_{\mathcal{O}_Y}$ on this open and finishes the proof. $\square$

- Remarks 2.3.**
- 1) Assume for example that X is a smooth projective curve of genus g , and $n = p$. Recall that $0 \neq \omega_L \in \Gamma(X, \Omega_X^1)$. In particular, if $g \geq 2$, necessarily $0 \neq \Omega_X^1/\mathcal{O}_X \cdot \omega_L$ is supported in codimension 1. So Ω_L^1 contains a non-trivial torsion subsheaf.
 - 2) The sheaf Ω_L^1 lies in $\sigma_*\Omega_Y^1$ but is not equal to it. Indeed, on the smooth locus of X (assuming X is reduced) the torsion free quotient of Ω_L^1 has rank equal to the dimension of X , while $\sigma_*\Omega_Y^1$ has rank $n \cdot \dim(X)$ on the étale locus of σ (which is non-empty if L itself is not a p -power line bundle).
 - 3) The class in $\text{Ext}_{\mathcal{O}_X}^2(\mathcal{O}_X, \mathcal{O}_X) = H^2(X, \mathcal{O}_X)$ defined by (2.7) vanishes. Indeed, let us decompose (2.7) as an extension of $\mathcal{O}_X$ by $\Omega_X^1/\mathcal{O}_X \cdot \omega_L$, followed by an extension of $\Omega_X^1/\mathcal{O}_X \cdot \omega_L$ by $\mathcal{O}_X \cdot \omega_L$. The first extension class in $H^1(X, \Omega_X^1/\mathcal{O}_X \cdot \omega_L)$ has cocycle $\frac{dv_\beta}{v_\beta} - \frac{dv_\alpha}{v_\alpha} = \frac{dg_{\alpha,\beta}}{g_{\alpha,\beta}}$ (see (2.8)), thus is the image of the Atiyah class of L in $H^1(X, \Omega_X^1)$. Thus the second boundary to $H^2(X, \mathcal{O}_X)$ dies.

Definition 2.4. We set $\Omega_L^0 := \mathcal{O}_X$ and for $i \geq 1$ we define the $\mathcal{O}_X$ -coherent sheaf Ω_L^i as the subsheaf of $\sigma_*\Omega_Y^i$ spanned by $\text{Im}(\Omega_X^i)$ and $\frac{dv_\alpha}{v_\alpha} \wedge \text{Im}(\Omega_X^{i-1})$.

Proposition 2.5. *The sheaf Ω_L^i is well defined. One has an exact sequence*

$$(2.10) \quad 0 \rightarrow \omega_L \wedge \Omega_X^{i-1} \rightarrow \Omega_X^i \xrightarrow{\sigma^*} \Omega_L^i \xrightarrow{s} \Omega_X^{i-1} \rightarrow 0$$

$$s\left(\frac{dv_\alpha}{v_\alpha} \wedge \beta\right) = \beta.$$

Furthermore, the differential $\sigma_*(d_Y)$ on $\sigma_*\Omega_Y^\bullet$ induces on $\oplus_{i \geq 0} \Omega_L^i$ the structure of a differential graded algebra $(\Omega_L^\bullet, d_L)$ so that $\sigma^* : (\Omega_X^\bullet, d_X) \rightarrow (\Omega_L^\bullet, d_L)$ is a morphism of differential graded algebras.

Proof. One proves (2.10) as one does (2.7). One has to see that $\sigma_*(d_Y)$ stabilizes $\Omega_L^\bullet$. As $0 = d_X(\omega_L) \in \Omega_X^2$, $0 = d_Y(\frac{dv_\alpha}{v_\alpha}) \in \sigma_*\Omega_Y^2$, (2.10) extends to an exact sequence of complexes

$$(2.11) \quad 0 \rightarrow (\omega_L \wedge \Omega_X^{\bullet-1}, -1 \wedge d_X) \rightarrow (\Omega_X^\bullet, d_X) \xrightarrow{\sigma^*} (\Omega_L^\bullet, d_L) \xrightarrow{s} (\Omega_X^{\bullet-1}, -d_X) \rightarrow 0.$$

This finishes the proof. $\square$

Remark 2.6. As $\frac{dg_{\alpha,\beta}}{g_{\alpha,\beta}} \in (\Omega_X^1)_{\text{clsd}}$ the same proof as in Remark 2.3, 3) shows that the extension class $\text{Ext}^2(\Omega_X^{\bullet-1}, \omega_L \wedge \Omega_X^{\bullet-1})$ defined by (2.11) dies.

In order to tie up with the notations of the Introduction, we set

$$(2.12) \quad \tau = \sigma^* : \Omega_X^\bullet \rightarrow \Omega_L^\bullet.$$

Proposition 2.7. *The formula $\nabla(e_\alpha) = -\frac{dv_\alpha}{v_\alpha} \otimes e_\alpha \in \Omega_L^1 \otimes_{\mathcal{O}_X} L$ defines a flat τ -connection ∇_L on L . So (L, ∇_L) is a class in $\mathbb{H}^1(X, \mathcal{O}_X^\times \xrightarrow{\tau d \log} \Omega_L^1 \xrightarrow{d_L} \Omega_L^2 \xrightarrow{d_L} \cdots)$, the group of isomorphism classes of line bundles with a flat τ -connection.*

Proof. Formula (2.6) implies that this defines a τ -connection. Flatness is obvious. A Čech cocycle for (L, ∇_L) is $(g_{\alpha,\beta}, \frac{dv_\alpha}{v_\alpha})$. $\square$

Remarks 2.8. 1) The same formal definitions 2.1 and 2.4 of $\Omega_L^\bullet$ when $(n, p) = 1$ yield $(\Omega_L^\bullet, d_L) = (\Omega_X^\bullet, d_X)$, and the flat τ -connection becomes the flat connection defined in (1.4) and (1.5). So Proposition 2.7 is a direct generalization of it.
2) Let X be proper reduced over a perfect field k , irreducible in the sense that $H^0(X, \mathcal{O}_X) = k$, and admitting a rational point $x \in X(k)$. A generalization of torsion line bundles to higher rank bundles is the notion of Nori finite bundles, that is bundles E which are trivialized over principal bundle $\sigma : Y \rightarrow X$

under a finite flat group scheme G (see [6] for the original definition and also [5] for a study of those bundles). So for the n -torsion line bundles considered in this section, $G \cong \mu_n$. If the characteristic of k is 0, then again σ is étale, the differential $d_Y : \mathcal{O}_Y \rightarrow \sigma^* \Omega_X^1 = \Omega_Y^1$ commutes with the action of G , inducing a connection $\nabla_E : E \rightarrow \Omega_X^1 \otimes_{\mathcal{O}_X} E$ and characteristic classes in our groups $\mathbb{H}^i(X, \mathcal{K}_i^m \xrightarrow{d\log} \Omega_X^i \xrightarrow{d} \Omega_X^{i+1} \cdots)$ (see [3]). If the characteristic of k is $p > 0$, then σ is étale if and only if G is smooth (which here means étale), in which case one can also construct those classes. If G is not étale, thus contains a non-trivial local subgroup scheme, then one should construct as in Proposition 2.5 a differential graded algebra $(\Omega_E^\bullet, d_E)$ with a map $(\Omega_X^\bullet, d_X) \xrightarrow{\tau} (\Omega_E^\bullet, d_E)$, so that E is endowed naturally with a flat τ -connection $\nabla_E : E \rightarrow \Omega_E^1 \otimes_{\mathcal{O}_X} E$. The techniques developed in [3] should then yield classes in the groups $\mathbb{H}^i(X, \mathcal{K}_i^m \xrightarrow{\tau d\log} \Omega_E^i \xrightarrow{d_E} \Omega_E^{i+1} \cdots)$.

REFERENCES

- [1] Bloch, S., Esnault, H.: Algebraic Chern-Simons theory, Am. J. of Mathematics **119** (1997), 903-952.
- [2] Bloch, S., Esnault, H.: A Riemann-Roch theorem for flat bundles, with values in the algebraic Chern-Simons theory, Annals of Mathematics **151** (2000), 1-46.
- [3] Esnault, H.: Algebraic Differential Characters, in Regulators in Analysis, Geometry and Number Theory, Progress in Mathematics, Birkhäuser Verlag, **171** (2000), 89-117.
- [4] Esnault, H.: Characteristic classes of flat bundles and determinant of the Gauß-Manin connection, Proceedings of the International Congress of Mathematicians, Beijing 2002, Higher Education Press, 471-483.
- [5] Esnault, H., Hai P. H., Sun, X.: On Nori's Fundamental Group Scheme, preprint 2006, 29 pages.
- [6] Nori, M.: The fundamental group scheme, Proc. Indian Acad. Sci. **91** (1982), 73-122.

UNIVERSITÄT DUISBURG-ESSEN, MATHEMATIK, 45117 ESSEN, GERMANY
E-mail address: esnault@uni-due.de