

Heegaard splittings and 1-relator groups

Joseph D. Masters

Abstract

We show that if M is a fibered, orientable 3-manifold, and if $\pi_1 M$ has a 1-relator presentation, then the presentation is induced by a Heegaard splitting of M . A corollary is that, for these manifolds, the rank of $\pi_1 M$ is equal to the (“restricted”) Heegaard genus of M . We also explore the analogy between 1-relator groups and Haken 3-manifolds, showing that every 1-relator group possesses a “1-relator hierarchy”.

1 Introduction

Let F be a closed, orientable surface. A *compression body* W is a cobordism rel ∂ between two surfaces $\partial_+ W$ and $\partial_- W$, such that $W \cong \partial_+ W \times I \cup 2\text{-handles} \cup 3\text{-handles}$, and $\partial_- W$ has no 2-sphere components. A *disk system* $\mathcal{D} = \{D_1, \dots, D_n\}$ for W is a set of compressing disks for $\partial_+ W$, such that W cut open along $\mathcal{D}$ is homeomorphic to $\partial_- \times I$ if $\partial_- \neq \emptyset$, or B^3 if $\partial_- = \emptyset$.

Let M be a compact, orientable 3-manifold. A *Heegaard splitting* of M is a pair (W, W') , where W, W' are compression bodies, such that $W \cup W' = M$, and $W \cap W' = \partial_+ W = \partial_+ W'$. The surface $\partial_+ W = F$ is called a *Heegaard surface* for M . The *genus* of the splitting (W, W') is the genus of F . The *Heegaard genus* of M , denoted $g(M)$, is the minimal genus among all Heegaard splittings of M .

In our setting, it will be convenient to restrict the class of Heegaard splittings. We shall define a *restricted Heegaard splitting* to be a Heegaard splitting (W, W') , where W is a handlebody. (This notion is closely related to the more standard notion of a “tunnel system”). The *restricted Heegaard genus*, denoted $g_0(M)$, is the minimal genus among all restricted Heegaard splittings. If ∂M has fewer than two components, then $g_0(M) = g(M)$.

Any restricted Heegaard splitting (W, W') determines a collection of finite presentations of $\pi_1 M$. Indeed, if we choose disk systems $\mathcal{D}$ and $\mathcal{E}$ for W and W' , respectively, we obtain a presentation for $\pi_1 M$, whose generators correspond to the elements of $\mathcal{D}$, and whose relators correspond to the attaching maps of the disks in $\mathcal{E}$. We say that the resulting presentation, P , is *induced by* (W, W') . Note that different choices of $\mathcal{D}$ or $\mathcal{E}$ would lead to different presentations, all of which are said to be induced by (W, W') . Any presentation obtained from P by deleting trivial relators is also said to be induced from (W, W') . We say that a presentation for a group is *geometric* if it is induced by a Heegaard splitting of some 3-manifold.

For example, if M is the exterior of the trefoil knot, then $\langle x_1, x_2 | x_1^2 x_2^{-3} \rangle$ is induced by a genus 2 Heegaard splitting of M . Note that the 1-relator presentation $\langle x_1 | id \rangle$ is induced by Heegaard splittings of two different 3-manifolds; it is induced by the genus 1 Heegaard splitting of $S^2 \times S^1$, and also by the genus 1 Heegaard splitting of $D^2 \times S^1$, where, in the latter case, the 2-handle is attached along a trivial curve of the Heegaard torus.

Our main theorem is the following:

Theorem 1.1. *If M is a compact, orientable 3-manifold which fibers over S^1 , then any 1-relator presentation for $\pi_1 M$ is induced by a Heegaard splitting of M .*

If G is any group, we let $rank(G)$ denote the minimal cardinality among all generating sets for G . We have the obvious inequality $rank \pi_1 M \leq g_0(M)$. There are examples where

this inequality is sharp (see [BZ]), however it is an open problem whether equality holds if M is hyperbolic.¹ Theorem 1.1 implies that equality does hold in the case where M is fibered, and G is a 1-relator group. This includes many hyperbolic examples.

Corollary 1.2. *Let M be a compact, orientable 3-manifold which fibers over S^1 , and suppose that $\pi_1 M$ has a 1-relator presentation. Then $\text{rank}(\pi_1 M) = g_0(M)$.*

Remark: The hypotheses of the corollary force $\text{rank}(\pi_1 M) \leq 2$. See Theorem 2.1.

Proof. Let $P = \langle x_1, \dots, x_n | w \rangle$ be the given presentation for $\pi_1 M$. By Theorem 1.1, P is induced by a Heegaard splitting of M .

First, suppose that $w = \text{id}$. There are now two possibilities. If $\partial M = \emptyset$, then we must have $P = \langle x_1 | \text{id} \rangle$, and $M \cong S^2 \times S^1$. Thus $\text{rank} \pi_1 M = g_0(M) = 1$. If $\partial M \neq \emptyset$, then W is a solid torus, and the 2-handle is attached along a trivial curve in ∂W . So in this case, $M \cong D^2 \times S^1$, and again $\text{rank} \pi_1 M = g_0(M) = 1$.

Suppose now that $w \neq \text{id}$. Recall that an element w in a free group F is *primitive* if it is contained in a free basis for F . If w is non-primitive, then $\pi_1 M$ is non-free (see [LS] Chapter II, Proposition 5.10), so $\text{rank}(\pi_1 M) = 2 = g_0(M)$. If w is primitive, then $\pi_1 M \cong \mathbb{Z}$, and $M \cong D^2 \times S^1$ or $S^2 \times S^1$. In this case $\text{rank}(\pi_1 M) = g_0(M) = 1$. $\diamond$

Say that a word w in a free group $F(n)$ is *geometric* if it can be represented by a simple closed curve on the boundary of a genus n handlebody.

Corollary 1.3. *If w is a non-geometric word in x_1, x_2 , and if $\langle x_1, x_2 | w \rangle$ has a map onto $\mathbb{Z}$ with finitely generated kernel, then $\langle x_1, x_2 | w \rangle$ is not the fundamental group of an orientable 3-manifold.*

Remark: For any given word, these two conditions are straightforward to verify. The algorithm to determine if a word in $F(n)$ is geometric is given in [Z], and an implementation, in the 2-generator case, is currently available at www.math.buffalo.edu/~jdmaster. The procedure to determine if $\ker(\phi)$ is finitely generated is discussed in Section 2. For example, Corollary 1.3 implies that the group $\langle x_1, x_2 | x_1 x_2 x_1 x_2 x_1^{-1} x_2^{-2} \rangle$ is not the fundamental group of an orientable 3-manifold.

Proof. (of Corollary 1.3) Suppose the kernel is finitely generated, and $\langle x_1, x_2 | w \rangle = \pi_1 M$ for an orientable 3-manifold M . By Scott's compact core theorem ([Sc]), we may assume that M is compact. By Stallings' Theorem ([S]), M is fibered. Thus, by Theorem 1.1, the given 1-relator presentation for $\pi_1 M$ is geometric. Thus the word w is geometric. $\diamond$

We expect that a version of Theorem 1.1 remains true without the hypothesis that M is fibered.

Conjecture 1.4. *Let M be a compact, orientable 3-manifold. Then*

- A. any 1-relator presentation of $\pi_1 M$ is geometric; and*
- B. if M has no 2-sphere boundary components, then any 1-relator presentation of $\pi_1 M$ is induced by a Heegaard splitting of M .*

Conjecture 1.4 A is equivalent to:

¹Very recently, Abert and Nikolov [AN] have shown that there are in fact hyperbolic examples for which inequality is sharp.

Conjecture 1.5. *The group $\langle x_1, \dots, x_n | w \rangle$ is the fundamental group of an orientable 3-manifold if and only if w is a geometric word.*

Conjecture 1.4 B implies that rank equals restricted Heegaard genus for orientable 3-manifolds with 1-relator fundamental groups. In particular, Conjecture 1.4 B implies the Poincaré Conjecture.

Evidence for Conjecture 1.4 A is given by the manifolds in the SnapPea census. There are 4303 examples given there of 1-relator presentations of orientable-3-manifold groups. We have used a computer to verify that all of these presentations are in fact geometric.

Section 2 contains some background material. The proof of Theorem 1.1 is contained in Section 3. Section 4 contains a general theorem about 1-relator groups, which is motivated by Conjecture 1.4. The final section contains some general discussion about 1-relator groups and Conjecture 1.4.

We wish to thank Tao Li for a useful conversation, and Ilya Kapovich for help with the references.

2 Background

Brown's criterion

Given a 1-relator group $G = \langle x_1, \dots, x_n | w \rangle$, where w is cyclically reduced, and a surjection $\phi : G \rightarrow \mathbb{Z}$. We shall describe a criterion, due to Brown, for determining if $\ker(\phi)$ is finitely generated.

Suppose that the exponent sum on x_1 in w is zero. This can always be arranged, by applying an appropriate free group automorphism to w (see Lemma 3.1). If $n \geq 3$, then it is easy to see that $\ker(\phi)$ is infinitely generated. If $n = 2$, then $\ker(\phi)$ is generated by $\{y_{i,j} = x_1^i x_j x_1^{-i}\}$, and we have $w = \prod y_{\lambda_j, \mu_j}$, for some finite sequence of integers (λ_j) . The following is proved in [B].

Theorem 2.1. *(Brown) $\ker(\phi)$ is finitely generated if and only if $n = 2$, and (λ_j) has a unique minimum and maximum.*

This result follows from Theorem 4.2 (and the discussion immediately preceding it) in [B]. The proof is an application of the Freiheitsatz (Theorem 2.2 below).

Remark: Theorem 2.1 provides another method of showing that certain 1-relator groups are not 3-manifold groups. Suppose G is as above, and that the sequence (λ_j) has

- i. a unique maximum (resp. minimum) and
- ii. a repeated minimum (resp. maximum).

Then G is not the fundamental group of a compact, orientable 3-manifold.

Indeed, if $G = \pi_1 M$ for a compact, orientable 3-manifold M , then the cover $\widetilde{M}$ corresponding to $\ker(\phi)$ has two ends. Assumption i. implies that one of the ends has a finitely generated group. Thus by an argument of Stallings (proof of Theorem 2 of [S]), M is fibered, and so $\ker(\phi)$ is finitely generated. But, by assumption ii and Proposition 2.1, $\ker(\phi)$ is not finitely generated, for a contradiction.

For example, this argument shows that a Baumslag-Solitar group of the type $\langle x_1, x_2 | x_2 x_1 x_2^p x_1^{-1} \rangle$ is not the fundamental group of a compact, orientable 3-manifold if $|p| > 1$ – a special case of a well-known result of Jaco [J].

Magnus's Freiheitssatz

One of the fundamental results about 1-relator groups is the following:

Theorem 2.2. (*Magnus's Freiheitssatz*) Let $G = \langle x_1, \dots, x_n | w \rangle$, where w is cyclically reduced, and let $X \subset \{x_1, \dots, x_n\}$. Suppose that w involves some x_i which is not in X . Then X freely generates a free subgroup of G .

For a proof, see, for example, [LS].

Uniqueness of the relator

The following result is Proposition II.5.8 in [LS]:

Theorem 2.3. (*Magnus*) If two elements w_1 and w_2 have the same normal closure in a free group F , then w_1 is conjugate to w_2 or w_2^{-1} .

In particular, if one fixes a generating set for a 1-relator group, then the relator is determined, up to conjugacy and inverse.

The Moldavansky Splitting

Suppose that $G = \langle x_1, \dots, x_n | w \rangle$, where the exponent sum of x_1 on w is zero. We shall now explain a method, due to Moldavansky, of representing G as an HNN-extension, with a 1-relator vertex group. Further details of the construction are given in [MS].

Let us represent G as the fundamental group of a 2-complex K , consisting of a bouquet of n circles, with a single 2-handle attached, according to the word w . Let $\phi : G \rightarrow \mathbb{Z}$ be the homomorphism which sends any word to the exponent sum of x_1 . Then $\ker(\phi)$ corresponds to an infinite cyclic cover $\tilde{K} \rightarrow K$, as pictured in Fig. 1, and $\pi_1 \tilde{K} = \ker(\phi)$ is generated by elements of the form $y_{i,j} = x_1^i x_j x_1^{-i}$, for $i \in \mathbb{Z}$ and $2 \leq j \leq n$. Let $\theta : \tilde{K} \rightarrow \tilde{K}$ be the covering transformation corresponding to the element x_1 .

Let $\ell : S^1 \rightarrow K$ be the attaching map of the 2-handle, and let $\tilde{\ell} : S^1 \rightarrow \tilde{K}$ be a lift. Then $\tilde{\ell}$ represents some word $\tilde{w}$, in the generators $y_{i,j}$. We may choose the lift so that the i -coordinates of the $y_{i,j}$'s involved in $\tilde{w}$ range from 0 to m .

Consider the connected sub-complex $\overline{K}$ of $\tilde{K}$ which contains the lifted 2-handle, and the 1-cells corresponding to $\{y_{i,j} : 0 \leq i \leq m, 2 \leq j \leq n\}$. Then $K = \overline{K}/(x \equiv \theta x)$, and this gives G the structure of an HNN-extension (see Fig. 1).

Proceeding algebraically now, consider the presentation:

$$(*) \quad \langle t, y_{i,j} \mid 0 \leq i \leq m, 2 \leq j \leq n | \tilde{w}, t y_{i,j} t^{-1} = y_{i+1,j} \mid 0 \leq i \leq m-1, 2 \leq j \leq n \rangle.$$

Then, by mapping t to x_1 and $y_{i,j}$ to $x_1^i x_j x_1^{-i}$, one may check that this is a presentation of G .

Now, let G_v be the following 1-relator group:

$$G_v = \langle y_{i,j} \mid 0 \leq i \leq m, 2 \leq j \leq n | \tilde{w} \rangle,$$

and let G_e be a free group:

$$G_e = \langle y_{i,j} \mid 0 \leq i \leq m-1, 2 \leq j \leq n \rangle.$$

(We have abused notation by re-using the symbol $y_{i,j}$.) Define maps $i^\pm : G_e \rightarrow G_v$, by $i^+ y_{i,j} = y_{i,j}$, and $i^- y_{i,j} = y_{i+1,j}$. By Theorem 2.2, these are both injections. Thus the




Figure 1: Construction of the Moldavsky splitting

presentation (*) expresses G as an HNN-extension with vertex group G_v , and edge group G_e .

For an illustration, suppose that $G = \langle x_1, x_2 | x_2 x_1 x_2^2 x_1 x_2^3 x_1^{-2} \rangle$. Then G is an HNN-extension, with vertex group $G_v = \langle y_{0,1}, y_{1,1}, y_{2,1} | y_{0,1} y_{1,1}^2 y_{2,1}^3 \rangle$, edge group $G_e = \langle y_{0,1}, y_{1,1} \rangle$, and inclusions $i^+ : G_e \rightarrow G_v$, sending $y_{i,j} \rightarrow y_{i,j}$ and $i^- : G_e \rightarrow G_v$, sending $y_{1,1} \rightarrow y_{2,1}$ and $y_{2,1} \rightarrow y_{3,1}$.

Heegaard splittings of bundles

Suppose M is a fibered 3-manifold, with a fiber F , which is a compact surface with non-empty boundary, and monodromy $f : F \rightarrow F$. Let $\{x_1, \dots, x_n\}$ be a free generating set for $\pi_1 F$, represented by a collection of embedded loops in F , meeting at a single point. Then $\pi_1 M$ has an HNN presentation:

$$(**) \quad \pi_1 M = \langle x_1, \dots, x_n, t | x_1 = t(f_* x_1) t^{-1}, \dots, x_n = t(f_* x_n) t^{-1} \rangle.$$

We have the following fact.

Lemma 2.4. *The presentation (**) for $\pi_1 M$ is induced by a Heegaard splitting of M .*

Proof. Let F_1 and F_2 be disjoint fibers in M . Let α be an arc, whose interior is disjoint from $F_1 \cup F_2$, with one endpoint in F_1 and the other endpoint in F_2 . Let S be the boundary of a regular neighborhood of $F_1 \cup F_2 \cup \alpha$. Then one may verify that S is a Heegaard surface inducing the presentation (*). $\diamond$

Modifying Heegaard splittings

Suppose (W, W') is a Heegaard splitting of M . We may obtain a new Heegaard splitting for M by attaching a trivial 1-handle to W , and drilling out the corresponding 1-handle from W' . We say that the new Heegaard splitting is a *stabilization* of (W, W') .

Suppose that there are essential disks $D \subset W$, and $D' \subset W'$, such that $|\partial D \cap \partial D'| = 1$. Let $C \subset \partial_+ W$ denote the boundary of a regular neighborhood of $\partial D \cup \partial D'$. Then there is a 2-sphere $S \subset M$, such that $S \cap \partial_+ W = C$. Let E be the disk $S \cap W$. Then we may form a new Heegaard surface for M by compressing $\partial_+ W$ along E . We say that the resulting Heegaard splitting is a *destabilization* of (W, W') .

There is a useful method for changing disk systems. Suppose $\mathcal{D}$ is a disk system for W , and that $D_1, D_2 \in \mathcal{D}$. Suppose that α is an arc in ∂W , starting on ∂D_1 and ending on ∂D_2 , with $\text{int } \alpha \cap D = \emptyset$ for all $D \in \mathcal{D}$. Let $\alpha \times I \subset \partial W$ be an embedded neighborhood of α , so that $(\alpha \times I) \cap \partial D_i$ is a neighborhood of $\alpha \cap D_i$, and $(\alpha \times I) \cap \partial D = \emptyset$ for $D \in \mathcal{D} - (D_1 \cup D_2)$. Then the *band sum* of D_1 and D_2 along α is the disk $E = D_1 \cup D_2 \cup N(\alpha)$. The set $\mathcal{E} = (\mathcal{D} - D_1) \cup E$ is then a new disk system for W .

Modifying group presentations

Suppose $P = \langle X | R \rangle$ is a presentation for a group G , where $X = \{x_1, \dots, x_m\}$ and $R = \{r_1, \dots, r_n\}$. One may obtain a new generating set by applying a free group automorphism to the x_i 's. We say that the resulting presentations are *Nielsen equivalent*.

For example, if $r \neq s$, and $1 \leq r, s \leq m$, let $\tau_{rs} : F(x_1, \dots, x_m) \rightarrow F(x_1, \dots, x_m)$ be the automorphism satisfying $\tau_{rs}x_r = x_r x_s$, and $\tau_{rs}x_i = x_i$ if $i \neq r$.

Let $\bar{x}_i = \tau_{rs}x_i$, and let $\bar{r}_i$ be the word obtained from r_i , by replacing x_r with $\bar{x}_r \bar{x}_s^{-1}$ and x_i with $\bar{x}_i$, for $i \neq r$. Then $\bar{P} = \langle \bar{x}_1, \dots, \bar{x}_m | \bar{r}_1, \dots, \bar{r}_n \rangle$ is a presentation for G .

Let $\tau_i : F(x_1, \dots, x_m) \rightarrow F(x_1, \dots, x_m)$ which sends x_i to x_i^{-1} , and stabilizes all the other generators. In this case, the new presentation for G is obtained from P by exchanging x_i and x_i^{-1} in all the relators $r_1, \dots, r_n$.

The automorphisms τ_{rs} and τ_i are called *Nielsen transformations* of $F(x_1, \dots, x_m)$. The operation which a Nielsen transformation induces on a group presentation is called a *Nielsen move*. For a proof of the following fact, see [LS], Chapter I.

Proposition 2.5. *The Nielsen transformations generate $\text{Aut}(F(x_1, \dots, x_m))$. In particular, if $\bar{P}$ is Nielsen equivalent to P , then $\bar{P}$ may be obtained from P by a sequence of Nielsen moves.*

We shall need the fact that Nielsen moves can be realized geometrically. The proof of the following proposition may be found in [G].

Proposition 2.6. *(Griffiths) Let $F(n) = \pi_1(W)$, for a handlebody W . Then every automorphism of $F(n)$ is induced by an automorphism of W .*

In addition to the Nielsen moves, it will be convenient to define the following operations on P (special examples of Tietze transformations), which give new presentations of $\pi_1 M$:

1. Add a new element x' to X , and also add x' to R . We say that P has been **stabilized**.
2. Suppose that there is a relation (not necessarily part of the presentation) in the group of the form $r = x^{-1}w$, where $x \in X$ and w is a word which does not involve x . Then remove x from X , remove r from R (if r is in R), and, wherever x appears in any relator $r' \in R - r$, replace it with w . We say that P has been **destabilized along x** .

Modifying presentations of 3-manifold groups

Suppose that P is a presentation for $\pi_1 M$ induced by a Heegaard splitting (W, W') of M ,

where W is a handlebody. By Proposition 2.6, any automorphism of $\pi_1 W$ is induced by a handlebody automorphism. Thus, if $\bar{P}$ is obtained from P by a Nielsen move, then $\bar{P}$ is also induced by the splitting (W, W') . It is easy to see that a stabilization of the splitting (W, W') , results in a stabilization of P . Similarly, a destabilization of (W, W') corresponds to a destabilization of P , as described in the following lemma.

Lemma 2.7. *Let (W, W') be a Heegaard splitting of M , let $\mathcal{D}$ and $\mathcal{D}'$ be disk systems for W and W' , respectively, and let $P = \langle x_1, \dots, x_g | R_1, \dots, R_s \rangle$ be the induced presentation of $\pi_1 M$. Let $D_1 \in \mathcal{D}$ be the disk which is dual to the generator x_1 , and suppose $D'_1 \in \mathcal{D}'$, with $|D_1 \cap D'_1| = 1$. Then P can be destabilized along x_1 so that the resulting presentation is induced by a destabilization of (W, W') .*

Proof. Let $C \subset \partial W$ be the boundary of a regular neighborhood of $\partial D_1 \cup \partial D'_1$. Then C bounds a disk in both W and in W' , so there is a sphere $S \subset M$ with $S \cap \partial W = C$. Let B be the ball bounded by S with $D_1 \cup D'_1 \subset B$. Then S is a reducing sphere for (W, W') , resulting in a destabilized Heegaard splitting (Z, Z') , where $Z = W \cup B$, and $Z' = \overline{W' - B}$.

We obtain a disk system for Z as follows. Let $\{p_1, \dots, p_n\} = \bigcup_{D_j \in \mathcal{D}} D_j \cap D'_1$, where $p_1 = D_1 \cap D'_1$, and the p_i 's are ordered consecutively along the circle $\partial D'_1$. Suppose $n \geq 2$, and let $D_j \in \mathcal{D}$ be the disk containing p_2 . Let α be an arc of $\partial D'_1$ with endpoints p_1 and p_2 , which is disjoint from p_i for $i \geq 3$. Then we replace D_j with the band sum of D_j and D_1 along α . Let $\mathcal{E}$ be the resulting disk system for W , and then $|\bigcup_{E \in \mathcal{E}} E \cap D'_1| = n - 1$. The resulting presentation of $\pi_1 M$ is obtained from P by applying a single Nielsen move—replacing the generator x_1 with $x_1 x_j^{\pm 1}$. Continuing, we obtain a disk system $\mathcal{E}$ for W , so that $E \cap D'_1 = \emptyset$ for all $E \in \mathcal{E} - D_1$. Then $\mathcal{E} - D_1$ is a system for Z . Let P^* be the presentation of $\pi_1 M$ induced by the systems $\mathcal{E}$ and $\mathcal{D}'$. Then P^* is obtained from P by applying a sequence of Nielsen moves, whose effect is to replace the generator x_1 with the relator corresponding to $\partial D'_1$. In P^* , the disk D'_1 corresponds to the relation $x_1 = id$.

We obtain a disk system for Z' as follows. Let $\{p_1, \dots, p_n\} = \bigcup_{D'_j \in \mathcal{D}'} D'_j \cap D_1$, where $p_1 = D'_1 \cap D_1$, and the p_i 's are ordered consecutively along the circle ∂D_1 . Suppose $n \geq 2$, and let $D'_j \in \mathcal{D}'$ be the disk containing p_2 . Let α be an arc of ∂D_1 with endpoints p_1 and p_2 , which is disjoint from p_i for $i \geq 3$. Then we replace D'_j with the band sum of D'_j and D'_1 along α . Let $\mathcal{E}'$ be the resulting disk system for W' , and then $|\bigcup_{E \in \mathcal{E}'} E \cap D_1| = n - 1$. The presentation of $\pi_1 M$ induced by $\mathcal{E}$ and $\mathcal{E}'$ is obtained from P^* by removing one occurrence of x_1 from one relation. Continuing, we obtain a disk system $\mathcal{E}'$, so that $D'_1 \in \mathcal{E}'$, and $E \cap D_1 = \emptyset$ for all $E \in \mathcal{E}' - D'_1$. Then $\mathcal{E}' - D'_1$ is a system for Z' . Let Q be the presentation of $\pi_1 M$ induced by the systems $\mathcal{E} - D_1$ and $\mathcal{E}' - D'_1$. Then Q is obtained from P^* by removing the generator x_1 , the relator $x_1 = id$, and all occurrences of x_1 in the remaining relators of P^* . It follows that Q is obtained from P by destabilizing along x_1 . $\diamond$

To prove Theorem 1.1, we require a technical version of Lemma 2.7 that works for immersed disks.

Lemma 2.8. *Let (W, W') be a Heegaard splitting of M , let $\mathcal{D}$ and $\mathcal{D}'$ be disk systems for W and W' , respectively, and let $P = \langle x_1, \dots, x_g | R_1, \dots, R_s \rangle$ be a presentation of $\pi_1 M$, where the generators are dual to the disks in $\mathcal{D}$, and each R_i is represented by an immersed disk in W' . Let $D_1 \in \mathcal{D}$ be the disk which is dual to the generator x_1 . Suppose there is a compressing disk D' for W' , so that $|D_1 \cap D'| = 1$. Then (W, W') can be destabilized to a splitting (Z, Z') , and P can be destabilized along x_1 to a presentation Q , where the*

generators of Q are dual to a disk system for Z , and the relations of Q correspond to immersed disks in Z' .

Proof. The proof is similar to the proof of Lemma 2.7. The disks D_1, D' determine a reducing sphere S , which bounds a ball B , giving a destabilized Heegaard splitting (Z, Z') , where $Z = W \cup B$, and $Z' = \overline{W'} - \overline{B}$. We obtain a disk system for Z as above.

If D'_i is an immersed disk representing R_i , we can remove intersection points of $\partial D'_i \cap \partial D_1$, by banding together D'_i and D' , along a subarc of ∂D_1 . Unlike the situation in Lemma 2.7, we also have to consider the possibility that D'_i meets D' . However, these intersections can be removed by banding together D'_i and D_1 , along a subarc of $\partial D'$. The result is an immersed disk E'_i in Z' , and the word represented by $\partial E'_i$ is the same as the word represented by $\partial D'_i$. Repeating this procedure for each E_i , we obtain immersed disks in the handlebody Z' , representing the relators in a presentation Q , obtained by destabilizing P along x_1 . $\diamond$

3 Proof of Theorem 1.1

Suppose

$$P = \langle x_1, \dots, x_n | w \rangle \cong G = \pi_1 M,$$

for some compact, orientable, fibered 3-manifold M . The first step is to put the word w into a standard form. We let $e_i(w)$ denote the exponent sum of the letter x_i in the word w . We may assume that $n \geq 2$, for otherwise $G \cong \mathbb{Z}$ and $w = id$, so M is either $S^2 \times S^1$ or $D^2 \times S^1$, and the presentation is induced by a genus 1 Heegaard splitting of M .

Since M is fibered, there is a map $\phi : G \rightarrow \mathbb{Z}$ with finitely generated kernel, and so, by Theorem 2.1, we may assume that $n = 2$, and the given presentation has the form:

$$(1) \quad \pi_1 M = \langle x_1, x_2 | w \rangle.$$

Suppose that w is primitive. Then (by Proposition 2.6), w is geometric, and $\pi_1 M \cong \mathbb{Z}$. Then $M \cong S^2 \times S^1$ or $D^2 \times S^1$, and the presentation is induced by a Heegaard splitting of M . So assume, from now on, that w is non-primitive.

Lemma 3.1. *There is an automorphism θ of the free group on x_1, x_2 , such that $\phi(\theta(x_1)) = 0, \phi(\theta(x_2)) = 1$.*

Proof. Let $\phi x_1 = \alpha$ and $\phi x_2 = \beta$. Let σ be the Nielsen transformation which stabilizes x_2 , and satisfies $\sigma x_1 = x_1 x_2$. Let τ be the Nielsen transformation which stabilizes x_1 , and satisfies $\tau(x_2) = x_2 x_1$. We have:

$$\begin{aligned} \phi(\sigma^{\pm 1} x_1) &= \alpha, \\ \phi(\sigma^{\pm 1} x_2) &= \beta \pm \alpha, \\ \phi(\tau^{\pm 1} x_1) &= \alpha \pm \beta, \\ \phi(\tau^{\pm 1} x_2) &= \beta. \end{aligned}$$

Thus by applying the Euclidean algorithm to (α, β) , we may find an automorphism θ , such that $\phi(\theta x_1) = 0$, and $\phi(\theta x_2) = \pm 1$. Replacing x_2 with x_2^{-1} if necessary, we have $\phi(\theta x_2) = 1$.

$\diamond$

We now return to the proof of Theorem 1.1. Let θ be the automorphism given by Lemma 3.1. By Proposition 2.6, the presentation $\langle x_1, x_2 | w \rangle$ is induced by a Heegaard splitting of M if and only if the presentation $\langle x_1, x_2 | \theta(w) \rangle$ is induced by a Heegaard splitting of M . Thus we may replace w with $\theta(w)$, and so assume that $\phi(x_1) = 0$ and $\phi(x_2) = 1$.

Let $y_i = x_2^i x_1 x_2^{-i}$. Since $\phi(w) = 0$, then we can represent the word w in the free group $F(x_1, x_2)$ as a product of y_i 's: i.e. $w = y_{r_1} \dots y_{r_k}$. Replacing w by a conjugate, we may assume that $\text{Min}\{r_1, \dots, r_k\} = 0$, and we let $n = \text{Max}\{r_1, \dots, r_k\}$. Then $w = w_2(y_0, \dots, y_n)$, for some word w_2 . Since ϕ has finitely generated kernel, then by Proposition 2.1, the letters y_0 and y_n appear only once each in the word w_2 . In particular, the word w_2 is primitive. Since w is non-primitive, we must have $n \geq 1$.

As explained in the previous section, the 1-relator group G has a Moldavansky splitting. In our case, the splitting has the following structure: the vertex group is $\mathcal{G}_v = \langle y_0, \dots, y_n | w_2 \rangle \subset G$; the stable letter is $t = x_2$; the edge group is $\mathcal{G}_e = \langle y_0, \dots, y_{n-1} \rangle \subset G$. The inclusion maps are:

$$\begin{aligned} i_+ : \mathcal{G}_e &\rightarrow \mathcal{G}_v \\ i_+(y_j) &= y_j; \\ i_- : \mathcal{G}_e &\rightarrow \mathcal{G}_v, \\ i_-(y_j) &= y_{j+1}. \end{aligned}$$

This splitting is represented by a triple $\mathcal{S} = (\mathcal{G}_v, \mathcal{G}_e, t)$. Since y_n appears only once in w_2 , then in fact $\mathcal{G}_v = \mathcal{G}_e$, and the splitting $\mathcal{S}$ decomposes G as a mapping torus of a free group automorphism $\psi : \mathcal{G}_e \rightarrow \mathcal{G}_e$. In particular, we have that $\mathcal{G}_e = \text{Ker}(\phi)$.

If $F \subset M$ is a non-separating incompressible surface, with dual loop ℓ , we let F^+ and F^- be the corresponding subsurfaces of $\partial(\overline{M-F})$. Let $p = \ell \cap F$ be a base-point for $\pi_1 F$, and let $p^\pm$ be the pre-images of p in $\partial(\overline{M-F})$. Let $j_+ : \pi_1(F^+, p^+) \rightarrow \pi_1(\overline{M-F}, p^+)$ be the map induced by inclusion. We say that the splitting $\mathcal{S}$ is *realized geometrically* if F and ℓ can be chosen so that $i_*(\overline{M-F}, p^+) = \mathcal{G}_v \subset \pi_1 M$, $j_{+*}(\pi_1 F^+, p^+) = \mathcal{G}_e$, and $[\ell] = t$.

Claim: The splitting $\mathcal{S}$ is realized geometrically.

Proof of claim:

The map $\phi : \pi_1 M \rightarrow \mathbb{Z}$ is dual to a fiber F , and the element $t = x_2$ is represented by a dual loop ℓ for F . Then $j_{+*}\pi_1(F^+, p^+) = \text{ker}(\phi) = \mathcal{G}_e$. Also, $\overline{M-F} \cong F \times [0, 1]$, and so $i_* : (\pi_1 \overline{M-F}, p^+) = \mathcal{G}_e = \mathcal{G}_v$, as required.

By the claim, the fiber of M has free fundamental group, and therefore M must have non-empty boundary. (In fact, by a result of Lyndon ([L]), any compact, orientable, irreducible 3-manifold with a 1-relator fundamental group must have non-empty boundary.)

We now wish to write down an explicit presentation for G , corresponding to the splitting $\mathcal{S}$. First, we may conjugate and invert w_2 , if necessary, so that it has the form $w_2 = y_n^{-1} w_3$, for some word w_3 which does not involve y_n . We may then eliminate the generator y_n from our presentation for G_v , obtaining the following presentation for $\pi_1 M$:

$$(2) \quad \pi_1 M \cong \langle y_0, \dots, y_{n-1}, t | y_0 t y_1^{-1} t^{-1}, \dots, y_{n-2} t y_{n-1}^{-1} t^{-1}, y_{n-1} t w_3^{-1} t^{-1} \rangle.$$

Our next task is to show that the presentation (2) is induced by a Heegaard splitting. By the claim, there is a fiber surface F in M , inducing the splitting $\mathcal{S}$, and hence inducing

the presentation (2). The difficulty is that, *a priori*, the given generators $y_0, \dots, y_{n-1}$ of the fiber group may not be representable by embedded loops in F .

Let (W, W') be the genus $n+1$ splitting for M given by Lemma 2.4. Let $\mathcal{D} = \{D_0, \dots, D_n\}$ and $\mathcal{E} = \{E_0, \dots, E_{n-1}\}$ be disk systems for W and W' , respectively, inducing a presentation of the following form:

$$(3) \quad \pi_1 M = \langle z_0, \dots, z_{n-1}, s | s z_0 s^{-1} f_* z_0^{-1}, \dots, s z_{n-1} s^{-1} f_* z_{n-1}^{-1} \rangle.$$

(So y_i is dual to D_i for $i \leq n-1$, and s is dual to D_n .) Since s and t are both represented by loops which are dual to F , then $s = tu$ for some $u \in i_* \pi_1 F$. It follows that f_* is the composition of ψ with an inner automorphism. Thus there is a fiber-preserving automorphism of M which takes the presentation (3) to:

$$(4) \quad \pi_1 M = \langle z_0, \dots, z_{n-1}, t | t z_0 t^{-1} \psi z_0^{-1}, \dots, t z_{n-1} t^{-1} \psi z_{n-1}^{-1} \rangle.$$

Thus presentation (4) is induced by a Heegaard splitting, which we still refer to as (W, W') . We also replace the disk systems $\{D_i\}$ and $\{E_i\}$ with their images under the automorphism.

Since $y_0, \dots, y_{n-1}$ and $z_0, \dots, z_{n-1}$ are both free bases for $i_* \pi_1 F$, then by Proposition 2.5 there is a sequence of Nielsen transformations taking $z_0, \dots, z_{n-1}$ to $y_0, \dots, y_{n-1}$. Suppose the first such move sends z_i to $z_i z_j$, and stabilizes all the other generators. Then the corresponding Nielsen move, applied to presentation (4), gives:

$$\pi_1 M = \langle \bar{z}_0, \dots, \bar{z}_{n-1}, t | t \bar{z}_0 t^{-1} \psi \bar{z}_0^{-1}, \dots, t \bar{z}_i \bar{z}_j^{-1} t^{-1} \psi (\bar{z}_i \bar{z}_j^{-1})^{-1}, \dots, t \bar{z}_{n-1} t^{-1} \psi \bar{z}_{n-1}^{-1} \rangle.$$

We now perform a Tietze transformation, multiplying (a conjugate of) the i th relation by (a conjugate of) the j th relation, to obtain the following presentation:

$$\pi_1 M = \langle \bar{z}_0, \dots, \bar{z}_{n-1}, t | t \bar{z}_0 t^{-1} \psi \bar{z}_0^{-1}, \dots, t \bar{z}_i t^{-1} \psi \bar{z}_i^{-1}, \dots, t \bar{z}_{n-1} t^{-1} \psi \bar{z}_{n-1}^{-1} \rangle.$$

To perform this move geometrically, we choose an (immersed) arc $\alpha \subset \partial W$, with one endpoint on ∂E_i and the other on ∂E_j , which is disjoint from all the disks $D_1, \dots, D_n$. Then we attach a regular neighborhood of α to E_i and E_j , obtaining an immersed disk whose boundary represents the relator $t \bar{z}_i t^{-1} \psi \bar{z}_i^{-1}$. Continuing, we may change the presentation (4) to the presentation (2), and we see that the relators $R_0, \dots, R_{n-1}$ of (2) are represented by *immersed* disks $E_0^\dagger, \dots, E_{n-1}^\dagger$, in W' .

By Dehn's Lemma, there is a compressing disk E_0^* for W' , so that $\partial E_0^* \subset \partial E_0^\dagger$. Let w^* be the word represented by ∂E_0^* .

Claim: The circle ∂E_0^* meets ∂D_0 in a single point.

Before proving the claim, we make a definition.

Definition: For any word u on a generating set $\mathcal{G}$, and any $g \in \mathcal{G}$, we define $n_g(u)$ to be the number of times the generator g , or its inverse, occurs in u . Note this is different from the exponent sum. For example, we have $n_t(R_0) = n_t(y_0 t y_1^{-1} t^{-1}) = 2$. Since $\partial E_0^* \subset \partial E_0^\dagger$, then $|\partial E_0^* \cap D_i| \leq |\partial E_0^\dagger \cap D_i|$ for each i . Therefore, for each generator g in (2), we have $n_g(w^*) \leq n_g(R_0) = n_g(y_0 t y_1^{-1} t^{-1})$.

Proof of claim: We have $|\partial E_0^* \cap \partial D_0| \leq |\partial E_0^\dagger \cap \partial D_0| = 1$. Suppose $\partial E_0^* \cap \partial D_0 = \emptyset$.

Thus $n_{y_0}(w^*) = 0$. Since $n_g(w^*) \leq n_g(y_0 t y_1^{-1} t^{-1})$ for each generator g , then $n_{y_1}(w^*) \leq 1$, $n_t(w^*) \leq 2$, and $n_g(w^*) = 0$ for all other generators g . Furthermore, since w^* represents the trivial element in $\pi_1 M$, the exponent sum of t in w^* must be zero.

We see that the only possibilities are that w^* is trivial, or else conjugate to $y_1^{\pm 1}$. The latter case is impossible, since the element y_1 is contained in a basis for the free group $\pi_1 F$. So w^* is the trivial word. But then there is a reducing sphere S for (W, W') which is disjoint from all the disks $D_0, \dots, D_n$; therefore some non-empty subset of the generators $y_0, \dots, y_{n-1}, t$ must be trivial in $\pi_1 M$. This is impossible, since $\{y_0, \dots, y_{n-1}\}$ is a basis for the free group $\pi_1 F$. This proves the claim.

By Lemma 2.8, we can destabilize (W, W') to a splitting (Z, Z') , and we can destabilize the presentation P along y_1 to obtain a presentation Q , where the relators of Q are represented by immersed disks in Z' . The presentation Q has a relator of the form $y_2 t y_3^{-1} t^{-1}$. Since y_2 appears once in this relator, we can repeat the previous argument, to obtain a reducing sphere for the splitting (Z, Z') , and a destabilization of the presentation Q along y_2 .

Continuing, we may reduce the Heegaard splitting (W, W') to a splitting (Z, Z') , where $\text{genus}(Z) = 2$, and $\pi_1 Z$ is generated by $y_{n-1} = x_2^{n-1} x_1 x_2^{-(n-1)}$ and $t = x_2$. Applying an automorphism of the handlebody Z , we may take the generators of $\pi_1 Z$ to be x_1, x_2 , and then the Heegaard splitting (Z, Z') determines a 1-relator presentation:

$$(6) \quad \pi_1 M \cong \langle x_1, x_2 | \overline{w} \rangle,$$

where $\overline{w}$ is a geometric word. Comparing the presentation (6) with the original presentation (1), and applying Theorem 2.3, we see that w is conjugate to $\overline{w}$ or $\overline{w}^{-1}$ in the free group $\langle x_1, x_2 \rangle$. Therefore w is geometric, and the given presentation (1) is induced by a Heegaard splitting of M .

4 Hierarchies for 1-relator groups

It was observed in [K] that 1-relator groups and Haken 3-manifolds share some common features. In this section, we shall make the analogy explicit, showing that every 1-relator group possesses a “1-relator hierarchy”.

Let G be a 1-relator group. We define a *1-relator hierarchy* for G to be a finite sequence of 1-relator groups $G_1, G_2, \dots, G_n$, such that $G_1 \cong G$, G_n is cyclic, and G_i splits as an HNN-extension with vertex group G_{i-1} .

Theorem 4.1. *Let G be a 1-relator group. Then G has a 1-relator hierarchy.*

Proof. The proof is by a “tower argument”, similar in spirit to the proof of Dehn’s Lemma. First we require some terminology. By a *loop* in a surface S , we understand an immersion from a circle into S . A loop in S is *essential* if it cannot be homotoped into ∂S . A loop in S is said to *fill* S if the complement of its image contains no essential loops.

Given a 1-relator group $G_1 = \langle x_{1,1}, \dots, x_{1,n_1} | w_1 \rangle$, with $n_1 \geq 2$. We view the free group on $x_{1,1}, \dots, x_{1,n_1}$ as $\pi_1 S_1$, for some compact surface S_1 , where the x_i ’s are represented by embedded loops meeting in a single point.

The word w_1 is represented by an immersed loop $\ell_1 \subset S_1$, and if w_1 is cyclically reduced, we may take ℓ_1 to minimize the number of self-intersections in its free homotopy class.

Case 1: The loop ℓ_1 does not fill S_1 .

In this case, there is an automorphism $\theta : S_1 \rightarrow S_1$, such that some generator $x_{1,i}$ does not appear in w_1 . After re-labeling, we may assume that the missing generator is $x_{1,1}$. We choose such a θ , and we replace the presentation $\langle x_{1,1}, \dots, x_{1,n_1} | w_1 \rangle$ with the presentation $\langle x_{1,1}, \dots, x_{1,n_1} | \theta_*(w_1) \rangle$, which defines an isomorphic group. We also replace ℓ_1 with $\theta(\ell_1)$.

The group G_1 then splits as a free product $G_1 \cong G_2 * \langle x_{1,1} \rangle$. We relabel the ordered list $(x_{1,2}, \dots, x_{1,n_1})$ by the ordered list $(x_{2,1}, \dots, x_{2,n_2})$, and let w_2 be the result of re-labeling $\theta(w_1)$. Then we have $G_2 \cong \langle x_{2,1}, \dots, x_{2,n_2} | w_2 \rangle$, where $n_2 = n_1 - 1$. We view this free product as an HNN-extension, with trivial edge groups.

There is a sub-surface $S_2 \subset S_1$, such that

$$i_* \pi_1 S_2 = \langle x_{2,1}, \dots, x_{2,n_2} \rangle \subset \pi_1 \tilde{S}_1.$$

The word w_2 is represented by a loop $\ell_2 \subset S_2$. In fact, in this case, we may choose $\ell_2 = \ell_1$.

Case 2: The loop ℓ_1 fills S_1 .

Since $b_1(S_1) \geq 2$, there is a map $\phi : G_1 \rightarrow \mathbb{Z}$, such that $w_1 \in \text{Ker}(\phi)$. By changing the generators of $\pi_1 S$, and re-writing w_1 in terms of the new generators, we may assume that ϕ is the exponent-sum function $e_{x_{1,1}}$.

Corresponding to the homomorphism $e_{x_{1,1}} : G_1 \rightarrow \mathbb{Z}$, there is an infinite cyclic cover $\tilde{S}_1 \rightarrow S_1$. Let $\tau : \tilde{S}_1 \rightarrow \tilde{S}_1$ be the corresponding covering transformation. Let $D \subset \tilde{S}_1$ be a compact fundamental domain for the action of τ . Let k_1 be minimal such that the surface $S_2 = D \cup tD \cup \dots \cup t^{k_1} D$ contains a lift of ℓ_1 . We have:

$$\pi_1 S_2 = \langle x_{1,2}, \dots, x_{1,n_1}, x_{1,1} x_{1,2} x_{1,1}^{-1}, \dots, x_{1,1}^{k_1} x_{1,n_1} x_{1,1}^{-k_1} \rangle.$$

We may choose these generators so they are represented by embedded loops in S_2 , meeting at a single point. We label the elements of this list consecutively as $(x_{2,1}, \dots, x_{2,n_2})$. Then we see (as in Section 2) that G_1 is an HNN-extension with vertex group

$$G_2 = \langle x_{2,1}, \dots, x_{2,n_2} | w_2 \rangle,$$

for some word w_2 , represented by a loop $\ell_2 \subset S_2$, which is a lift of ℓ_1 . The edge group is the free group $F(r_2)$, where $r_2 = (n_1 - 1)k_1$, and the images of the edge group are generated by $\{x_{2,1}, \dots, x_{2,r_2}\}$ and $\{x_{2,n_1}, \dots, x_{2,n_2}\}$, respectively. By the Freiheitsatz, these are in fact free subgroups of G_2 .

Suppose that by the above procedure we have constructed a splitting of G_1 , with vertex group $G_2 = \langle x_{2,1}, \dots, x_{2,n_2} | w_2 \rangle$, where G_2 is represented by a loop ℓ_2 in a surface S_2 . If $n_2 \geq 2$, we may apply the above process to G_2 . Continuing, we obtain a sequence of 1-relator groups $G_1, G_2, \dots$, where $G_i = \langle x_{i,1}, \dots, x_{i,n_i} | w_i \rangle$, and G_i is a vertex group for G_{i-1} . The sequence terminates at G_k if and only if $n_k = 1$.

Topologically, for each i we have surfaces S_i and $\tilde{S}_i$, (where possibly $\tilde{S}_i = S_i$) so that S_{i+1} is a subsurface of $\tilde{S}_i$. We also have a loop $\ell_i \subset S_i$ representing the word w_i , such that ℓ_{i+1} is a homeomorphic lift of ℓ_i .

Claim: The sequence $G_1, G_2, \dots$ terminates after finitely many steps.

Proof of claim:

Note that when a curve is lifted to a finite cover, the number of self-intersections can never go up. Therefore the number of self-intersections of ℓ_{i+1} is less than or equal to the number of self-intersections of ℓ_i for all i . Thus if the sequence $G_1, G_2, \dots$ is infinite, then the sequence $\ell_1, \ell_2, \dots$ must contain infinitely many consecutive terms for which the number of self-intersections remains constant. We claim that this is impossible.

Indeed, suppose the number of self-intersections of ℓ_{i+1} is the same as the number of self-intersections of ℓ_i , and suppose that $\ell_{i+1} \subset S_{i+1}$ is constructed as in Case 2. Then the lifts of ℓ_i to $\tilde{S}_i$ are all disjoint. Let C be the boundary of a regular neighborhood of one of these lifts in $\tilde{S}_i$. Since each lift of ℓ_i is essential, and since no lift of ℓ_i fills $\tilde{S}_i$, then some component of C must be essential. Thus C projects to a collection of loops in S_i , which are disjoint from ℓ_i , at least one of which must be essential. Thus ℓ_i does not fill S_i . Therefore, G_{i+1} is constructed from G_i as in Case 1, and so $n_{i+1} = n_i - 1$.

Similarly, if the number of self-intersections of $\ell_i, \dots, \ell_{i+k}$ are all the same, then $n_{i+k} = n_i - k$. Thus if the sequence of G_i 's is infinite, we arrive at a contradiction. $\diamond$

5 Concluding remarks

5.1 On 1-relator groups

Often, facts about 1-relator groups are proved by induction on the length of the relator. Proofs break into two cases: if $G = \langle x_1, \dots, x_n | w \rangle$, and the word w has zero exponent sum on one of its letters, then $G \cong G_2 *_t$, where G_2 is a 1-relator group, whose relator is shorter than the relator of G . One may then apply the induction hypothesis and the HNN structure to prove things about G .

If w does not have this form, one shows that G embeds nicely in a certain 1-relator group $\hat{G}$, with $\hat{G} \cong \hat{G}_2 *_t$; where $\hat{G}_2$ is a 1-relator group, whose relator is shorter than w . One may then apply the induction hypothesis to $\hat{G}_2$, and attempt to transfer this information to G .

Theorem 4.1 makes it possible to prove facts about 1-relator groups by inducting, instead, on the length of a hierarchy. This allows one to avoid the embedding step, and give proofs which are, to a 3-manifold topologist, more intuitive.

5.2 On Conjecture 1.4

Our main motivation for presenting Proposition 4.1 is provided by Conjecture 1.4. The standard method for 1-relator groups appears problematic in this case, since there is no guarantee that the embedding step can be performed geometrically. Proposition 4.1 gives hope that the conjecture might be approachable by inducting, instead, on the length of a hierarchy.

However, the proof given for the fibered case does not directly generalize. Indeed, a key point in the proof of Theorem 1.1 is that if M is fibered, then the Moldavansky splitting is realized geometrically. But, in general, it is *not* true that the Moldavansky splitting of a 1-relator 3-manifold group is geometric.

Let M be the manifold $M017$ in the Snappea census. Then the presentation

$$\pi_1 M = \langle x_1, x_2 | x_1^2 x_2 x_1^3 x_2 x_1^2 x_2^{-2} \rangle$$

is induced by a Heegaard splitting of M . Since $b_1(M) = 1$ and $\partial M \cong T^2$, there is a unique (up to isotopy) longitude curve $\lambda \subset \partial M$, whose image is trivial in $H_1(M, \mathbb{Q})$. One may compute that in $\pi_1 M$, the loop λ represents the element $x_2^{-4} x_1^2 x_2 x_1^2 (x_1^3 x_2 x_1^2)^3$. Thus $[\lambda]$ has order 7 in $H_1(M, \mathbb{Z}) \cong \mathbb{Z} \times \mathbb{Z}/7$. It follows that any non-separating surface in M must have at least seven boundary components, and thus have a fundamental group of rank at least 6. However, for this example, the Moldavansky edge group has rank 2.

References

- [AN] M. Abert and N. Nikolov “Rank gradient, cost of groups and the rank versus Heegard genus problem”, pre-print.
- [BZ] M. Boileau and H. Zieschang, “Heegaard genus of closed orientable Seifert 3-manifolds”, *Invent. Math.* **76** (1984), no. 3, 455–468.
- [B] K. S. Brown, “Trees, valuations, and the Bieri-Neumann-Strebel invariant”, *Invent. Math.* **90** (1987), no. 3, 479–504.
- [G] H. B. Griffiths, “Automorphisms of a 3-dimensional handlebody”, *Abh. Math. Sem. Univ. Hamburg* **26** (1963/1964) 191–210.
- [J] W. Jaco, “Roots, relations and centralizers in three-manifold groups”, *Geometric topology (Proc. Conf., Park City, Utah, 1974)*, pp. 283–309. Lecture Notes in Math., Vol. 438, Springer, Berlin, 1975.
- [K] I. Kapovich, “Howson property and one-relator groups”, *Comm. Algebra* **27**, no. 3 (1999), 1057–1072.
- [L] R. C. Lyndon, Cohomology theory of groups with a single defining relation, *Ann. of Math. (2)* **52** (1950), 650–665.
- [LS] R. C. Lyndon and P. E. Schupp, *Combinatorial Group Theory*, Springer-Verlag, 1977.
- [MS] J. McCool and P. Schupp, “On one relator groups and HNN extensions”, *J. Austral. Math. Soc.* **16** (1973), 249–256.
- [Sc] Scott, G. P. Compact submanifolds of 3-manifolds, *J. London Math. Soc. (2)* **7** (1973), 246–250.
- [S] J. Stallings, “On fibering certain 3-manifolds”, *Topology of 3-manifolds and related topics*, Prentice Hall, 1961.
- [Z] H. Zieschang, “On Heegaard diagrams of 3-manifolds”, *Asterisque* **163-164** (1988), 247–280.

Math Department
SUNY at Buffalo
jdmaster@buffalo.edu