Self-complementary plane partitions by Proctor's minuscule method

Greg Kuperberg*
Department of Mathematics, University of Chicago, Chicago, IL 95616†

A method of Proctor [4] realizes the set of arbitrary plane partitions in a box and the set of symmetric plane partitions as bases of linear representations of Lie groups. We extend this method by realizing transposition and complementation of plane partitions as natural linear transformations of the representations, thereby enumerating symmetric plane partitions, self-complementary plane partitions, and transpose-complement plane partitions in a new way.

arXiv:math/9411239v1 [math.CO] 1 Nov 1994

While investigating minuscule representations of Lie algebras, Robert Proctor [4] developed one of the simplest methods for enumerating symmetry classes of plane partitions. The method yields an enumeration of arbitrary plane partitions, using the representation theory of $\mathrm{SL}(n)$, and symmetric plane partitions, using the (projective) representation theory of $\mathrm{SO}(2n+1)$. (Proctor also provided solutions to two previously open symmetry classes, namely TCPP's and SSCPP's in the notation of Stembridge, in a related representation-theoretic investigation.)

The author and John Stembridge have independently discovered a way to extend this method to enumerate self-complementary plane partitions (first enumerated by Stanley [6]). In this paper, we will present a brief proof of this enumeration; Stembridge [7] gives a longer exposition and another generalization of the new proof. For both of us, the proof was inspired by, and partially explains, Stembridge's $q=-1$ phenomenon [8], which in this case is the observation that the number of self-complementary plane partitions is the number of plane partitions with an even number of cubes minus the number with an odd number of cubes.

In addition, the author has found a new proof of the enumeration of symmetric plane partitions which extends to an enumeration of TCPP's, while Stembridge's analysis also enumerates SSCPP's and self-complementary chains of order ideals in other minuscule posets. Unfortunately, symmetry classes involving cyclic symmetry have no known treatment by the minuscule method. Table I gives a summary of which enumerations are given in which paper.

No. [6]	Case	Directly treated by... This paper	Stembridge [7]	Proctor [4]
1	P	Yes	Yes	Yes
2	S	Yes	Yes	Yes
3	CS	No	No	No
4	TS	No	No	No
5	SC	Yes	Yes	No
6	TC	Yes	No	No
7	SSC	No	Yes	No
8	CSTC	No	No	No
9	CSSC	No	No	No
10	TSSC	No	No	No

Table I: Proctor's minuscule method and its extensions

1. CONVENTIONS




Figure 1: A plane partition in a box

A plane partition in an $a \times b \times c$ box is a collection of unit cubes in the rectangular solid $[0,a] \times [0,b] \times [0,c]$ in three dimensions which is stable under gravitational attraction towards the origin; Figure 1 shows an example of a plane partition.

The two simplest symmetry operations on the set of plane partitions are transposition τ, which consists of switching two coordinate axes; and rotation ρ, which consists of cyclically permuting all three axes. There is also a third operation, complementation κ, which consists of taking all cubes which are in the box but not in the diagram of the partition and reversing all three coordinates of the box.

A partition in an $a \times b$ rectangle is the analogue of a plane partition in two dimensions, as shown in Figure 2. The operations of complementation and transposition act on partitions. The set of partitions has a partial ordering given by set-theoretic inclusion, and it will be convenient to think of a plane partition as an ordered chain of c partitions. (Adjacent terms may be equal in such a chain.) Transposition of each partition in such a chain is equivalent to transposition of the plane partition, and the same is true of complementation provided the chain is also reversed.

*Supported by NSF Postdoctoral Fellowship DMS #9107908
†Current email: greg@math.ucdavis.edu




Figure 2: A partition in a rectangle

A partition P in an $a \times b$ rectangle can also be viewed as a a binary sequence, i.e., a function b_P from $\{1,\ldots,a+b\}$ to $\{0,1\}$ whose values sum to a. Specifically, the boundary of P is a path of $a+b$ steps from the upper left corner to the lower right corner, and we let $b_P(n) = 1$ if the n'th step goes down and $b_P(n) = 0$ if it goes to the right. Transposing or complementing a partitions has a simple effect on its binary sequence:

$$\begin{aligned} b_{\kappa(P)}(n) &= b_P(a+b+1-n) \\ b_{\kappa\tau(P)}(n) &= 1-b_P(n). \end{aligned}$$

Finally, if T is a plane partition (or a partition), $|T|$ is the number of cubes (squares) in T, and the q-weight of T is defined to be $q^{|T|}$.

2. ARBITRARY PLANE PARTITIONS

This section is a review of the following result:

Theorem 2.1. *The set of plane partitions in an $a \times b \times c$ box can be regarded as a basis of an irreducible representation of* $\mathrm{SL}(a+b)$.

This result is essentially due to Young, Schur, and Weyl, but it was first noted in the context of plane partitions by Stanley and Proctor, and Proctor generalized it to symmetric plane partitions and $\mathrm{SO}(2a+1)$.

Consider a vector space V over $\mathbb{C}$ (some other choices of the ground field work also) of dimension $a+b$ and with basis $x_1,\ldots,x_{a+b}$. The vector space $\Lambda^a V$ of alternating forms over V of degree a has a basis indexed by partitions in an $a \times b$ box. Namely, the basis element corresponding to P is a wedge product formed from the binary sequence of P:

$$x_P = \bigwedge_{n=1}^{a+b} x_n^{b_P(n)}.$$

Consider the vector space $S^c(\Lambda^a V)$ of symmetric forms over $\Lambda^a V$ of degree c, i.e., homogeneous polynomials of degree c in the x_P's, treated as independent variables. If a plane partition T is viewed as a chain of partitions $P_1 \geq P_2 \geq \ldots \geq P_c$, then we can define an element p_T of $S^c(\Lambda^a V)$ by

$$p_T = \prod_n x_{P_n}.$$

These elements are not a basis of $S^c(\Lambda^a V)$, but merely a linearly independent set.

We view V, $\Lambda^a V$, and $S^c(\Lambda^a V)$ as linear representations of $\mathrm{SL}(a+b) = \mathrm{SL}(V)$. The representation $S^c(\Lambda^a V)$ has a unique highest weight vector with weight $c\lambda_a$. Here λ_n is the nth fundamental weight, the weight corresponding to the nth vertex from the end of the Dynkin diagram of $\mathrm{SL}(a+b)$, which is a chain of length $a+b-1$. Therefore, the irreducible representation $V(c\lambda_a)$ appears as a subrepresentation with multiplicity 1, and there is an equivariant projection

$$\pi : S^c(\Lambda^a V) \to V(c\lambda_a).$$

(Those not familiar with the representation theory of simple Lie algebras should refer to Serre [5] or Bourbaki [1].)

We now invoke the following fundamental result:

Theorem 2.2. *The vectors $x_T = \pi(p_T)$, ranging over all plane partitions T in an $a \times b \times c$ box, are a basis for the irreducible representation $V(c\lambda_a)$ of* $\mathrm{SL}(a+b)$.

The proof follows from the theory of semistandard tableaux, originally due to Young, as applied by Weyl to Lie groups [3].

Using this result, the Weyl dimension formula gives us the number of plane partitions in a box. In the notation of Propp, the number is:

$$N(a,b,c) = \frac{H(a+b+c)H(a)H(b)H(c)}{H(a+b)H(a+c)H(b+c)}$$

Here $H(n) = 1!2!3!\ldots(n-1)!$ is the hyperfactorial function. More generally, the Weyl q-dimension formula gives the trace of the element

$$D_q = q^{(1-a-b)/2} \begin{pmatrix} 1 & 0 & 0 & & 0 \\ 0 & q & 0 & \cdots & 0 \\ 0 & 0 & q^2 & & 0 \\ \vdots & & & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & q^{a+b-1} \end{pmatrix}$$

in $\mathrm{SL}(a+b)$ in its action on $V(c\lambda_a)$ (or any other irreducible representation), which is important to us because of the following lemma:

Lemma 2.3. *The the action of $D_q \in \mathrm{SL}(a+b)$ on $V(c\lambda_a)$ is diagonal in the basis $\{x_T\}$, and the eigenvalue of x_T is $q^{|T|-abc/2}$.*

Proof. We will work our way up from x_n for an integer n to x_P for a partition P to p_T and then to x_T. The element D_q is, by definition, diagonal in its action on V, and the eigenvalue of x_n is $q^{n-(a+b+1)/2}$.

Since x_P is a (wedge) product of x_n's, it is also an eigenvector of D_q, and its eigenvalue is the product of the eigenvalues. If P_0 is the empty partition, then

$$x_{P_0} = x_1 \wedge x_2 \wedge \ldots\ldots x_a,$$

and the eigenvalue is therefore $q^{a(a+1)/2-a(a+b+1)/2} = q^{-ab/2}$. Consider the effect of an elementary move on P which consists of adding a single square to it. This is equivalent to replacing some factor x_n in x_P by x_{n+1}, which results in an extra factor of q. Therefore, the eigenvalue is $q^{|P|-ab/2}$ in the general case.

Since p_T is a (symmetric) product of c different x_P's, it is also an eigenvector of D_q. Its eigenvalue is the product of the eigenvalues of its factors, namely $q^{|T|-abc/2}$.

Finally, D_q commutes with the projection π (because π is equivariant under the action of all of $\mathrm{SL}(a+b)$). Thus, if p_T is an eigenvector with a certain eigenvalue, $x_T = \pi(p_T)$ must be an eigenvector with the same eigenvalue. □

The logic of the proof of Lemma 2.3 will be repeated several more times in the rest of the paper with successively shorter explanations.

The q-dimension formula yields the trace of D_q, which by the lemma is $q^{-abc/2}$ times the total q-weight of all plane partitions. The formula for the q-weight is:

$$N(a,b,c)_q = \frac{H(a+b+c)_q H(a)_q H(b)_q H(c)_q}{H(a+b)_q H(a+c)_q H(b+c)_q},$$

where

$$H(n)_q = (1)_q!(2)_q!\ldots(n-1)_q!$$

is the q-hyperfactorial function,

$$(n)_q! = (1)_q(2)_q\ldots(n)_q$$

is the q-factorial function, and

$$(n)_q = 1+q+q^2+\ldots+q^{n-1}$$

is a q-integer.

3. SELF-COMPLEMENTARY PLANE PARTITIONS

Theorem 3.1 (Stembridge, K). *If the plane partitions in an $a\times b\times c$ box are a basis for a vector space, the operation of complementation is linearly conjugate to the operation of negating partitions with an odd number of cubes.*

We consider the effect of the group element of the form:

$$K = i^{a+b-1}\begin{pmatrix} 0&0&0&0&1\\ 0&0&0&1&0\\ 0&0&1&0&0\\ 0&1&0&0&0\\ 1&0&0&0&0 \end{pmatrix}$$

in $\mathrm{SL}(a+b)$ on the x_P's and the x_T's. By definition, it sends x_n to $i^{a+b-1}x_{a+b+1-n}$, and its effect on a partition is to reverse its binary sequence, which is the complementation operation. Specifically:

$$K(x_P) = i^{a(a+b-1)}(-1)^{a(a-1)/2}x_{\kappa(P)} = i^{ab}x_{\kappa(P)}.$$

The factor of $(-1)^{a(a-1)/2}$ comes from a reversal of the factors in x_P, which is an antisymmetric product. Since p_T is a symmetric product of x_P's, the effect of K is:

$$K(p_T) = i^{abc}p_{\kappa(T)}$$

Finally, since K commutes with π, it has the same effect on the x_T's. Therefore the trace of K in $V(c\lambda_a)$ is, up to a factor of i^{abc}, the number of plane partitions fixed by κ. $V(b\lambda_a)$.

But K is conjugate to D_{-1} and therefore has the same trace. (Note that the notation D_q is an abuse because the matrix is really a function of $q^{1/2}$. To obtain D_{-1}, we set $q^{1/2} = i$.) In conclusion,

Corollary 3.2 (Stanley).

$$N_\kappa(a,b,c) = N(a,b,c)_{-1}$$

This equation is easily equivalent to the formula given by Stanley [6].

4. SYMMETRIC PLANE PARTITIONS

Theorem 4.1. *If the plane partitions in an $a\times a\times c$ box are interpreted as a basis for an irreducible representation of $\mathrm{SL}(2a)$ as in section 2, transposition is induced by the Hodge star operator relative to a symplectic inner product.*

Proof. We proceed with the notation of the section 2, taking $a=b$. For any non-degenerate bilinear form B on V, the group $\mathrm{SL}(V)$ possesses an outer automorphism $\sigma_B : A \mapsto A^{-T}$ (inverse transpose), where the transpose $A\mapsto A^T$ is taken relative to the form B. (Usually, B is either symmetric or antisymmetric, but it need not be either in the definition of σ_B.) Explicitly, B induces an isomorphism $\alpha_B : V \to V^*$, where V^* is the dual vector space, given by

$$\alpha_B(v) = B(\cdot, v).$$

Then $\sigma_B(A)$ is the composition

$$\alpha_B^{-1}A^{*-1}\alpha_B,$$

where A^* is the adjoint of A. Even more explicitly, if A and B are given by matrices M_A and M_B,

$$M_{\sigma_B(A)} = M_B^{-1}M_A^{-T}M_B,$$

where M^{-T} is the usual inverse transpose of the matrix M. We consider the group $\mathbb{Z}/2\ltimes_{\sigma_B}\mathrm{SL}(V)$, which is the same group for all choices of B because σ_B differs from $\sigma_{B'}$ by an inner automorphism of $\mathrm{SL}(V)$.

The automorphism σ_B can be given a compatible linear action on $\Lambda^a V$, called the Hodge star operator, as follows: First, we pick a specific volume form on V, i. e. an isomorphism between $\Lambda^{2a}V$ and $\mathbb{C}$, relative to which B has determinant 1. This identification induces a bilinear form on $\Lambda^a V$ given by the wedge product. There is a second bilinear form induced

by B (and denoted by the same letter), because $\Lambda^a V \subseteq V^{\otimes a}$ and $V^{\otimes a}$ has a bilinear form which is just a multilinear extension of B on V. More explicitly, if $v_1,\ldots,v_a$ and $w_1,\ldots,w_a$ are two sequences of vectors,

$$B(v_1\wedge\ldots\wedge v_a, w_1\wedge\ldots\wedge w_a) = \sum_{\sigma\in S_a}(-1)^\sigma \prod_{n=1}^{a} B(v_n, w_{\sigma(n)}) \tag{1}$$

Both bilinear operations can be interpreted as functions from $\Lambda^a V$ to its dual, and the Hodge star operator is either one composed with the inverse of the other. More explicitly, the Hodge star is defined by the equation

$$B(\nu, *\omega) = \nu\wedge\omega$$

for $\nu,\omega\in\Lambda^a V$. If it is assigned to σ_B, $\Lambda^a V$ becomes a representation of $\mathbb{Z}/2\ltimes_{\sigma_B}\mathrm{SL}(V)$.

With respect to the given basis for V, we define volume in the standard way, and we define B by a matrix of the form:

$$\begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 1\\ 0 & 0 & 0 & 0 & -1 & 0\\ 0 & 0 & 0 & 1 & 0 & 0\\ 0 & 0 & -1 & 0 & 0 & 0\\ 0 & 1 & 0 & 0 & 0 & 0\\ -1 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}.$$

In general, B is defined by the equations

$$B(x_n, x_k) = \begin{cases} (-1)^n & k = 2a+1-n\\ 0 & \text{otherwise}\end{cases}. \tag{2}$$

With this choice of B,

$$B(x_P, x_{P'}) = 0$$

unless $P' = \kappa(P)$, and

$$B(x_P, x_{\kappa(P)}) = (-1)^{|P|},$$

by multilinearity. The sign in this equation merits some explanation. A factor of $(-1)^{|P|+a(a-1)/2}$ comes from multiplying a factors of equation 2, while another factor of $(-1)^{a(a-1)/2}$ appears because in the sole non-zero term in equation 1, σ is the reversing permutation.

On the other hand,

$$x_P\wedge x_{P'} = 0$$

unless $b_P(n) = 1 - b_{P'}(n)$, because otherwise the wedge product has repeated linear factors. This happens precisely when $P' = \kappa\tau(P)$. In this case:

$$x_P\wedge x_{\kappa\tau(P)} = (-1)^{|P|} x_1\wedge\ldots\wedge x_{2a}.$$

The sign comes from permuting the wedge factors. Composing the wedge product (as a scalar-valued bilinear function) with B, we obtain

$$*x_P = x_{\tau(P)},$$

which establishes that the Hodge star operation as the linear extension of transposition of partitions. We can extend the Hodge star multilinearly to the polynomial space $S^c(\Lambda^a V)$ over a-forms, and it immediately follows that

$$*\, p_T = p_{\tau(T)}. \tag{3}$$

We would like to infer the conclusion of the theorem,

$$*x_T = x_{\tau(T)},$$

from equation 3. To do so, we need to know that the Hodge star operator commutes with the projection $\pi : S^c(\Lambda^a V)\to V(c\lambda_a)$. In general, given a representation R of a semidirect product $\mathbb{Z}/2\ltimes_\sigma G$, with G reductive, either σ, acting on G, fixes the character of a G-irreducible summand V of R, in which case σ, acting on R, leaves V invariant; or σ does not fix the character of V, in which case $\sigma(V)$ is a disjoint G-irreducible summand of R. In our case, $V(c\lambda_a)$ appears with multiplicity 1, and π is characterized as the projection whose image is $V(c\lambda_a)$ and whose kernel is the direct sum of all other $SL(V)$-irreducible summands. With the given choice of B, the automorphism σ_B is induced by reversing the Dynkin diagram of $SL(V)$, which fixes its middle fundamental weight λ_a and therefore also fixes $c\lambda_a$ and the entire character of $V(c\lambda_a)$. Therefore both the kernel and image of π are invariant under the Hodge star action of σ_B, which implies that π and $*$ commute. □

As stated in the proof, σ_B is a Dynkin diagram automorphism. The character theory of semi-direct products arising from Dynkin diagram automorphisms is described by Neil Chriss [2], who explained to the author that although this theory is known to several representation theorists, it may not have been previously published. The group $\mathbb{Z}/2\ltimes_{\sigma_B}\mathrm{SL}(2a)$ has two components. The character of a representation on the identity component is just the usual character of $\mathrm{SL}(2a)$. The character on the σ_B component, when non-zero, equals the character of an associated representation of the dual Lie group, in this case $\mathrm{SO}(2a+1)$, to the subgroup fixed by the outer automorphism, in this case $\mathrm{Sp}(2a)$. The representation associated to $V_{\mathrm{SL}(2a)}(c\lambda_a)$ is the projective representation $V_{\mathrm{SO}(2a+1)}(c\lambda_a)$, where λ_a is now the weight corresponding to the short root of B_a, the root system of $\mathrm{SO}(2a+1)$. In particular, the trace of σ_B is the dimension of $V_{\mathrm{SO}(2a+1)}(c\lambda_a)$, as given by the Weyl dimension formula, and the trace of $\sigma_B D_q$ is the q-dimension, as given by the Weyl q-dimension formula. From the construction of these linear operators, their traces are $N_\tau(a,a,c)$, the number of τ-invariant plane partitions, and their total q-weight, $N_\tau(a,a,c)_q$, times $q^{-a^2c/2}$, respectively.

By analogy with the results described in section 2, Proctor [4] also realizes τ-invariant plane partitions as a basis for the same projective representation $V_{\mathrm{SO}(2a+1)}(c\lambda_a)$, and the q-dimension again equals the trace of the linear transformation which multiplies each plane partition by its q-weight times $q^{-a^2c/2}$. Since the result of this section arrives at the same Lie group character, if not an explicit representation, it provides another proof which is similar to but not the same as Proctor's

proof of the product formulas for $N_\tau(a,a,c)$ and $N_\tau(a,a,c)_q$. Converting the Weyl formulas by an extension of Propp's notation, they are

$$N_\tau(a,a,c)_q = \frac{H_2(2a+b)_{q^2}H(a)_{q^2}(2a+b-1)_q!!}{H(a+b)_{q^2}H_2(b+1)_{q^2}(b-1)_q!!(2a-1)_q!!}$$

Here

$$H_2(n) = (n-2)_q!(n-4)_q!(n-6)_q!\ldots$$

is a staggered q-hyperfactorial, and

$$n_q!! = (n)_q(n-2)_q(n-4)_q\ldots$$

is a staggered q-factorial. (The formula was first proved by George Andrews.)

5. TRANSPOSE-COMPLEMENT PLANE PARTITIONS

In this section, we combine the results of the previous two sections to enumerate $\kappa\tau$-invariant plane partitions. In the representation $V(c\lambda_a)$ of $\mathbb{Z}/2 \ltimes_{\sigma_B} \mathrm{SL}(2a)$, which has plane partitions in an $a \times a \times c$ box as a basis, the group element σ_B acts as the linear extension of τ, while the group element K acts as i^{a^2c} times the linear extension of κ. Therefore their product acts as i^{a^2c} times $\kappa\tau$. If $\sigma_B K$ were conjugate to $\sigma_B D_{-1}$, it would imply by the results of the last two sections that

$$N_{\tau\kappa}(a,a,c) = N_\tau(a,a,c)_{-1},$$

which easily implies the usual formula due to Proctor and mentioned by Stanley [6]. It is easy to check that a group element with a matrix of the form

$$\frac{1}{\sqrt{2}}\begin{pmatrix} 1 & 0 & 0 & 1 \\ 0 & 1 & -1 & 0 \\ 0 & 1 & 1 & 0 \\ -1 & 0 & 0 & 1 \end{pmatrix}$$

conjugates K to D_{-1} and commutes with σ_B. (The general expression for this element is $\frac{1}{\sqrt{2}}(I + D_{-1}K^{-1})$, or its matrix is $\frac{1}{\sqrt{2}}(I+M_B)$.) Therefore $\sigma_B K$ and $\sigma_B D_{-1}$ are also conjugate. The conclusion is

Theorem 5.1.

$$N_{\kappa\tau}(a,a,c) = N_\tau(a,a,c)_{-1}$$

6. SYMMETRIC, SELF-COMPLEMENTARY PLANE PARTITIONS

In this paper, we have constructed a linear representation $V_{\mathrm{SL}(a+b)}(c\lambda_a)$ with a basis indexed by plane partitions. We have recognized the linear extension of complementation κ as part of the group action on this representation. When $a = b$, we have also recognized the linear extension of transposition τ as part of the group action of a semidirect product of $\mathrm{SL}(a+b)$. The traces of these linear transformations tell us the number of elements fixed by κ or τ or $\kappa\tau$. The author briefly thought that this construction would lead to an enumeration of plane partitions fixed by both κ and τ. this is a naive hope. Consider a permutation representation of the Klein group generated by τ and κ with one free orbit and two fixed points, and consider a second permutation representation with three orbits with two points, with stabilizers τ, κ, and $\kappa\tau$. These two representations are different, but the linear representations they induce are isomorphic.

Thus, in the absence of extra structure, the linear representation $V_{\mathrm{SL}(2a)}(c\lambda_a)$ gives no information about κ,τ-invariant plane partitions. However, Stembridge enumerates κ,τ-invariant plane partitions by expressing κ as an involution on the representation $V_{\mathrm{SO}(2a+1)}(c\lambda_a)$. The author conjectures that latter representation can be realized as a natural vector subspace of the former one.

[1] Nicolas Bourbaki, *Groupes et algébres de Lie*, vol. 4–6, Hermann, Paris, 1968.

[2] Neil Chriss, *Representations of Hecke algebras arising from unramified groups*, Ph.D. thesis, University of Chicago, 1993.

[3] Gordon James and Adalbert Kerber, *The representation theory of the symmetric group*, Addison-Wesley, Reading, Mass., 1981.

[4] Robert A. Proctor, *Bruhat lattices, plane partition generating functions, and minuscule representations*, Europ. J. Combin. **5** (1984), no. 4, 331–350.

[5] Jean-Pierre Serre, *Complex semismiple Lie algebras*, Springer-Verlag, New York, 1987.

[6] Richard P. Stanley, *Symmetries of plane partitions*, J. Combin. Theory Ser. A **43** (1986), no. 1, 103–113.

[7] John R. Stembridge, *On minuscule representations, plane partitions, and involutions in complex Lie groups*, Duke Math. J. **73** (1994), no. 2, 469–490.

[8] ———, *Some hidden relations involving the ten symmetry classes of plane partitions*, J. Combin. Theory Ser. A **68** (1994), 372–409.